

Full wwPDB X-ray Structure Validation Report (i)

Aug 15, 2023 – 02:01 PM EDT

PDB ID : 1U1Y

Title : Crystal structure of a complex between WT bacteriophage MS2 coat protein

and an F5 aptamer RNA stemloop with 2aminopurine substituted at the-10

position

Authors: Horn, W.T.; Convery, M.A.; Stonehouse, N.J.; Adams, C.J.; Liljas, L.;

Phillips, S.E.; Stockley, P.G.

Deposited on : 2004-07-16

Resolution : 2.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
11126112	(# Entries)	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
Clashscore	141614	3438 (2.90-2.82)
Ramachandran outliers	138981	3348 (2.90-2.82)
Sidechain outliers	138945	3351 (2.90-2.82)
RNA backbone	3102	1088 (3.12-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain					
1	R	17		71%		29%		
1	S	17	29%	35%	24%	6%	6%	
2	A	129		80%		17%	•	
2	В	129		81%		16%	•	
2	С	129		84%		12%	•	



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3755 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a RNA chain called 5'-R(*CP*CP*GP*GP*(2PR)P*GP*GP*AP*UP*CP*AP*CP*AP*CP*GP*G)-3'.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	D	17	Total	С	N	О	Р	0	0	0
1	11	17	359	162	70	111	16	U	0	0
1	C	16	Total	С	N	О	Р	0	0	0
1	S	10	345	154	67	108	16	U	0	U

• Molecule 2 is a protein called Coat protein.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	Λ 190	129	Total	С	N	О	S	0	0	0
	A	129	964	606	165	189	4	0	0	
2	D	129	Total	С	N	О	S	0	0	0
	Б	129	964	606	165	189	4	U	0	
2	С	129	Total	С	N	О	S	0	0	0
	2 C	129	964	606	165	189	4	U	U	U

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	R	15	Total O 15 15	0	0
3	S	12	Total O 12 12	0	0
3	A	38	Total O 38 38	0	0
3	В	51	Total O 51 51	0	0
3	С	43	Total O 43 43	0	0



Residue-property plots (i) 3


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 2: Coat protein

 $\bullet \ \mathrm{Molecule} \ 1: \ 5'-R(^*\mathrm{CP}^*\mathrm{CP}^*\mathrm{GP}^*\mathrm{GP}^*(2\mathrm{PR})\mathrm{P}^*\mathrm{GP}^*\mathrm{GP}^*\mathrm{AP}^*\mathrm{UP}^*\mathrm{CP}^*\mathrm{AP}^*\mathrm{CP}^*\mathrm{CP}^*\mathrm{AP}^*\mathrm{CP}^*\mathrm{GP}$)-3'

Chain R: 71% 29% $\bullet \ \mathrm{Molecule} \ 1: \ 5'-R(^*\mathrm{CP}^*\mathrm{CP}^*\mathrm{GP}^*\mathrm{GP}^*(2\mathrm{PR})\mathrm{P}^*\mathrm{GP}^*\mathrm{GP}^*\mathrm{AP}^*\mathrm{UP}^*\mathrm{CP}^*\mathrm{AP}^*\mathrm{CP}^*\mathrm{CP}^*\mathrm{AP}^*\mathrm{CP}^*\mathrm{GP}$)-3'Chain S: 29% 24% 35% 6% • Molecule 2: Coat protein Chain A: 80% 17% • Molecule 2: Coat protein Chain B: 81% 16%





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants	288.61Å 288.61Å 654.68Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 - 2.85	Depositor
% Data completeness	71.2 (30.00-2.85)	Depositor
(in resolution range)	71.2 (90.00 2.00)	Берозпог
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.188 , 0.196	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3755	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 2PR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	R	0.28	0/376	0.82	$1/581 \ (0.2\%)$	
1	S	0.96	2/361~(0.6%)	2.14	$7/559 \ (1.3\%)$	
2	A	0.33	0/981	0.62	0/1337	
2	В	0.33	0/981	0.64	0/1337	
2	С	0.33	0/981	0.61	0/1337	
All	All	0.43	$2/3680 \ (0.1\%)$	0.93	8/5151 (0.2%)	

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	Ideal(A)
1	S	115	С	C4'-C3'	11.27	1.65	1.53
1	S	115	С	C3'-O3'	6.29	1.50	1.42

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	S	115	С	N1-C1'-C2'	-40.17	61.78	114.00
1	S	115	С	C1'-O4'-C4'	-8.31	103.25	109.90
1	S	115	С	C6-N1-C2	-7.63	117.25	120.30
1	S	115	С	O4'-C1'-C2'	7.17	114.05	107.60
1	S	115	С	C4'-C3'-O3'	6.85	126.70	113.00
1	S	115	С	C5'-C4'-O4'	-6.17	101.70	109.10
1	R	11	A	N9-C1'-C2'	5.50	121.15	114.00
1	S	115	С	O4'-C1'-N1	5.42	112.54	108.20

There are no chirality outliers.

There are no planarity outliers.



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	R	359	0	183	1	0
1	S	345	0	175	19	0
2	A	964	0	964	21	0
2	В	964	0	964	15	0
2	С	964	0	964	13	0
3	A	38	0	0	5	0
3	В	51	0	0	6	0
3	С	43	0	0	8	0
3	R	15	0	0	0	0
3	S	12	0	0	5	0
All	All	3755	0	3250	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:S:115:C:O2'	1:S:116:G:C5'	1.76	1.32
1:S:115:C:O3'	3:S:122:HOH:O	1.63	1.12
1:S:115:C:O2'	1:S:116:G:H5'	1.36	1.11
1:S:115:C:O2'	1:S:116:G:O5'	1.63	1.11
2:C:20:VAL:HG13	2:C:32:TRP:HB3	1.37	1.04
2:B:20:VAL:HG13	2:B:32:TRP:HB3	1.41	1.02
1:S:115:C:HO2'	1:S:116:G:C5'	1.65	1.00
2:A:20:VAL:HG13	2:A:32:TRP:HB3	1.53	0.88
1:S:115:C:C3'	3:S:122:HOH:O	2.24	0.80
1:S:111:A:H2'	3:S:119:HOH:O	1.95	0.65
2:C:20:VAL:CG1	2:C:32:TRP:HB3	2.23	0.62
1:S:115:C:HO2'	1:S:116:G:P	2.19	0.62
2:C:61:LYS:HG2	2:C:87:ASN:HD22	1.65	0.61
2:A:49:ARG:HH12	2:A:57:LYS:HE2	1.65	0.61
2:C:56:ARG:HD2	3:C:161:HOH:O	2.00	0.61
2:C:16:GLY:HA3	3:C:168:HOH:O	2.01	0.60
2:A:56:ARG:HD2	2:B:38:ARG:HD3	1.83	0.60

Continued on next page...



 $Continued\ from\ previous\ page...$

Continuea from prev		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ (\mathring{\rm A})$	overlap (Å)
2:B:27:ASN:HB2	3:B:151:HOH:O	1.99	0.60
2:C:87:ASN:HB3	3:C:149:HOH:O	2.01	0.60
2:B:15:THR:HG23	3:B:166:HOH:O	2.02	0.60
2:A:57:LYS:HE3	2:A:89:GLU:CD	2.24	0.58
2:A:57:LYS:HD3	3:A:163:HOH:O	2.03	0.57
2:A:1:ALA:HB3	3:A:158:HOH:O	2.06	0.56
2:C:49:ARG:HD2	3:C:155:HOH:O	2.06	0.56
1:S:109:U:H5'	1:S:110:C:C5	2.43	0.54
2:A:52:SER:OG	2:A:53:ALA:N	2.41	0.53
1:S:108:A:H2'	1:S:110:C:C5	2.43	0.53
1:S:109:U:C5'	1:S:110:C:H5	2.20	0.53
2:A:33:ILE:HD13	2:A:38:ARG:NH1	2.24	0.53
2:B:20:VAL:HG13	2:B:32:TRP:CB	2.27	0.52
1:S:106:G:H5"	3:S:124:HOH:O	2.08	0.52
2:B:3:ASN:ND2	2:B:3:ASN:H	2.08	0.52
1:S:115:C:O2'	1:S:116:G:P	2.67	0.51
2:A:54:GLN:HB2	3:A:132:HOH:O	2.09	0.51
2:B:1:ALA:O	2:B:2:SER:CB	2.58	0.51
2:C:125:ASN:HA	3:C:158:HOH:O	2.10	0.50
2:A:86:LEU:HD21	2:A:88:MET:CE	2.42	0.50
2:C:56:ARG:NH1	3:C:166:HOH:O	2.44	0.50
2:B:83:ARG:HD2	3:B:165:HOH:O	2.10	0.50
2:A:49:ARG:NH1	2:A:57:LYS:HE2	2.27	0.49
2:A:33:ILE:HD13	2:A:38:ARG:HH11	1.78	0.48
2:B:56:ARG:NE	3:B:141:HOH:O	2.40	0.47
1:S:104:G:C6	1:S:106:G:N7	2.83	0.47
2:A:8:VAL:HA	2:A:19:THR:HA	1.95	0.47
2:A:57:LYS:HE3	2:A:89:GLU:OE1	2.16	0.46
2:B:67:VAL:O	2:B:70:GLN:HB3	2.17	0.45
1:S:112:C:O2'	1:S:113:C:H5'	2.16	0.45
2:C:57:LYS:NZ	3:C:164:HOH:O	2.49	0.45
2:A:103:LEU:HD12	2:A:103:LEU:HA	1.84	0.45
2:B:62:VAL:N	3:B:176:HOH:O	2.51	0.43
2:C:70:GLN:O	2:C:76:GLU:HA	2.19	0.43
1:S:109:U:H5'	1:S:110:C:H5	1.80	0.43
2:A:60:ILE:HB	2:A:88:MET:HG2	2.00	0.43
2:A:56:ARG:HD3	3:A:160:HOH:O	2.18	0.42
2:C:20:VAL:HG13	2:C:32:TRP:CB	2.28	0.42
2:A:61:LYS:HZ3	2:A:87:ASN:HD21	1.67	0.42
1:S:109:U:C5'	1:S:110:C:C5	3.00	0.42
2:B:28:GLY:HA3	3:C:157:HOH:O	2.19	0.41

Continued on next page...



n previous	paae
	n previous

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
2:A:61:LYS:NZ	2:A:87:ASN:HD21	2.19	0.41
2:B:87:ASN:HB2	3:B:164:HOH:O	2.19	0.41
2:B:30:ALA:O	2:B:45:THR:HA	2.20	0.41
1:R:16:G:O2'	1:R:17:G:H5'	2.20	0.41
1:S:112:C:H5'	3:S:121:HOH:O	2.20	0.41
2:C:15:THR:C	2:C:17:ASP:H	2.24	0.40
1:S:109:U:H5"	1:S:110:C:H5	1.87	0.40
2:A:43:LYS:HE2	3:A:138:HOH:O	2.21	0.40
2:B:118:ILE:HB	2:B:119:PRO:HD3	2.04	0.40
2:A:31:GLU:HG2	2:A:32:TRP:N	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
2	A	127/129 (98%)	120 (94%)	7 (6%)	0	100	100
2	В	127/129 (98%)	123 (97%)	3 (2%)	1 (1%)	19	46
2	С	127/129 (98%)	113 (89%)	14 (11%)	0	100	100
All	All	381/387 (98%)	356 (93%)	24 (6%)	1 (0%)	41	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	В	2	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar



resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	hain Analysed Rotameric Outliers		Percentiles		
2	A	106/106 (100%)	98 (92%)	8 (8%)	13	34
2	В	106/106 (100%)	96 (91%)	10 (9%)	8	23
2	С	106/106 (100%)	95 (90%)	11 (10%)	7	19
All	All	318/318 (100%)	289 (91%)	29 (9%)	9	25

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	4	PHE
2	A	8	VAL
2	A	19	THR
2 2 2 2	A	20	VAL
2	A	25	PHE
2	A	48	VAL THR VAL PHE VAL
2	A	52	SER
2 2 2 2 2 2	A A A A A B B B	128 3 4	SER ILE ASN PHE VAL PHE SER
2	В	3	ASN
2	В	4	PHE
2	В	20	VAL
2 2 2 2 2	В	25	PHE
2	В	35	SER
2	В	38	ARG
2	В	48	ARG VAL ARG LEU ASP THR VAL
2	В	56	ARG
2	В	77	LEU
2	В	114	ASP
2	С	77 114 15	THR
2	С	20	VAL
2	С	23	SER
2	С	35	SER
2	C	38	ARG
2	С	48	VAL ARG
2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	B B C C C C C C C C C C C C C C C C C C	49	ARG
2	С	54	GLN ARG
2	C	56	ARG
2	С	99	SER
2	С	102	GLU



Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	87	ASN
2	В	3	ASN
2	В	36	ASN
2	В	70	GLN
2	С	6	GLN
2	С	87	ASN

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	R	14/17 (82%)	1 (7%)	0
1	S	14/17 (82%)	5 (35%)	1 (7%)
All	All	28/34 (82%)	6 (21%)	1 (3%)

All (6) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	R	6	G
1	S	106	G
1	S	109	U
1	S	112	С
1	S	115	С
1	S	116	G

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	S	115	C

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection.



RMSZ is the roo	ot-mean-square o	of all Z	scores of the	bond le	ngths	or angl	ലട)
	o incan square o	n an Δ	BCOLCB OI UIIC	DOMA IC	115 0110	(Or angr	$-\omega_{I}$.

Mol	Tuna	Chain	Dag	Link	Bo	ond leng	ths	В	ond ang	les
MIOI	Type	Chain	Res	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	2PR	S	105	1	17,23,24	0.71	0	16,33,36	1.41	3 (18%)
1	2PR	R	5	1	17,23,24	0.67	0	16,33,36	1.34	2 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	2PR	S	105	1	-	0/3/21/22	0/3/3/3
1	2PR	R	5	1	-	0/3/21/22	0/3/3/3

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	S	105	2PR	C2-N3-C4	-3.61	111.23	115.36
1	R	5	2PR	C2-N3-C4	-3.59	111.26	115.36
1	S	105	2PR	N2-C2-N1	-2.47	114.93	117.44
1	R	5	2PR	N2-C2-N1	-2.40	115.00	117.44
1	S	105	2PR	O3'-C3'-C2'	-2.12	103.33	110.90

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

