



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 23, 2024 – 03:29 PM EDT

PDB ID : 1U5Y
Title : Crystal structure of murine APRIL, pH 8.0
Authors : Wallweber, H.J.; Compaan, D.M.; Starovasnik, M.A.; Hymowitz, S.G.
Deposited on : 2004-07-28
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	140	 9% 72% 24% ..
1	B	140	 14% 74% 21% ..
1	D	140	 11% 69% 19% 5% 7%

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 3220 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tumor necrosis factor ligand superfamily member 13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	137	1080	684	197	194	5	0	0	0
1	B	137	1062	672	191	194	5	0	0	0
1	D	130	997	636	180	177	4	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	102	GLY	-	cloning artifact	UNP Q9D777
A	103	SER	-	cloning artifact	UNP Q9D777
B	102	GLY	-	cloning artifact	UNP Q9D777
B	103	SER	-	cloning artifact	UNP Q9D777
D	102	GLY	-	cloning artifact	UNP Q9D777
D	103	SER	-	cloning artifact	UNP Q9D777


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	23	Total	O	0	0
			23	23		
2	B	30	Total	O	0	0
			30	30		
2	D	28	Total	O	0	0
			28	28		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tumor necrosis factor ligand superfamily member 13



- Molecule 1: Tumor necrosis factor ligand superfamily member 13



- Molecule 1: Tumor necrosis factor ligand superfamily member 13



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	123.34Å 81.17Å 53.59Å 90.00° 111.79° 90.00°	Depositor
Resolution (Å)	30.00 – 2.30 28.63 – 2.30	Depositor EDS
% Data completeness (in resolution range)	94.7 (30.00-2.30) 94.7 (28.63-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.88 (at 2.31Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.201 , 0.268 0.207 , 0.268	Depositor DCC
R_{free} test set	2213 reflections (10.67%)	wwPDB-VP
Wilson B-factor (Å ²)	29.8	Xtrriage
Anisotropy	0.521	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 43.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3220	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/1105	0.88	6/1497 (0.4%)
1	B	0.56	0/1086	0.80	1/1474 (0.1%)
1	D	0.58	0/1018	0.83	5/1380 (0.4%)
All	All	0.56	0/3209	0.84	12/4351 (0.3%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	214	ASP	CB-CG-OD2	8.54	125.98	118.30
1	D	144	ASP	CB-CG-OD2	7.40	124.96	118.30
1	B	214	ASP	CB-CG-OD2	7.27	124.84	118.30
1	D	214	ASP	CB-CG-OD2	6.97	124.58	118.30
1	A	144	ASP	CB-CG-OD2	6.85	124.46	118.30
1	D	123	ASP	CB-CG-OD2	6.63	124.27	118.30
1	A	164	ASP	CB-CG-OD2	6.21	123.89	118.30
1	A	121	ASP	CB-CG-OD2	6.00	123.70	118.30
1	D	150	ASP	CB-CG-OD2	5.91	123.62	118.30
1	D	164	ASP	CB-CG-OD2	5.11	122.90	118.30
1	A	196	ASP	CB-CG-OD2	5.11	122.90	118.30
1	A	194	ASP	CB-CG-OD2	5.00	122.80	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1080	0	1069	9	0
1	B	1062	0	1040	21	0
1	D	997	0	974	15	0
2	A	23	0	0	0	0
2	B	30	0	0	3	0
2	D	28	0	0	1	0
All	All	3220	0	3083	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:145:ILE:HG22	1:B:217:THR:HG22	1.60	0.83
1:A:175:ARG:HD2	1:A:214:ASP:OD1	1.88	0.74
1:B:106:HIS:NE2	2:B:3:HOH:O	2.23	0.72
1:D:115:ASN:C	1:D:115:ASN:HD22	1.95	0.69
1:A:106:HIS:NE2	2:B:3:HOH:O	2.27	0.67
1:B:186:ARG:NH1	1:B:188:ILE:HD11	2.11	0.66
1:D:116:ILE:HG22	1:D:117:THR:H	1.63	0.62
1:A:186:ARG:HD2	1:A:188:ILE:HD11	1.82	0.61
2:B:3:HOH:O	1:D:106:HIS:NE2	2.30	0.60
1:B:145:ILE:CG2	1:B:217:THR:HG22	2.31	0.59
1:B:191:MET:N	1:B:192:PRO:HD3	2.20	0.57
1:B:190:SER:C	1:B:192:PRO:HD3	2.25	0.56
1:D:115:ASN:C	1:D:115:ASN:ND2	2.59	0.55
1:D:169:MET:HB2	2:D:47:HOH:O	2.06	0.55
1:D:175:ARG:HG2	1:D:184:LEU:HD21	1.90	0.53
1:D:220:ILE:HD12	1:D:220:ILE:N	2.24	0.53
1:B:145:ILE:HB	1:B:215:ILE:HG23	1.91	0.52
1:B:145:ILE:HB	1:B:215:ILE:CG2	2.39	0.52
1:D:115:ASN:ND2	1:D:128:MET:HB2	2.25	0.51
1:B:190:SER:HG	1:D:198:ALA:N	2.09	0.51
1:B:159:GLN:HA	1:B:202:CYS:O	2.11	0.49
1:D:156:LEU:HD13	1:D:238:PHE:CE1	2.47	0.49
1:B:147:ARG:NE	1:B:149:TRP:HZ2	2.10	0.47
1:B:167:PHE:O	1:B:169:MET:HG2	2.15	0.47
1:B:156:LEU:HD13	1:B:238:PHE:CE1	2.50	0.46
1:A:114:VAL:HG12	1:A:128:MET:O	2.16	0.46
1:D:157:TYR:CZ	1:D:237:GLY:HA3	2.52	0.45
1:B:168:THR:O	1:B:168:THR:HG23	2.18	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:220:ILE:HD12	1:B:220:ILE:N	2.33	0.43
1:D:113:PRO:HD3	1:D:234:THR:O	2.18	0.43
1:A:170:GLY:HA3	1:A:187:CYS:O	2.19	0.43
1:A:127:VAL:O	1:A:144:ASP:HA	2.19	0.43
1:B:183:THR:HB	1:D:232:HIS:CD2	2.54	0.42
1:B:156:LEU:HD13	1:B:238:PHE:CD1	2.55	0.42
1:A:197:ARG:NE	1:D:167:PHE:HA	2.36	0.41
1:B:128:MET:HA	1:B:144:ASP:HA	2.02	0.41
1:D:170:GLY:HA3	1:D:187:CYS:O	2.20	0.41
1:A:175:ARG:HD3	1:A:210:LEU:HD22	2.02	0.41
1:B:175:ARG:HA	1:B:215:ILE:O	2.20	0.41
1:B:147:ARG:NE	1:B:149:TRP:CZ2	2.89	0.41
1:A:220:ILE:N	1:A:220:ILE:HD12	2.35	0.41
1:B:191:MET:N	1:B:192:PRO:CD	2.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	135/140 (96%)	131 (97%)	2 (2%)	2 (2%)	10	10
1	B	135/140 (96%)	125 (93%)	8 (6%)	2 (2%)	10	10
1	D	126/140 (90%)	117 (93%)	6 (5%)	3 (2%)	6	4
All	All	396/420 (94%)	373 (94%)	16 (4%)	7 (2%)	8	7

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	167	PHE
1	B	192	PRO
1	D	167	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	120	ALA
1	D	119	LYS
1	B	166	THR
1	D	165	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/122 (97%)	101 (86%)	17 (14%)	3	3
1	B	115/122 (94%)	105 (91%)	10 (9%)	10	12
1	D	104/122 (85%)	92 (88%)	12 (12%)	5	6
All	All	337/366 (92%)	298 (88%)	39 (12%)	5	6

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	105	LYS
1	A	107	SER
1	A	116	ILE
1	A	118	SER
1	A	122	SER
1	A	137	ARG
1	A	145	ILE
1	A	156	LEU
1	A	166	THR
1	A	174	SER
1	A	180	ARG
1	A	181	ARG
1	A	186	ARG
1	A	189	ARG
1	A	190	SER
1	A	222	ARG
1	A	232	HIS
1	B	108	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	114	VAL
1	B	116	ILE
1	B	123	ASP
1	B	140	GLU
1	B	142	GLN
1	B	145	ILE
1	B	174	SER
1	B	193	SER
1	B	202	CYS
1	D	108	VAL
1	D	114	VAL
1	D	115	ASN
1	D	118	SER
1	D	123	ASP
1	D	128	MET
1	D	164	ASP
1	D	166	THR
1	D	174	SER
1	D	175	ARG
1	D	188	ILE
1	D	232	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	142	GLN
1	D	115	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	137/140 (97%)	0.50	13 (9%) 8 11	6, 13, 27, 38	0
1	B	137/140 (97%)	0.71	19 (13%) 2 4	2, 12, 29, 39	0
1	D	130/140 (92%)	0.58	15 (11%) 4 7	4, 12, 32, 39	0
All	All	404/420 (96%)	0.60	47 (11%) 4 6	2, 12, 29, 39	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	120	ALA	6.8
1	B	120	ALA	6.8
1	D	123	ASP	6.5
1	B	106	HIS	5.7
1	D	105	LYS	5.5
1	D	122	SER	5.4
1	D	121	ASP	5.2
1	D	190	SER	5.0
1	D	167	PHE	5.0
1	B	105	LYS	4.8
1	B	122	SER	4.6
1	B	121	ASP	4.6
1	A	105	LYS	4.5
1	B	192	PRO	4.5
1	B	191	MET	4.4
1	A	196	ASP	4.2
1	A	106	HIS	4.2
1	B	149	TRP	4.2
1	B	166	THR	3.9
1	D	120	ALA	3.8
1	A	197	ARG	3.8
1	A	121	ASP	3.7
1	D	178	GLN	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	199	TYR	3.5
1	B	193	SER	3.2
1	D	117	THR	3.2
1	A	122	SER	3.1
1	D	179	GLY	3.1
1	B	165	VAL	2.9
1	A	179	GLY	2.9
1	D	223	ALA	2.9
1	D	119	LYS	2.6
1	B	118	SER	2.6
1	D	106	HIS	2.5
1	A	137	ARG	2.5
1	B	167	PHE	2.5
1	B	124	VAL	2.4
1	D	199	TYR	2.4
1	B	168	THR	2.3
1	A	195	PRO	2.3
1	B	179	GLY	2.3
1	D	169	MET	2.3
1	B	178	GLN	2.2
1	A	205	ALA	2.2
1	A	203	TYR	2.1
1	B	180	ARG	2.0
1	B	181	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.