

Full wwPDB X-ray Structure Validation Report (i)

Oct 18, 2023 – 05:28 PM EDT

PDB ID : 1UAD

Title: Crystal structure of the RalA-GppNHp-Sec5 Ral-binding domain complex

Authors: Fukai, S.; Matern, H.T.; Scheller, R.H.; Brunger, A.T.

Deposited on : 2003-03-09

Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{\rm A})}) \end{array}$
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	A	175	6% 85%		13%	, ··
1	В	175	77%		19%	
2	С	99	5% 69%	21%	•	7%
2	D	99	62%	30%		7%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 4420 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Ras-related protein Ral-A.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	173	Total			0	S	43	0	0
			1389	870	234	279	6			
1	B	173	Total	С	N	О	S	29	0	0
1	Ъ	175	1389	870	234	279	6	29	U	U

• Molecule 2 is a protein called Exocyst complex component Sec 5.


Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
9	C	92	Total	С	N	О	S	0	0	0
2		92	677	424	121	128	4	U	U	U
9	D	92	Total	С	N	О	S	0	0	0
2	D	92	677	424	121	128	4	U	0	U

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	В	1	Total Mg 1 1	0	0

• Molecule 4 is PHOSPHOAMINOPHOSPHONIC ACID-GUANYLATE ESTER (three-letter code: GNP) (formula: C₁₀H₁₇N₆O₁₃P₃).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
4	Λ	1	Total	С	N	О	Р	0	0	
4	A	1	32	10	6	13	3	U	0	
4	D	1	Total	С	N	О	Р	0	0	
4	Б	1	32	10	6	13	3	U	0	

• Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	78	Total O 78 78	0	0
5	В	83	Total O 83 83	0	0
5	С	48	Total O 48 48	0	0
5	D	13	Total O 13 13	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Ras-related protein Ral-A





4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 4	Depositor
Cell constants	117.42Å 117.42Å 102.68Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.51 - 2.10	Depositor
rtesolution (A)	41.51 - 2.10	EDS
% Data completeness	99.3 (41.51-2.10)	Depositor
(in resolution range)	99.4 (41.51-2.10)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$< I/\sigma(I) > 1$	3.86 (at 2.10Å)	Xtriage
Refinement program	CNS 1.1	Depositor
Ρ. Р.	0.219 , 0.254	Depositor
R, R_{free}	0.207 , 0.238	DCC
R_{free} test set	4051 reflections $(10.04%)$	wwPDB-VP
Wilson B-factor (Å ²)	35.5	Xtriage
Anisotropy	0.377	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.32 , 45.3	EDS
L-test for twinning ²	$< L > = 0.48, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	0.034 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4420	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.02% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GNP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.36	0/1409	0.57	0/1893	
1	В	0.34	0/1409	0.55	0/1893	
2	С	0.32	0/688	0.61	0/933	
2	D	0.27	0/688	0.59	0/933	
All	All	0.33	0/4194	0.57	0/5652	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1389	0	1349	14	0
1	В	1389	0	1350	26	0
2	С	677	0	707	22	0
2	D	677	0	707	26	0
3	A	1	0	0	0	0
3	В	1	0	0	0	0
4	A	32	0	13	0	0
4	В	32	0	13	1	0
5	A	78	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	В	83	0	0	2	0
5	С	48	0	0	1	0
5	D	13	0	0	2	0
All	All	4420	0	4139	87	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 11.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1 2:D:78:THR:HG23 1:B:80:ASP:HB3 2:C:41:LEU:HD23	2:D:81:GLY:H 1:B:84:ARG:HH12 2:C:52:ALA:HB2	1.32 1.51	overlap (Å) 0.94
1:B:80:ASP:HB3 2:C:41:LEU:HD23	1:B:84:ARG:HH12 2:C:52:ALA:HB2		
2:C:41:LEU:HD23	2:C:52:ALA:HB2	1.51	
			0.75
O CLOO A CREATER	- 0	1.74	0.69
2:C:69:ASN:ND2	2:C:71:LYS:HE2	2.10	0.67
1:B:112:LEU:HD22	1:B:118:GLU:HA	1.76	0.66
2:D:75:ILE:HG21	2:D:83:ARG:HE	1.61	0.66
2:D:78:THR:HG21	5:D:104:HOH:O	1.96	0.65
2:C:42:THR:HG22	5:C:108:HOH:O	1.97	0.64
2:D:53:GLU:HG2	2:D:55:MET:HE3	1.78	0.64
2:C:69:ASN:HD22	2:C:71:LYS:HE2	1.64	0.62
2:D:66:GLN:HA	2:D:92:LEU:HD21	1.82	0.62
2:D:41:LEU:HD23	2:D:52:ALA:HB2	1.82	0.61
2:D:47:ASN:ND2	2:D:49:LEU:H	1.99	0.61
2:D:44:CYS:SG	2:D:72:GLY:HA2	2.43	0.58
2:C:19:ILE:HD12	2:C:93:LEU:HB2	1.84	0.58
2:D:78:THR:CG2	2:D:81:GLY:H	2.12	0.58
1:B:52:ARG:HD3	1:B:65:ASP:OD2	2.05	0.56
2:C:21:TRP:NE1	2:C:63:ARG:HD3	2.19	0.56
1:A:58:ASP:OD2	1:A:176:ARG:NH2	2.33	0.56
1:B:54:LYS:HD3	1:B:61:GLU:OE1	2.06	0.55
1:B:79:ARG:HH11	1:B:79:ARG:HG3	1.72	0.55
2:D:66:GLN:HB3	5:D:106:HOH:O	2.08	0.52
2:D:4:SER:O	2:D:5:ARG:HD3	2.09	0.52
1:B:107:PHE:O	1:B:111:ILE:HG13	2.10	0.51
1:B:23:GLY:HA2	4:B:1200:GNP:O3G	2.11	0.51
2:D:64:VAL:HG12	2:D:65:GLY:N	2.25	0.51
1:B:80:ASP:O	1:B:84:ARG:HG2	2.11	0.51
1:B:94:SER:OG	1:B:128:LYS:HD2	2.11	0.51
2:D:44:CYS:HB3	2:D:67:ALA:CB	2.41	0.51
1:B:70:ALA:HB1	1:B:72:GLN:HG2	1.94	0.50



 $Continued\ from\ previous\ page...$

Continued from prev		Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \ ({\rm \AA})$	overlap (Å)
2:C:71:LYS:HD3	2:C:71:LYS:H	1.77	0.50
2:D:41:LEU:HD13	2:D:76:VAL:HG22	1.93	0.49
2:D:47:ASN:HD22	2:D:48:CYS:N	2.11	0.49
2:D:21:TRP:N	2:D:64:VAL:O	2.42	0.48
2:D:55:MET:HA	2:D:55:MET:HE2	1.96	0.48
2:C:21:TRP:CD1	2:C:63:ARG:HD3	2.49	0.48
1:A:112:LEU:HD22	1:A:118:GLU:HA	1.96	0.48
2:C:20:PRO:HD3	2:C:93:LEU:O	2.15	0.47
2:C:71:LYS:HD3	2:C:71:LYS:N	2.30	0.47
2:D:36:THR:O	2:D:79:LYS:HE3	2.15	0.46
1:B:79:ARG:HD2	1:B:83:PHE:CZ	2.50	0.46
1:A:79:ARG:HG3	1:A:79:ARG:HH11	1.81	0.46
1:B:110:GLN:O	1:B:114:VAL:HG23	2.15	0.46
2:D:20:PRO:HD3	2:D:93:LEU:O	2.15	0.46
1:A:145:ARG:HD3	1:A:149:TRP:CH2	2.51	0.45
1:B:108:ARG:HD3	1:B:149:TRP:CE2	2.52	0.45
2:C:75:ILE:HD13	2:C:83:ARG:NH1	2.32	0.45
2:D:53:GLU:HG2	2:D:55:MET:CE	2.44	0.45
1:B:22:SER:O	1:B:25:VAL:HG22	2.17	0.45
1:A:132:GLU:OE1	1:A:135:ARG:NE	2.47	0.44
2:D:46:HIS:HB2	2:D:66:GLN:O	2.17	0.44
2:D:44:CYS:HB3	2:D:67:ALA:HB2	1.99	0.44
1:B:143:LYS:HE2	5:B:1282:HOH:O	2.16	0.44
1:A:117:ASP:OD1	1:A:178:ARG:NH2	2.51	0.44
1:B:52:ARG:HD2	1:B:65:ASP:HB2	1.98	0.44
2:D:7:PRO:HA	2:D:8:PRO:HD3	1.94	0.44
1:A:77:ALA:HA	1:A:80:ASP:OD2	2.17	0.44
1:A:79:ARG:HG3	1:A:79:ARG:NH1	2.33	0.44
1:B:145:ARG:HD3	1:B:149:TRP:CH2	2.53	0.43
2:C:19:ILE:HG23	2:C:20:PRO:N	2.33	0.43
1:A:47:LYS:HE2	2:C:15:PRO:CB	2.49	0.43
1:B:40:VAL:HG12	1:B:42:ASP:OD1	2.18	0.43
2:D:47:ASN:HD22	2:D:47:ASN:C	2.20	0.43
1:B:127:ASN:O	1:B:128:LYS:HB2	2.18	0.42
2:C:47:ASN:ND2	2:C:49:LEU:H	2.16	0.42
1:B:63:GLN:NE2	5:B:1264:HOH:O	2.47	0.42
1:B:115:LYS:HA	1:B:115:LYS:HE2	2.00	0.42
1:B:117:ASP:OD2	1:B:119:ASN:N	2.46	0.42
1:B:95:ILE:HD13	1:B:129:SER:HA	2.02	0.42
2:C:64:VAL:HG12	2:C:65:GLY:N	2.34	0.42
2:C:75:ILE:HD13	2:C:83:ARG:HH12	1.85	0.42



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash		
Atom-1	Atom-2	${f distance}({ m \AA})$	overlap (Å) 0.41 0.41 0.41 0.41 0.41 0.41 0.41 0.41 0.41 0.41 0.41 0.41 0.41 0.40		
2:C:69:ASN:HB3	2:C:71:LYS:HE2	2.01	0.41		
2:C:72:GLY:O	2:C:89:SER:HA	2.20	0.41		
1:A:18:ILE:CD1	1:A:86:GLY:HA3	2.50	0.41		
1:A:83:PHE:CB	1:A:114:VAL:HG21	2.51	0.41		
1:A:132:GLU:OE1	1:A:132:GLU:HA	2.21	0.41		
2:C:64:VAL:HG11	2:C:92:LEU:HD13	2.02	0.41		
2:C:70:ASP:OD2	2:C:71:LYS:HD3	2.20	0.41		
1:A:120:VAL:O	1:A:121:PRO:C	2.59	0.41		
1:B:72:GLN:O	1:B:72:GLN:HG3	2.21	0.41		
2:C:94:LYS:HA	2:C:95:PRO:HD3	1.96	0.41		
2:D:13:ILE:O	2:D:13:ILE:HG23	2.21	0.41		
1:A:138:SER:OG	1:A:141:GLU:HB2	2.20	0.40		
1:B:79:ARG:HG3	1:B:79:ARG:NH1	2.36	0.40		
1:B:108:ARG:CZ	1:B:112:LEU:HD11	2.51	0.40		
2:D:37:ASP:OD2	2:D:78:THR:OG1	2.38	0.40		
2:C:67:ALA:H	2:C:92:LEU:CD2	2.35	0.40		

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	171/175 (98%)	162 (95%)	9 (5%)	0	100	100
1	В	171/175 (98%)	164 (96%)	7 (4%)	0	100	100
2	\mathbf{C}	90/99~(91%)	89 (99%)	1 (1%)	0	100	100
2	D	90/99 (91%)	89 (99%)	1 (1%)	0	100	100
All	All	522/548 (95%)	504 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	150/152~(99%)	147 (98%)	3 (2%)	55 60
1	В	150/152~(99%)	142 (95%)	8 (5%)	22 20
2	С	77/83 (93%)	74 (96%)	3 (4%)	32 33
2	D	77/83 (93%)	76 (99%)	1 (1%)	69 75
All	All	454/470 (97%)	439 (97%)	15 (3%)	38 40

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73	GLU
1	A	115	LYS
1	A	141	GLU
1	В	41	GLU
1	В	42	ASP
1	В	43	TYR
1	В	52	ARG
1	В	80	ASP
1	В	119	ASN
1	В	133	ASP
1	В	178	ARG
2	С	19	ILE
2	С	47	ASN
2	С	71	LYS
2	D	47	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	163	ASN
1	В	63	GLN
1	В	72	GLN
1	В	119	ASN



Continued from previous page...

Mol	Chain	Res	Type
2	С	47	ASN
2	D	47	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Ty	True	ype Chain		Link	Bond lengths			Bond angles		
MIOI	Type	Type Chain Res	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GNP	A	200	3	29,34,34	2.23	12 (41%)	33,54,54	2.83	8 (24%)
4	GNP	В	1200	3	29,34,34	2.21	11 (37%)	33,54,54	2.89	8 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GNP	A	200	3	-	4/14/38/38	0/3/3/3



$Continued\ from\ previous\ page...$

\mathbf{Mol}	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GNP	В	1200	3	-	3/14/38/38	0/3/3/3

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$Ideal(\AA)$
4	В	1200	GNP	C2-N2	4.85	1.43	1.33
4	A	200	GNP	C2-N2	4.80	1.43	1.33
4	A	200	GNP	PG-N3B	4.55	1.75	1.63
4	В	1200	GNP	PG-N3B	4.17	1.74	1.63
4	A	200	GNP	C6-N1	3.98	1.40	1.33
4	В	1200	GNP	C6-N1	3.83	1.39	1.33
4	В	1200	GNP	C5-C6	3.22	1.46	1.41
4	A	200	GNP	PG-O1G	3.13	1.51	1.46
4	A	200	GNP	PB-O2B	-2.91	1.48	1.56
4	В	1200	GNP	PG-O3G	-2.89	1.49	1.56
4	В	1200	GNP	PB-O2B	-2.88	1.49	1.56
4	A	200	GNP	O4'-C4'	-2.84	1.38	1.45
4	A	200	GNP	PG-O3G	-2.82	1.49	1.56
4	A	200	GNP	C5-C6	2.80	1.46	1.41
4	В	1200	GNP	PG-O1G	2.64	1.50	1.46
4	В	1200	GNP	O6-C6	-2.52	1.18	1.24
4	В	1200	GNP	O4'-C4'	-2.50	1.39	1.45
4	В	1200	GNP	PB-N3B	2.40	1.69	1.63
4	A	200	GNP	O6-C6	-2.39	1.18	1.24
4	A	200	GNP	PG-O2G	-2.34	1.50	1.56
4	В	1200	GNP	PG-O2G	-2.22	1.50	1.56
4	A	200	GNP	PB-N3B	2.10	1.68	1.63
4	A	200	GNP	C8-N7	-2.06	1.31	1.34

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
4	A	200	GNP	C5-C6-N1	-9.08	111.01	123.43
4	В	1200	GNP	C5-C6-N1	-9.04	111.06	123.43
4	В	1200	GNP	O1G-PG-N3B	-7.25	101.09	111.77
4	A	200	GNP	C2-N3-C4	-7.15	107.19	115.36
4	В	1200	GNP	C2-N3-C4	-7.09	107.26	115.36
4	A	200	GNP	O1G-PG-N3B	-7.02	101.44	111.77
4	В	1200	GNP	N2-C2-N1	-5.12	109.28	117.25
4	A	200	GNP	N2-C2-N1	-4.96	109.54	117.25
4	В	1200	GNP	N3-C2-N1	4.89	133.75	127.22
4	A	200	GNP	N3-C2-N1	4.78	133.61	127.22



Continued from previous page.					
	C'	ontinued	from	previous	page

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$Ideal(^{o})$
4	В	1200	GNP	O2B-PB-O1B	3.63	117.53	109.92
4	A	200	GNP	O2B-PB-O1B	3.44	117.14	109.92
4	В	1200	GNP	O2G-PG-O1G	-2.26	107.76	113.45
4	A	200	GNP	O2G-PG-O1G	-2.10	108.16	113.45
4	В	1200	GNP	C2-N1-C6	2.02	119.14	115.93
4	A	200	GNP	C2-N1-C6	2.02	119.14	115.93

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	200	GNP	PG-N3B-PB-O1B
4	A	200	GNP	PG-N3B-PB-O3A
4	A	200	GNP	PA-O3A-PB-O1B
4	A	200	GNP	PA-O3A-PB-O2B
4	В	1200	GNP	PG-N3B-PB-O1B
4	В	1200	GNP	PG-N3B-PB-O3A
4	В	1200	GNP	PA-O3A-PB-O2B


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	1200	GNP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	168/175 (96%)	0.28	11 (6%) 18 23	23, 37, 71, 86	0
1	В	170/175~(97%)	0.61	9 (5%) 26 32	24, 42, 71, 89	0
2	С	92/99~(92%)	0.34	5 (5%) 25 31	22, 35, 62, 77	0
2	D	92/99 (92%)	0.75	10 (10%) 5 7	34, 53, 78, 86	0
All	All	522/548 (95%)	0.48	35 (6%) 17 22	22, 42, 73, 89	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	11	SER	5.5
2	D	70	ASP	5.1
1	В	82	TYR	4.4
1	A	12	LEU	4.3
2	С	95	PRO	3.9
1	A	183	SER	3.6
1	В	72	GLN	3.6
2	D	71	LYS	3.4
2	D	67	ALA	3.3
2	С	19	ILE	3.3
1	В	43	TYR	3.2
1	A	43	TYR	3.1
1	A	133	ASP	3.1
1	A	78	ILE	3.0
2	D	69	ASN	2.9
1	A	82	TYR	2.8
2	D	68	LYS	2.8
1	A	80	ASP	2.7
2	D	50	LEU	2.6
1	В	78	ILE	2.6
2	D	65	GLY	2.6



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	19	ILE	2.4
2	С	4	SER	2.3
1	В	42	ASP	2.3
1	A	180	MET	2.3
1	A	40	VAL	2.2
2	D	4	SER	2.2
1	В	169	PHE	2.2
1	В	133	ASP	2.2
2	С	94	LYS	2.2
1	В	11	SER	2.2
1	В	18	ILE	2.1
1	A	77	ALA	2.0
2	С	21	TRP	2.0
2	D	75	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B-factors}({f \AA}^2)$	Q<0.9
3	MG	В	300	1/1	0.87	0.24	53,53,53,53	0
3	MG	A	300	1/1	0.95	0.15	38,38,38,38	0
4	GNP	В	1200	32/32	0.96	0.10	40,45,50,52	0
4	GNP	A	200	32/32	0.97	0.09	26,30,36,37	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around GNP B 1200:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around GNP A 200:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

