



wwPDB X-ray Structure Validation Summary Report

Oct 3, 2023 – 06:33 AM EDT

PDB ID : 6UB1
Title : Crystal structure of a GH128 (subgroup III) curdlan-specific exo-beta-1,3-galactanase from *Blastomyces gilchristii* (BgGH128_III) in complex with laminaribiose at -3 and -2 subsites
Authors : Costa, P.A.C.R.; Santos, C.R.; Domingues, M.N.; Lima, E.A.; Mandelli, F.; Murakami, M.T.
Deposited on : 2019-09-11
Resolution : 1.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : **FAILED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8197 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLYCOSIDE HYDROLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	230	1750	1120	295	329	6	0	2	0
1	B	230	1755	1122	297	330	6	0	2	0
1	C	230	1756	1123	296	331	6	0	2	0
1	D	230	1759	1125	296	332	6	0	3	0

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	MET	-	initiating methionine	UNP A0A179UGT5
A	3	GLY	-	expression tag	UNP A0A179UGT5
A	4	SER	-	expression tag	UNP A0A179UGT5
A	5	SER	-	expression tag	UNP A0A179UGT5
A	6	HIS	-	expression tag	UNP A0A179UGT5
A	7	HIS	-	expression tag	UNP A0A179UGT5
A	8	HIS	-	expression tag	UNP A0A179UGT5
A	9	HIS	-	expression tag	UNP A0A179UGT5
A	10	HIS	-	expression tag	UNP A0A179UGT5
A	11	HIS	-	expression tag	UNP A0A179UGT5
A	12	SER	-	expression tag	UNP A0A179UGT5
A	13	SER	-	expression tag	UNP A0A179UGT5
A	14	GLY	-	expression tag	UNP A0A179UGT5
A	15	LEU	-	expression tag	UNP A0A179UGT5
A	16	VAL	-	expression tag	UNP A0A179UGT5
A	17	PRO	-	expression tag	UNP A0A179UGT5
A	18	ALA	-	expression tag	UNP A0A179UGT5
A	19	GLY	-	expression tag	UNP A0A179UGT5
A	20	SER	-	expression tag	UNP A0A179UGT5
A	21	HIS	-	expression tag	UNP A0A179UGT5
A	22	MET	-	expression tag	UNP A0A179UGT5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	2	MET	-	initiating methionine	UNP A0A179UGT5
B	3	GLY	-	expression tag	UNP A0A179UGT5
B	4	SER	-	expression tag	UNP A0A179UGT5
B	5	SER	-	expression tag	UNP A0A179UGT5
B	6	HIS	-	expression tag	UNP A0A179UGT5
B	7	HIS	-	expression tag	UNP A0A179UGT5
B	8	HIS	-	expression tag	UNP A0A179UGT5
B	9	HIS	-	expression tag	UNP A0A179UGT5
B	10	HIS	-	expression tag	UNP A0A179UGT5
B	11	HIS	-	expression tag	UNP A0A179UGT5
B	12	SER	-	expression tag	UNP A0A179UGT5
B	13	SER	-	expression tag	UNP A0A179UGT5
B	14	GLY	-	expression tag	UNP A0A179UGT5
B	15	LEU	-	expression tag	UNP A0A179UGT5
B	16	VAL	-	expression tag	UNP A0A179UGT5
B	17	PRO	-	expression tag	UNP A0A179UGT5
B	18	ALA	-	expression tag	UNP A0A179UGT5
B	19	GLY	-	expression tag	UNP A0A179UGT5
B	20	SER	-	expression tag	UNP A0A179UGT5
B	21	HIS	-	expression tag	UNP A0A179UGT5
B	22	MET	-	expression tag	UNP A0A179UGT5
C	2	MET	-	initiating methionine	UNP A0A179UGT5
C	3	GLY	-	expression tag	UNP A0A179UGT5
C	4	SER	-	expression tag	UNP A0A179UGT5
C	5	SER	-	expression tag	UNP A0A179UGT5
C	6	HIS	-	expression tag	UNP A0A179UGT5
C	7	HIS	-	expression tag	UNP A0A179UGT5
C	8	HIS	-	expression tag	UNP A0A179UGT5
C	9	HIS	-	expression tag	UNP A0A179UGT5
C	10	HIS	-	expression tag	UNP A0A179UGT5
C	11	HIS	-	expression tag	UNP A0A179UGT5
C	12	SER	-	expression tag	UNP A0A179UGT5
C	13	SER	-	expression tag	UNP A0A179UGT5
C	14	GLY	-	expression tag	UNP A0A179UGT5
C	15	LEU	-	expression tag	UNP A0A179UGT5
C	16	VAL	-	expression tag	UNP A0A179UGT5
C	17	PRO	-	expression tag	UNP A0A179UGT5
C	18	ALA	-	expression tag	UNP A0A179UGT5
C	19	GLY	-	expression tag	UNP A0A179UGT5
C	20	SER	-	expression tag	UNP A0A179UGT5
C	21	HIS	-	expression tag	UNP A0A179UGT5
C	22	MET	-	expression tag	UNP A0A179UGT5

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	2	MET	-	initiating methionine	UNP A0A179UGT5
D	3	GLY	-	expression tag	UNP A0A179UGT5
D	4	SER	-	expression tag	UNP A0A179UGT5
D	5	SER	-	expression tag	UNP A0A179UGT5
D	6	HIS	-	expression tag	UNP A0A179UGT5
D	7	HIS	-	expression tag	UNP A0A179UGT5
D	8	HIS	-	expression tag	UNP A0A179UGT5
D	9	HIS	-	expression tag	UNP A0A179UGT5
D	10	HIS	-	expression tag	UNP A0A179UGT5
D	11	HIS	-	expression tag	UNP A0A179UGT5
D	12	SER	-	expression tag	UNP A0A179UGT5
D	13	SER	-	expression tag	UNP A0A179UGT5
D	14	GLY	-	expression tag	UNP A0A179UGT5
D	15	LEU	-	expression tag	UNP A0A179UGT5
D	16	VAL	-	expression tag	UNP A0A179UGT5
D	17	PRO	-	expression tag	UNP A0A179UGT5
D	18	ALA	-	expression tag	UNP A0A179UGT5
D	19	GLY	-	expression tag	UNP A0A179UGT5
D	20	SER	-	expression tag	UNP A0A179UGT5
D	21	HIS	-	expression tag	UNP A0A179UGT5
D	22	MET	-	expression tag	UNP A0A179UGT5

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-3)-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	E	2	Total	C	O	0	0	0
			23	12	11			
2	F	2	Total	C	O	0	0	0
			23	12	11			
2	G	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is beta-D-glucopyranose (three-letter code: BGC) (formula: C₆H₁₂O₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	C O	0	0
			12	6 6		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	263	Total	O	0	0
			263	263		
4	B	273	Total	O	0	0
			273	273		
4	C	266	Total	O	0	0
			266	266		
4	D	294	Total	O	0	0
			294	294		

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics i

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	56.32Å 60.80Å 63.55Å 69.67° 78.24° 77.86°	Depositor
Resolution (Å)	35.49 – 1.60	Depositor
% Data completeness (in resolution range)	86.3 (35.49-1.60)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.60 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.160 , 0.197	Depositor
Wilson B-factor (Å ²)	13.9	Xtrriage
Anisotropy	0.551	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.035 for -h,-l,-k	Xtrriage
Total number of atoms	8197	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 90.48 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.2589e-08. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	E	1	2	12,12,12	1.58	2 (16%)	17,17,17	2.03	5 (29%)
2	BGC	E	2	2	11,11,12	0.62	0	15,15,17	2.24	8 (53%)
2	BGC	F	1	2	12,12,12	1.25	2 (16%)	17,17,17	1.96	7 (41%)
2	BGC	F	2	2	11,11,12	1.13	1 (9%)	15,15,17	2.66	5 (33%)
2	BGC	G	1	2	12,12,12	1.81	2 (16%)	17,17,17	3.19	11 (64%)
2	BGC	G	2	2	11,11,12	0.87	0	15,15,17	1.96	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	E	1	2	-	0/2/22/22	0/1/1/1
2	BGC	E	2	2	-	1/2/19/22	0/1/1/1
2	BGC	F	1	2	-	0/2/22/22	0/1/1/1
2	BGC	F	2	2	-	2/2/19/22	0/1/1/1
2	BGC	G	1	2	-	1/2/22/22	0/1/1/1
2	BGC	G	2	2	-	2/2/19/22	0/1/1/1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	1	BGC	O2-C2	3.94	1.52	1.43
2	E	1	BGC	O1-C1	3.90	1.52	1.39
2	G	1	BGC	O1-C1	3.46	1.50	1.39
2	E	1	BGC	O2-C2	3.00	1.50	1.43
2	F	1	BGC	O2-C2	2.67	1.49	1.43

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	1	BGC	O5-C1-C2	-6.45	98.77	110.28
2	F	2	BGC	O2-C2-C3	5.73	121.62	110.14
2	F	2	BGC	C1-O5-C5	5.41	119.53	112.19
2	G	2	BGC	C2-C3-C4	-4.46	103.17	110.89
2	G	1	BGC	O1-C1-C2	4.35	121.28	109.03

There are no chirality outliers.


5 of 6 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	F	2	BGC	O5-C5-C6-O6
2	F	2	BGC	C4-C5-C6-O6
2	G	2	BGC	O5-C5-C6-O6
2	G	2	BGC	C4-C5-C6-O6
2	E	2	BGC	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





4.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BGC	B	301	-	12,12,12	1.00	0	17,17,17	2.37	7 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BGC	B	301	-	-	2/2/22/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	301	BGC	O4-C4-C3	-5.14	98.47	110.35
3	B	301	BGC	O2-C2-C1	3.23	116.66	109.16
3	B	301	BGC	C1-C2-C3	3.12	116.80	110.31
3	B	301	BGC	O1-C1-O5	-2.97	101.47	110.38
3	B	301	BGC	O5-C1-C2	2.93	115.52	110.28

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	301	BGC	O5-C5-C6-O6
3	B	301	BGC	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



4.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

4.8 Polymer linkage issues

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

5.4 Ligands

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers

EDS failed to run properly - this section is therefore empty.