



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 5, 2023 – 10:08 AM EDT

PDB ID : 3UCL  
Title : Cyclohexanone-bound crystal structure of cyclohexanone monooxygenase in the Rotated conformation  
Authors : Yachnin, B.J.; Berghuis, A.M.  
Deposited on : 2011-10-27  
Resolution : 2.36 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	573	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CYH	A	543	-	-	X	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4215 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclohexanone monooxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	511	3986	2534	672	773	7	0	0	0

There are 33 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-32	MET	-	expression tag	UNP C0STX7
A	-31	GLY	-	expression tag	UNP C0STX7
A	-30	SER	-	expression tag	UNP C0STX7
A	-29	SER	-	expression tag	UNP C0STX7
A	-28	HIS	-	expression tag	UNP C0STX7
A	-27	HIS	-	expression tag	UNP C0STX7
A	-26	HIS	-	expression tag	UNP C0STX7
A	-25	HIS	-	expression tag	UNP C0STX7
A	-24	HIS	-	expression tag	UNP C0STX7
A	-23	HIS	-	expression tag	UNP C0STX7
A	-22	HIS	-	expression tag	UNP C0STX7
A	-21	HIS	-	expression tag	UNP C0STX7
A	-20	ASP	-	expression tag	UNP C0STX7
A	-19	TYR	-	expression tag	UNP C0STX7
A	-18	ASP	-	expression tag	UNP C0STX7
A	-17	ILE	-	expression tag	UNP C0STX7
A	-16	PRO	-	expression tag	UNP C0STX7
A	-15	THR	-	expression tag	UNP C0STX7
A	-14	THR	-	expression tag	UNP C0STX7
A	-13	GLU	-	expression tag	UNP C0STX7
A	-12	ASN	-	expression tag	UNP C0STX7
A	-11	LEU	-	expression tag	UNP C0STX7
A	-10	TYR	-	expression tag	UNP C0STX7
A	-9	PHE	-	expression tag	UNP C0STX7
A	-8	GLN	-	expression tag	UNP C0STX7
A	-7	GLY	-	expression tag	UNP C0STX7
A	-6	SER	-	expression tag	UNP C0STX7

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	LEU	-	expression tag	UNP C0STX7
A	-4	GLU	-	expression tag	UNP C0STX7
A	-3	ALA	-	expression tag	UNP C0STX7
A	-2	SER	-	expression tag	UNP C0STX7
A	-1	MET	-	expression tag	UNP C0STX7
A	0	HIS	-	expression tag	UNP C0STX7

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	53	27	9	15	2	0	0

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula:  $C_{21}H_{28}N_7O_{17}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	48	21	7	17	3	0	0

- Molecule 4 is CYCLOHEXANONE (three-letter code: CYH) (formula: C<sub>6</sub>H<sub>10</sub>O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	7	6	1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	121	Total 121	O 121	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cyclohexanone monooxygenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.54Å 67.11Å 131.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.61 – 2.36 30.61 – 2.36	Depositor EDS
% Data completeness (in resolution range)	97.3 (30.61-2.36) 97.4 (30.61-2.36)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.66 (at 2.36Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.191 , 0.264 0.189 , 0.260	Depositor DCC
$R_{free}$ test set	2061 reflections (10.20%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.8	Xtrriage
Anisotropy	0.185	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 36.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4215	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, FAD, CYH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.80	0/4089	0.78	4/5578 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	265	ARG	NE-CZ-NH1	5.74	123.17	120.30
1	A	265	ARG	NE-CZ-NH2	-5.66	117.47	120.30
1	A	506	LEU	CB-CG-CD1	-5.26	102.05	111.00
1	A	342	ARG	NE-CZ-NH2	-5.07	117.77	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	225	GLU	Peptide
1	A	226	GLN	Peptide
1	A	434	PHE	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3986	0	3751	68	0
2	A	53	0	31	0	0
3	A	48	0	25	8	0
4	A	7	0	10	6	0
5	A	121	0	0	5	0
All	All	4215	0	3817	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:542:NAP:H3D	4:A:543:CYH:C5	1.77	1.14
3:A:542:NAP:H3D	4:A:543:CYH:H51	1.20	1.11
1:A:167:THR:HG21	1:A:189:THR:HG21	1.29	1.10
1:A:428:LEU:H	1:A:444:GLN:HE22	1.01	0.99
1:A:167:THR:CG2	1:A:189:THR:HG21	2.00	0.90
1:A:428:LEU:H	1:A:444:GLN:NE2	1.72	0.87
4:A:543:CYH:H21	5:A:601:HOH:O	1.77	0.84
1:A:435:THR:HG21	1:A:440:SER:OG	1.80	0.82
1:A:270:GLN:HE22	1:A:306:LYS:NZ	1.81	0.79
3:A:542:NAP:H3D	4:A:543:CYH:H52	1.65	0.78
1:A:156:GLY:O	1:A:159:THR:HB	1.87	0.75
1:A:223:ASN:ND2	1:A:226:GLN:CB	2.50	0.74
1:A:428:LEU:N	1:A:444:GLN:HE22	1.84	0.73
1:A:84:ILE:HG12	1:A:89:ILE:HG13	1.73	0.71
1:A:225:GLU:HG3	1:A:227:ILE:HB	1.76	0.68
1:A:59:ASP:OD2	1:A:329:ARG:NH1	2.27	0.68
1:A:435:THR:HG23	1:A:436:ASN:O	1.94	0.67
3:A:542:NAP:C3D	4:A:543:CYH:H51	2.12	0.66
1:A:215:VAL:O	1:A:332:CYS:HA	1.97	0.64
1:A:145:LEU:HD22	1:A:385:VAL:HB	1.80	0.63
1:A:390:ARG:HD3	1:A:403:ASN:ND2	2.13	0.63
1:A:335:GLY:O	1:A:338:GLU:HG2	1.99	0.62

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:270:GLN:HE22	1:A:306:LYS:HZ1	1.45	0.61
1:A:172:GLU:HA	1:A:172:GLU:OE1	2.02	0.60
1:A:188:SER:HB3	4:A:543:CYH:H32	1.84	0.59
1:A:53:TYR:OH	1:A:192:GLN:NE2	2.37	0.58
1:A:487:THR:O	1:A:488:LYS:CB	2.53	0.57
1:A:353:ASN:HD21	1:A:367:ASP:H	1.53	0.56
1:A:390:ARG:HH11	1:A:403:ASN:HD22	1.52	0.56
1:A:15:GLY:O	1:A:20:GLY:HA3	2.06	0.54
1:A:127:THR:HG22	1:A:133:VAL:HG22	1.90	0.53
1:A:145:LEU:HD21	3:A:542:NAP:N7N	2.24	0.53
1:A:431:ASN:HA	1:A:506:LEU:O	2.08	0.53
1:A:84:ILE:HG23	1:A:89:ILE:HD11	1.90	0.52
1:A:223:ASN:HD22	1:A:226:GLN:CB	2.23	0.51
1:A:270:GLN:HE22	1:A:306:LYS:HZ2	1.54	0.51
1:A:63:HIS:CD2	1:A:230:ILE:HD13	2.46	0.50
1:A:86:GLN:HB3	1:A:87:PRO:HD3	1.93	0.50
1:A:478:THR:OG1	1:A:506:LEU:HD11	2.12	0.48
1:A:51:ASN:O	1:A:86:GLN:HG3	2.12	0.48
1:A:144:GLY:HA2	5:A:611:HOH:O	2.14	0.47
1:A:84:ILE:HG12	1:A:89:ILE:CG1	2.42	0.47
1:A:145:LEU:HD21	3:A:542:NAP:H72N	1.80	0.46
1:A:525:ASP:OD2	1:A:528:ARG:NH1	2.42	0.46
1:A:475:GLU:HA	1:A:475:GLU:OE2	2.16	0.46
1:A:201:VAL:HG21	1:A:204:LEU:HB2	1.96	0.45
1:A:159:THR:HG22	1:A:360:ALA:N	2.32	0.45
1:A:280:ARG:HD2	5:A:584:HOH:O	2.17	0.45
1:A:409:GLN:H	1:A:409:GLN:HG3	1.39	0.45
1:A:159:THR:HG21	1:A:359:THR:HA	1.99	0.45
1:A:16:ALA:HB2	1:A:106:PHE:CE2	2.52	0.44
1:A:435:THR:CG2	1:A:436:ASN:O	2.63	0.44
1:A:127:THR:CG2	1:A:133:VAL:HG22	2.48	0.44
1:A:145:LEU:N	5:A:613:HOH:O	2.51	0.44
1:A:85:THR:HB	1:A:87:PRO:HD2	2.01	0.43
1:A:63:HIS:HB2	1:A:230:ILE:HD11	2.00	0.43
1:A:84:ILE:HG13	1:A:88:GLU:HB2	2.00	0.43
1:A:63:HIS:HB2	1:A:230:ILE:CD1	2.49	0.42
1:A:223:ASN:O	1:A:224:PRO:C	2.58	0.42
1:A:507:PHE:HZ	3:A:542:NAP:O7N	2.02	0.42
1:A:198:ALA:N	1:A:199:PRO:HD2	2.35	0.42
1:A:26:LYS:HD2	1:A:26:LYS:HA	1.80	0.42
1:A:432:GLY:HA3	5:A:632:HOH:O	2.19	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:212:GLN:HE21	1:A:212:GLN:HB3	1.62	0.41
1:A:12:VAL:HG13	1:A:38:PHE:CE2	2.56	0.41
1:A:23:ALA:O	1:A:27:LEU:HG	2.20	0.41
1:A:303:ILE:O	1:A:306:LYS:HB2	2.20	0.41
1:A:507:PHE:CZ	3:A:542:NAP:O7N	2.74	0.41
1:A:86:GLN:HB3	1:A:86:GLN:HE21	1.70	0.41
1:A:150:ASN:HB2	1:A:383:ASP:N	2.36	0.41
1:A:366:GLU:OE1	1:A:366:GLU:HA	2.21	0.41
1:A:121:GLU:OE1	1:A:135:ARG:NH2	2.54	0.40
1:A:238:TRP:O	1:A:242:LYS:HG3	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	505/573 (88%)	486 (96%)	16 (3%)	3 (1%)	25 27

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	224	PRO
1	A	153	ASN
1	A	152	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	409/472 (87%)	385 (94%)	24 (6%)	19	22

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	THR
1	A	47	THR
1	A	52	ARG
1	A	86	GLN
1	A	103	ARG
1	A	107	LYS
1	A	127	THR
1	A	145	LEU
1	A	167	THR
1	A	174	LYS
1	A	193	VAL
1	A	223	ASN
1	A	225	GLU
1	A	230	ILE
1	A	234	TYR
1	A	264	GLU
1	A	293	GLU
1	A	331	LEU
1	A	369	VAL
1	A	386	ASP
1	A	409	GLN
1	A	435	THR
1	A	485	LEU
1	A	506	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	75	GLN
1	A	86	GLN
1	A	192	GLN
1	A	212	GLN
1	A	223	ASN
1	A	270	GLN
1	A	353	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	403	ASN
1	A	409	GLN
1	A	444	GLN
1	A	458	ASN
1	A	482	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	CYH	A	543	-	7,7,7	1.51	1 (14%)	8,8,8	1.25	0
3	NAP	A	542	-	45,52,52	1.92	11 (24%)	56,80,80	1.56	8 (14%)
2	FAD	A	541	-	53,58,58	1.46	7 (13%)	68,89,89	1.44	14 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CYH	A	543	-	-	-	0/1/1/1
3	NAP	A	542	-	-	8/31/67/67	0/5/5/5
2	FAD	A	541	-	-	3/30/50/50	0/6/6/6

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	541	FAD	C4X-N5	6.33	1.43	1.30
3	A	542	NAP	C2N-N1N	5.67	1.41	1.35
3	A	542	NAP	O4D-C1D	5.20	1.48	1.41
3	A	542	NAP	C3N-C7N	4.84	1.57	1.50
2	A	541	FAD	C10-N1	3.57	1.40	1.33
3	A	542	NAP	C6N-N1N	3.52	1.44	1.35
2	A	541	FAD	C10-N10	2.88	1.43	1.37
3	A	542	NAP	PN-O5D	2.85	1.70	1.59
2	A	541	FAD	C5'-C4'	2.71	1.55	1.51
4	A	543	CYH	C2-C1	2.69	1.56	1.50
3	A	542	NAP	P2B-O2X	2.59	1.64	1.54
3	A	542	NAP	PA-O2A	2.26	1.65	1.55
3	A	542	NAP	O4D-C4D	2.22	1.50	1.45
3	A	542	NAP	O4B-C1B	2.22	1.44	1.41
2	A	541	FAD	C7M-C7	2.20	1.55	1.51
2	A	541	FAD	PA-O2A	2.16	1.65	1.55
3	A	542	NAP	C5A-N7A	-2.13	1.32	1.39
3	A	542	NAP	C5D-C4D	2.09	1.58	1.51
2	A	541	FAD	O2'-C2'	-2.07	1.39	1.43

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	542	NAP	N3A-C2A-N1A	-4.80	121.18	128.68
3	A	542	NAP	O4D-C4D-C5D	4.47	124.09	109.37
2	A	541	FAD	O4B-C1B-C2B	-4.13	100.90	106.93
2	A	541	FAD	N3A-C2A-N1A	-4.11	122.25	128.68
3	A	542	NAP	O7N-C7N-C3N	3.76	124.13	119.63
3	A	542	NAP	O4D-C1D-C2D	-3.21	102.23	106.93
2	A	541	FAD	C4X-C10-N1	-2.89	118.02	124.73
2	A	541	FAD	C9A-C5X-N5	-2.85	119.34	122.43
2	A	541	FAD	C4-N3-C2	-2.72	120.62	125.64
2	A	541	FAD	C10-N1-C2	2.70	122.31	116.90
2	A	541	FAD	C4X-C4-N3	2.67	119.97	113.19
3	A	542	NAP	C6N-N1N-C2N	-2.62	119.58	121.97
2	A	541	FAD	O2'-C2'-C1'	-2.61	103.49	109.80

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	541	FAD	C10-C4X-N5	-2.45	119.66	124.86
3	A	542	NAP	PN-O5D-C5D	2.21	134.61	121.68
2	A	541	FAD	O4-C4-C4X	-2.17	120.86	126.60
2	A	541	FAD	C5X-C9A-N10	2.15	120.18	117.95
2	A	541	FAD	C4A-C5A-N7A	-2.05	107.26	109.40
3	A	542	NAP	C3N-C7N-N7N	-2.04	115.30	117.75
3	A	542	NAP	C5N-C4N-C3N	2.02	122.73	120.34
2	A	541	FAD	C4X-C10-N10	2.01	119.42	116.48
2	A	541	FAD	P-O3P-PA	-2.01	125.94	132.83

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	541	FAD	N10-C1'-C2'-O2'
2	A	541	FAD	N10-C1'-C2'-C3'
3	A	542	NAP	C5D-O5D-PN-O1N
3	A	542	NAP	C3D-C4D-C5D-O5D
3	A	542	NAP	O4D-C4D-C5D-O5D
3	A	542	NAP	C4D-C5D-O5D-PN
3	A	542	NAP	C5D-O5D-PN-O2N
2	A	541	FAD	O4B-C4B-C5B-O5B
3	A	542	NAP	C2B-O2B-P2B-O2X
3	A	542	NAP	C5D-O5D-PN-O3
3	A	542	NAP	O4B-C4B-C5B-O5B

There are no ring outliers.


2 monomers are involved in 10 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	543	CYH	6	0
3	A	542	NAP	8	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and



any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	511/573 (89%)	-0.13	10 (1%) 65 75	14, 27, 42, 62	1 (0%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	224	PRO	7.8
1	A	383	ASP	5.0
1	A	5	THR	3.7
1	A	221	PRO	2.7
1	A	131	GLY	2.6
1	A	6	THR	2.6
1	A	384	ALA	2.3
1	A	448	ILE	2.3
1	A	7	HIS	2.2
1	A	172	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	CYH	A	543	7/7	0.72	0.31	39,41,42,43	0
3	NAP	A	542	48/48	0.93	0.14	21,30,53,58	0
2	FAD	A	541	53/53	0.97	0.10	12,23,27,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.