



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 02:46 am BST

PDB ID : 4UG2  
Title : Thermostabilised HUMAN A2a Receptor with CGS21680 bound  
Authors : Lebon, G.; Edwards, P.C.; Leslie, A.G.W.; Tate, C.G.  
Deposited on : 2015-03-21  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	325	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 81%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: grey;"></div> </div>
1	B	325	<div style="display: flex; align-items: center;"> <div style="width: 0%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 80%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 15%; height: 10px; background-color: grey;"></div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4542 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THERMOSTABILISED HUMAN A2A RECEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	292	2262	1498	377	368	19	0	0	0
1	B	276	2142	1427	351	345	19	0	0	0

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	318	ALA	-	expression tag	UNP P29274
A	319	ALA	-	expression tag	UNP P29274
A	320	GLU	-	expression tag	UNP P29274
A	321	ASN	-	expression tag	UNP P29274
A	322	LEU	-	expression tag	UNP P29274
A	323	TYR	-	expression tag	UNP P29274
A	324	PHE	-	expression tag	UNP P29274
A	325	GLN	-	expression tag	UNP P29274
A	48	ALA	LEU	engineered mutation	UNP P29274
A	54	LEU	ALA	engineered mutation	UNP P29274
A	65	ALA	THR	engineered mutation	UNP P29274
A	89	ALA	GLN	engineered mutation	UNP P29274
A	154	ALA	ASN	engineered mutation	UNP P29274
B	318	ALA	-	expression tag	UNP P29274
B	319	ALA	-	expression tag	UNP P29274
B	320	GLU	-	expression tag	UNP P29274
B	321	ASN	-	expression tag	UNP P29274
B	322	LEU	-	expression tag	UNP P29274
B	323	TYR	-	expression tag	UNP P29274
B	324	PHE	-	expression tag	UNP P29274
B	325	GLN	-	expression tag	UNP P29274
B	48	ALA	LEU	engineered mutation	UNP P29274
B	54	LEU	ALA	engineered mutation	UNP P29274
B	65	ALA	THR	engineered mutation	UNP P29274
B	89	ALA	GLN	engineered mutation	UNP P29274

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	154	ALA	ASN	engineered mutation	UNP P29274

- Molecule 2 is (2S)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLB) (formula: C<sub>21</sub>H<sub>40</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			25	21	4		
2	B	1	Total	C	O	0	0
			25	21	4		

- Molecule 3 is 2-[P-(2-CARBOXYETHYL)PHENYLETHYL-AMINO]-5'-N-ETHYLCARB OXAMIDO ADENOSINE (three-letter code: NGI) (formula: C<sub>23</sub>H<sub>29</sub>N<sub>7</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
3	A	1	Total	C	N	O	0	0
			36	23	7	6		
3	B	1	Total	C	N	O	0	0
			36	23	7	6		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	7	Total	O	0	0
			7	7		
4	B	9	Total	O	0	0
			9	9		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.25Å 105.94Å 125.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	125.86 – 2.60 52.97 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.9 (125.86-2.60) 99.0 (52.97-2.60)	Depositor EDS
$R_{merge}$	0.19	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.30 (at 2.61Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.238 , 0.271 0.235 , 0.265	Depositor DCC
$R_{free}$ test set	1234 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.5	Xtrriage
Anisotropy	0.225	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 44.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	4542	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OLB, NGI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.51	1/2318 (0.0%)	0.52	0/3162
1	B	0.50	2/2194 (0.1%)	0.50	0/2994
All	All	0.50	3/4512 (0.1%)	0.51	0/6156

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	143	TRP	CD2-CE2	5.41	1.47	1.41
1	B	32	TRP	CD2-CE2	5.21	1.47	1.41
1	B	129	TRP	CD2-CE2	5.07	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2262	0	2330	17	0
1	B	2142	0	2213	9	0
2	A	25	0	40	1	0
2	B	25	0	40	0	0
3	A	36	0	28	1	0
3	B	36	0	28	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	7	0	0	0	0
4	B	9	0	0	0	0
All	All	4542	0	4679	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:13:GLU:HG3	1:B:60:ILE:HG23	1.74	0.69
1:A:13:GLU:HG3	1:A:60:ILE:HG23	1.80	0.63
1:B:66:ILE:HG23	1:B:168:PHE:HB2	1.90	0.53
1:B:66:ILE:CG2	1:B:168:PHE:HB2	2.38	0.53
1:B:174:MET:HE2	1:B:256:THR:HB	1.91	0.53
1:A:6:SER:O	1:A:10:ILE:HG12	2.12	0.49
1:A:10:ILE:HD11	1:A:68:THR:HG21	1.94	0.49
1:B:13:GLU:HG3	1:B:60:ILE:CG2	2.42	0.49
1:B:10:ILE:HD13	1:B:64:ILE:HG23	1.95	0.48
1:A:13:GLU:HG3	1:A:60:ILE:CG2	2.44	0.46
1:A:66:ILE:CG2	1:A:168:PHE:HB2	2.45	0.45
1:A:92:ILE:HG21	1:A:185:CYS:O	2.15	0.45
1:B:6:SER:O	1:B:10:ILE:HG12	2.17	0.44
1:A:55:VAL:HA	1:A:59:ALA:HB3	1.98	0.44
1:A:208:LEU:HD23	1:A:225:LEU:HD21	2.00	0.44
3:A:1311:NGI:O1	3:A:1311:NGI:H8	2.18	0.43
1:A:174:MET:HE3	1:A:174:MET:HB3	1.95	0.43
1:A:184:ALA:HB2	2:A:1310:OLB:H36	1.99	0.43
3:B:1307:NGI:O1	3:B:1307:NGI:H8	2.18	0.43
1:A:65:ALA:HB1	1:A:70:PHE:CD1	2.55	0.42
1:A:107:ARG:HD2	1:A:199:ARG:HG3	2.02	0.41
1:A:182:PHE:CD1	1:A:250:HIS:HB3	2.56	0.41
1:A:208:LEU:HG	1:A:225:LEU:HD11	2.02	0.41
1:B:296:ARG:O	1:B:300:ARG:HG3	2.21	0.41
1:B:92:ILE:HG21	1:B:185:CYS:O	2.21	0.40
1:A:66:ILE:HA	1:A:66:ILE:HD13	1.97	0.40
1:A:13:GLU:CG	1:A:60:ILE:HG23	2.51	0.40
1:A:88:THR:O	1:A:91:SER:HB3	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	288/325 (89%)	285 (99%)	3 (1%)	0	100	100
1	B	270/325 (83%)	267 (99%)	3 (1%)	0	100	100
All	All	558/650 (86%)	552 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	240/269 (89%)	236 (98%)	4 (2%)	60	81
1	B	227/269 (84%)	226 (100%)	1 (0%)	91	97
All	All	467/538 (87%)	462 (99%)	5 (1%)	73	88

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	275	VAL
1	A	282	VAL
1	A	293	ARG
1	A	309	ARG
1	B	282	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NGI	A	1311	-	33,39,39	0.73	0	39,55,55	1.54	7 (17%)
3	NGI	B	1307	-	33,39,39	0.71	0	39,55,55	1.51	6 (15%)
2	OLB	A	1310	-	24,24,24	0.92	1 (4%)	25,25,25	0.95	2 (8%)
2	OLB	B	1306	-	24,24,24	0.97	1 (4%)	25,25,25	0.86	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NGI	A	1311	-	-	2/16/38/38	0/4/4/4
3	NGI	B	1307	-	-	2/16/38/38	0/4/4/4
2	OLB	A	1310	-	-	10/24/24/24	-
2	OLB	B	1306	-	-	10/24/24/24	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1306	OLB	O20-C1	4.53	1.46	1.33
2	A	1310	OLB	O20-C1	4.25	1.45	1.33

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1311	NGI	C11-N7-C23	5.51	121.53	115.28
3	B	1307	NGI	C11-N7-C23	4.70	120.61	115.28
3	A	1311	NGI	N7-C11-N5	-4.05	119.83	126.23
3	B	1307	NGI	N7-C11-N5	-3.66	120.44	126.23
3	B	1307	NGI	C9-C10-N5	-2.95	119.07	121.01
2	A	1310	OLB	O20-C1-C2	2.93	121.11	111.91
3	B	1307	NGI	C6-C7-C4	2.80	104.90	101.64
3	B	1307	NGI	C7-C6-C5	2.60	104.89	100.98
3	A	1311	NGI	N6-C11-N5	2.58	121.66	117.19
2	B	1306	OLB	O20-C1-C2	2.51	119.79	111.91
3	A	1311	NGI	C6-C7-C4	2.44	104.48	101.64
3	A	1311	NGI	C7-C6-C5	2.39	104.58	100.98
2	A	1310	OLB	O20-C1-O19	-2.23	117.96	123.59
3	B	1307	NGI	C11-N5-C10	2.18	120.99	116.52
3	A	1311	NGI	C9-C10-N5	-2.17	119.58	121.01
3	A	1311	NGI	C11-N5-C10	2.03	120.69	116.52

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1310	OLB	O19-C1-O20-C21
2	A	1310	OLB	C2-C1-O20-C21
2	B	1306	OLB	C11-C12-C13-C14
2	A	1310	OLB	C13-C14-C15-C16
2	B	1306	OLB	C13-C14-C15-C16
2	B	1306	OLB	C1-C2-C3-C4
2	A	1310	OLB	C6-C7-C8-C9
2	A	1310	OLB	C5-C6-C7-C8
2	B	1306	OLB	O20-C21-C22-C24
2	B	1306	OLB	C15-C16-C17-C18
2	A	1310	OLB	C11-C12-C13-C14
2	A	1310	OLB	C3-C4-C5-C6
2	B	1306	OLB	O20-C21-C22-O23
2	A	1310	OLB	C15-C16-C17-C18

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
2	B	1306	OLB	C3-C4-C5-C6
2	B	1306	OLB	O20-C1-C2-C3
2	B	1306	OLB	C2-C3-C4-C5
3	A	1311	NGI	O1-C3-C4-O2
3	B	1307	NGI	O1-C3-C4-O2
3	A	1311	NGI	N1-C3-C4-O2
3	B	1307	NGI	N1-C3-C4-O2
2	A	1310	OLB	C12-C13-C14-C15
2	B	1306	OLB	C10-C11-C12-C13
2	A	1310	OLB	C4-C5-C6-C7


There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1311	NGI	1	0
3	B	1307	NGI	1	0
2	A	1310	OLB	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	292/325 (89%)	0.21	16 (5%) 25 19	24, 36, 66, 85	0
1	B	276/325 (84%)	-0.07	3 (1%) 80 78	24, 34, 62, 82	0
All	All	568/650 (87%)	0.07	19 (3%) 46 39	24, 35, 65, 85	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	110	LEU	5.0
1	A	155	HIS	3.6
1	A	292	ILE	3.6
1	A	304	ARG	3.3
1	A	300	ARG	3.2
1	A	36	ASN	3.0
1	A	297	GLN	2.9
1	B	206	ARG	2.9
1	A	261	ASP	2.7
1	B	147	GLY	2.5
1	A	224	THR	2.5
1	B	71	CYS	2.5
1	A	148	GLN	2.4
1	A	161	GLU	2.4
1	A	221	ALA	2.3
1	A	32	TRP	2.1
1	A	208	LEU	2.1
1	A	301	LYS	2.1
1	A	162	GLY	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	OLB	A	1310	25/25	0.84	0.24	43,45,55,57	0
2	OLB	B	1306	25/25	0.86	0.22	42,45,58,59	0
3	NGI	A	1311	36/36	0.94	0.17	28,29,45,46	0
3	NGI	B	1307	36/36	0.95	0.18	30,31,50,51	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around OLB B 1306:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around NGI A 1311:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.