



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 24, 2023 – 12:42 PM EDT

PDB ID : 5UJ2  
Title : Crystal structure of HCV NS5B genotype 2A JFH-1 isolate with S15G E86Q E87Q C223H V321I mutations and Delta8 neta hairpoin loop deletion in complex with GS-639476 (diphsohate version of GS-9813), Mn<sup>2+</sup> and symmetrical primer template 5'-AUAAAUUU  
Authors : Edwards, T.E.; Fox III, D.; Appleby, T.C.; Murakami, E.; Rey, A.; McGrath, M.E.  
Deposited on : 2017-01-16  
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)
RNA backbone	3102	1007 (3.16-2.64)


The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	P	8	50% 25% 25%
1	T	8	38% 62%

*Continued on next page...*

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
 Validation Pipeline (wwPDB-VP) : 2.35.1

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	A	572	 A horizontal bar chart showing the quality of chain. The bar is divided into segments: a small red segment at the beginning labeled '2%', a large green segment labeled '87%', a small yellow segment labeled '6%', and a small grey segment at the end labeled '7%'.

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 4370 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (5'-R(\*AP\*UP\*AP\*AP\*AP\*UP\*UP\*U)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	T	8	Total 147	C 66	N 23	O 51	P 7	0	0	1
1	P	6	Total 105	C 47	N 16	O 37	P 5	0	0	1

- Molecule 2 is a protein called Genome polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	A	534	Total 4061	C 2567	N 714	O 754	S 26	0	1	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	initiating methionine	UNP R9TEU1
A	0	SER	-	expression tag	UNP R9TEU1
A	15	GLY	SER	engineered mutation	UNP R9TEU1
A	86	GLN	GLU	engineered mutation	UNP R9TEU1
A	87	GLN	GLU	engineered mutation	UNP R9TEU1
A	223	HIS	CYS	engineered mutation	UNP R9TEU1
A	321	ILE	VAL	engineered mutation	UNP R9TEU1
A	?	-	ASN	deletion	UNP R9TEU1
A	?	-	PHE	deletion	UNP R9TEU1
A	?	-	GLU	deletion	UNP R9TEU1
A	?	-	MET	deletion	UNP R9TEU1
A	?	-	TYR	deletion	UNP R9TEU1
A	?	-	GLY	deletion	UNP R9TEU1
A	?	-	SER	deletion	UNP R9TEU1
A	?	-	VAL	deletion	UNP R9TEU1
A	445	GLY	SER	conflict	UNP R9TEU1
A	571	LEU	-	expression tag	UNP R9TEU1
A	572	GLU	-	expression tag	UNP R9TEU1

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	573	HIS	-	expression tag	UNP R9TEU1
A	574	HIS	-	expression tag	UNP R9TEU1
A	575	HIS	-	expression tag	UNP R9TEU1
A	576	HIS	-	expression tag	UNP R9TEU1
A	577	HIS	-	expression tag	UNP R9TEU1
A	578	HIS	-	expression tag	UNP R9TEU1

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	P	1	Total Mn 1 1	0	0
3	A	2	Total Mn 2 2	0	0

- Molecule 4 is (1S)-1-(4-aminoimidazo[2,1-f][1,2,4]triazin-7-yl)-1,4-anhydro-2-deoxy-2-fluoro-5-O-[(S)-hydroxy(phosphonoxy)phosphoryl]-2-methyl-D-ribose (three-letter code: 8B4) (formula: C<sub>11</sub>H<sub>16</sub>FN<sub>5</sub>O<sub>9</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	P	1	Total C F N O P 28 11 1 5 9 2	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Cl 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	P	2	Total O 2 2	0	0
6	A	23	Total O 23 23	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RNA (5'-R(\*AP\*UP\*AP\*AP\*AP\*UP\*UP\*U)-3')



- Molecule 1: RNA (5'-R(\*AP\*UP\*AP\*AP\*AP\*UP\*UP\*U)-3')



- Molecule 2: Genome polyprotein



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	141.76Å 141.76Å 91.11Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.90 41.35 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.2 (50.00-2.90) 99.2 (41.35-2.90)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.43 (at 2.90Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.184 , 0.227 0.192 , 0.228	Depositor DCC
$R_{free}$ test set	1167 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	63.1	Xtrriage
Anisotropy	0.017	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 35.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.039 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4370	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	60.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, 8B4, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	P	0.33	0/116	0.66	0/179
1	T	0.35	0/163	0.68	0/252
2	A	0.45	0/4159	0.67	0/5672
All	All	0.45	0/4438	0.67	0/6103

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	P	105	0	52	1	0
1	T	147	0	74	2	0
2	A	4061	0	3985	16	0
3	A	2	0	0	0	0
3	P	1	0	0	0	0
4	P	28	0	0	1	0
5	A	1	0	0	0	0
6	A	23	0	0	0	0
6	P	2	0	0	0	0
All	All	4370	0	4111	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:85:LEU:C	2:A:85:LEU:HD23	2.25	0.57
1:P:6:U:H2'	1:P:7:U:C6	2.40	0.57
2:A:18:GLU:CB	2:A:399:THR:HG21	2.37	0.55
2:A:203:TYR:CD2	2:A:314:LEU:HD13	2.44	0.53
2:A:203:TYR:CE2	2:A:314:LEU:HD13	2.47	0.50
1:T:4:A:H2'	1:T:5:A:C8	2.48	0.49
2:A:21:LEU:HB3	2:A:22:PRO:HD2	1.95	0.49
2:A:399:THR:HG22	2:A:401:ARG:H	1.78	0.49
1:T:2:U:O2	2:A:283:GLY:HA2	2.14	0.48
2:A:86:GLN:O	2:A:90:GLN:HG2	2.14	0.47
2:A:132:THR:O	2:A:259:ARG:NE	2.38	0.45
4:P:602:8B4:O2B	4:P:602:8B4:O3'	2.35	0.44
2:A:60:LEU:HD13	2:A:64:TYR:CE1	2.53	0.44
2:A:174:ALA:HB1	2:A:253:ILE:HD13	1.99	0.44
2:A:219:TYR:HB3	2:A:320:LEU:HD21	1.99	0.44
2:A:90:GLN:HA	2:A:90:GLN:HE21	1.83	0.43
2:A:84:THR:O	2:A:85:LEU:C	2.56	0.42
2:A:48:ARG:HG2	2:A:159:LEU:HG	2.01	0.41
2:A:524:TYR:CZ	2:A:539:LEU:HD21	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	A	533/572 (93%)	513 (96%)	19 (4%)	1 (0%)	47 78

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	130	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	A	428/486 (88%)	414 (97%)	14 (3%)	38 72

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	5	SER
2	A	19	GLU
2	A	29	SER
2	A	56	ARG
2	A	90	GLN
2	A	112	SER
2	A	117	ASN
2	A	121	SER
2	A	194	GLN
2	A	320	LEU
2	A	324	SER
2	A	336	LEU
2	A	473	SER
2	A	483	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	90	GLN

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	P	4/8 (50%)	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	T	6/8 (75%)	2 (33%)	0
All	All	10/16 (62%)	2 (20%)	0

All (2) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	T	7	U
1	T	8	U

There are no RNA pucker outliers to report.

#### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	8B4	P	602	3	18,30,30	0.70	0	24,48,48	1.35	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	8B4	P	602	3	-	2/12/36/36	0/3/3/3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	P	602	8B4	N3-C2-N1	-4.57	121.42	128.59

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	P	602	8B4	PA-O3A-PB-O1B
4	P	602	8B4	PA-O3A-PB-O2B

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	P	602	8B4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	P	6/8 (75%)	-0.08	0 <a href="#">100</a> <a href="#">100</a>	72, 74, 98, 123	0
1	T	8/8 (100%)	0.00	0 <a href="#">100</a> <a href="#">100</a>	57, 60, 100, 134	0
2	A	534/572 (93%)	-0.24	12 (2%) <a href="#">62</a> <a href="#">59</a>	38, 57, 83, 105	0
All	All	548/588 (93%)	-0.23	12 (2%) <a href="#">62</a> <a href="#">59</a>	38, 57, 84, 134	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	534	LEU	4.5
2	A	532	THR	3.4
2	A	537	THR	3.4
2	A	536	LEU	2.6
2	A	0	SER	2.5
2	A	509	ALA	2.5
2	A	535	LYS	2.4
2	A	57	THR	2.3
2	A	533	LYS	2.2
2	A	530	VAL	2.2
2	A	527	ASN	2.1
2	A	524	TYR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	CL	A	603	1/1	0.90	0.17	70,70,70,70	0
3	MN	A	602	1/1	0.95	0.08	100,100,100,100	0
3	MN	P	601	1/1	0.99	0.12	46,46,46,46	0
4	8B4	P	602	28/28	0.99	0.14	45,55,58,59	0
3	MN	A	601	1/1	0.99	0.13	53,53,53,53	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.