Full wwPDB X-ray Structure Validation Report (i)

Oct 4, 2023 – 04:46 AM EDT

PDB ID	:	6UKG
Title	:	HhaI endonuclease in Complex With DNA in space group P21 (pH 4.2)
Authors	:	Horton, J.R.; Cheng, X.
Deposited on	:	2019-10-04
Resolution	:	1.16 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1758 (1.20-1.12)
Clashscore	141614	1832 (1.20-1.12)
Ramachandran outliers	138981	1768 (1.20-1.12)
Sidechain outliers	138945	1768 (1.20-1.12)
RSRZ outliers	127900	1724 (1.20-1.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	258	9% 90% 10%
2	A	23	39% 35% 48% 17%
3	B	23	52% 48% 48% .

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 5961 atoms, of which 2633 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HhaI Restriction Endonuclease.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	X	258	4256	1384	2124	345	395	8	0	9	0

- Molecule 2 is a DNA chain called DNA (5'-D(*CP*TP*GP*TP*TP*GP*CP*GP*CP*TP*TP*GP*GP*A)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P			
2	A	14	693	214	246	80	132	21	0	9	0

- Molecule 3 is a DNA chain called DNA (5'-D(*CP*CP*AP*AP*GP*CP*GP*CP*AP*AP*CP*AP*G)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P			
3	B	13	692	215	245	85	127	20	0	9	0

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	X	1	Total O P 5 4 1	0	0

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	X	1	Total C H O 10 2 6 2	0	0
5	X	1	Total C H O 10 2 6 2	0	0
5	B	1	Total C H O 10 2 6 2	0	0

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
6	X	1	10	8	2	0	1
			Total	O	S		
6	X	1	5	4	1	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	X	227	227	227	0	0
			Total	O		
7	A	25	25	25	0	0
			Total	O		
7	B	18	18	18	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: HhaI Restriction Endonuclease

Chain X:





• Molecule 2: DNA (5'-D(*CP*TP*GP*TP*TP*GP*CP*GP*CP*TP*TP*GP*GP*A)-3')

Chain A:



• Molecule 3: DNA (5'-D(*CP*CP*AP*AP*GP*CP*GP*CP*AP*AP*CP*AP*G)-3')

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	61.85Å 37.55Å 69.92Å 90.00° 110.26° 90.00°	Depositor
Resolution (Å)	31.52 – 1.16 31.52 – 1.12	Depositor EDS
% Data completeness (in resolution range)	88.9 (31.52-1.16) 79.7 (31.52-1.12)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.26 (at 1.12Å)	Xtriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, R_{free}	0.141 , 0.157 0.141 , 0.156	Depositor DCC
R_{free} test set	1919 reflections (2.00%)	wwPDB-VP
Wilson B-factor (Å^2)	11.4	Xtriage
Anisotropy	0.067	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.50 , 56.4	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	5961	wwPDB-VP
Average B, all atoms (Å^2)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.20% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, PO4, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	X	0.69	13/2216 (0.6%)	0.59	0/2981
2	A	2.71	6/491 (1.2%)	1.82	14/735 (1.9%)
3	B	1.70	5/491 (1.0%)	1.75	11/730 (1.5%)
All	All	1.38	24/3198 (0.8%)	1.13	25/4446 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	A	0	1

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	12[B]	DC	O3'-P	40.48	2.09	1.61
2	A	4[B]	DA	O3'-P	-36.13	1.17	1.61
3	B	10[B]	DT	O3'-P	21.71	1.87	1.61
3	B	2[B]	DT	O3'-P	16.76	1.81	1.61
3	B	12[B]	DG	O3'-P	-14.49	1.43	1.61
2	A	3[B]	DC	O3'-P	13.81	1.77	1.61
3	B	13[B]	DG	O3'-P	-13.56	1.44	1.61
2	A	1[B]	DT	O3'-P	-10.77	1.48	1.61
2	A	2[B]	DC	O3'-P	-9.18	1.50	1.61
1	X	145	GLU	CD-OE2	7.67	1.34	1.25
1	X	7	GLU	CD-OE2	7.33	1.33	1.25
1	X	91	GLU	CD-OE2	6.94	1.33	1.25
1	X	194	GLU	CD-OE2	6.76	1.33	1.25
1	X	142	GLU	CD-OE2	6.69	1.33	1.25
1	X	194	GLU	CD-OE1	-6.59	1.18	1.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	5	GLU	CD-OE2	6.51	1.32	1.25
1	X	46	GLU	CD-OE2	6.29	1.32	1.25
3	B	3[B]	DG	O3'-P	6.19	1.68	1.61
1	X	7	GLU	CD-OE1	-5.77	1.19	1.25
1	X	5	GLU	CD-OE1	-5.75	1.19	1.25
1	X	91	GLU	CD-OE1	-5.63	1.19	1.25
1	X	110	GLU	CD-OE1	-5.39	1.19	1.25
2	A	10[B]	DA	O3'-P	-5.33	1.54	1.61
1	X	46	GLU	CD-OE1	-5.14	1.20	1.25

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	B	3[B]	DG	OP1-P-O3'	-20.39	60.35	105.20
3	B	2[B]	DT	P-O3'-C3'	19.26	142.81	119.70
2	A	3[B]	DC	O3'-P-O5'	-17.79	70.19	104.00
2	A	11[B]	DA	OP1-P-O3'	-17.69	66.28	105.20
2	A	3[B]	DC	OP1-P-O3'	16.13	140.69	105.20
2	A	11[B]	DA	O3'-P-O5'	15.80	134.02	104.00
3	B	10[B]	DT	P-O3'-C3'	-12.85	104.28	119.70
3	B	3[B]	DG	OP2-P-O3'	11.28	130.02	105.20
2	A	11[B]	DA	P-O3'-C3'	-10.57	107.01	119.70
3	B	12[B]	DG	O3'-P-O5'	-10.05	84.90	104.00
3	B	12[B]	DG	OP1-P-O3'	9.52	126.15	105.20
2	A	10[B]	DA	P-O3'-C3'	8.42	129.81	119.70
3	B	3[B]	DG	O3'-P-O5'	7.96	119.11	104.00
2	A	12[B]	DC	OP2-P-O3'	7.86	122.49	105.20
2	A	1[B]	DT	P-O3'-C3'	-7.35	110.88	119.70
3	B	13[B]	DG	P-O3'-C3'	7.09	128.21	119.70
2	A	4[B]	DA	O3'-P-O5'	7.00	117.30	104.00
3	B	10[B]	DT	OP2-P-O3'	6.45	119.39	105.20
3	B	9	DC	O4'-C4'-C3'	-6.38	101.95	104.50
2	A	3[B]	DC	P-O3'-C3'	6.33	127.30	119.70
2	A	10[B]	DA	O3'-P-O5'	6.08	115.55	104.00
3	B	2[B]	DT	O3'-P-O5'	5.75	114.93	104.00
2	A	2[B]	DC	P-O3'-C3'	-5.64	112.93	119.70
2	A	1[B]	DT	OP2-P-O3'	5.39	117.05	105.20
2	A	12[B]	DC	O3'-P-O5'	-5.34	93.85	104.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	A	8	DG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	2132	2124	2116	19	1
2	A	447	246	247	2	0
3	B	447	245	247	0	0
4	X	5	0	0	0	0
5	B	4	6	6	0	0
5	X	8	12	12	0	0
6	X	15	0	0	1	0
7	A	25	0	0	1	0
7	B	18	0	0	0	0
7	X	227	0	0	4	1
All	All	3328	2633	2628	21	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:12[A]:DG:OP2	7:A:101:HOH:O	1.85	0.93
1:X:25[B]:LYS:HG2	1:X:33:SER:O	1.90	0.71
1:X:153[B]:THR:HG23	7:X:1044:HOH:O	1.97	0.65
1:X:105:ILE:HD12	1:X:153[B]:THR:HG22	1.81	0.61
1:X:245:LEU:HD22	7:X:935:HOH:O	2.00	0.60
1:X:220[A]:LYS:NZ	7:X:905:HOH:O	2.33	0.60
1:X:122:LYS:CG	1:X:141:ILE:HD11	2.33	0.58
1:X:25[B]:LYS:HD2	1:X:26:GLY:N	2.19	0.57
1:X:122:LYS:HG3	1:X:141:ILE:HD11	1.92	0.52
1:X:122:LYS:HG3	1:X:141:ILE:CD1	2.40	0.51
1:X:25[B]:LYS:HD3	1:X:32:THR:HG23	1.92	0.50
1:X:105:ILE:HD12	1:X:153[B]:THR:CG2	2.40	0.50
1:X:122:LYS:HG2	1:X:141:ILE:HD11	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:25[B]:LYS:CE	7:X:914:HOH:O	2.61	0.48
1:X:137:ILE:HG23	1:X:245:LEU:CD2	2.45	0.47
1:X:25[B]:LYS:HD2	1:X:26:GLY:O	2.15	0.46
2:A:11[A]:DT:H2"	2:A:12[A]:DG:C8	2.52	0.44
1:X:238:ASN:ND2	6:X:803[A]:SO4:O4	2.52	0.43
1:X:25[B]:LYS:HD3	1:X:32:THR:CG2	2.50	0.42
1:X:78:ILE:CG2	1:X:83:GLN:HG3	2.50	0.41
1:X:174:GLN:HG2	1:X:178:LYS:HE2	2.03	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:239:ASN:HD22	7:X:906:HOH:O[2_555]	1.52	0.08

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	X	266/258 (103%)	260 (98%)	6 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	X	240/237 (101%)	239 (100%)	1 (0%)	91 75

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	229	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	EDO	X	804	-	3,3,3	0.44	0	2,2,2	0.30	0
4	PO4	X	801	-	4,4,4	0.89	0	6,6,6	0.49	0
5	EDO	X	802	-	3,3,3	0.47	0	2,2,2	0.34	0
5	EDO	B	1801	-	3,3,3	0.45	0	2,2,2	0.14	0
6	SO4	X	805	-	4,4,4	0.15	0	6,6,6	0.09	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
6	SO4	X	803[B]	-	4,4,4	0.18	0	6,6,6	0.15	0
6	SO4	X	803[A]	-	4,4,4	0.18	0	6,6,6	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	X	802	-	-	0/1/1/1	-
5	EDO	B	1801	-	-	0/1/1/1	-
5	EDO	X	804	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	X	804	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	X	803[A]	SO4	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	X	258/258 (100%)	0.78	23 (8%)	9	10	8, 15, 29, 52	0
2	A	23/23 (100%)	2.17	9 (39%)	0	1	9, 20, 45, 49	18 (78%)
3	B	22/23 (95%)	2.44	12 (54%)	0	0	12, 24, 44, 48	18 (81%)
All	All	303/304 (99%)	1.01	44 (14%)	2	3	8, 16, 35, 52	36 (11%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	213	ASN	11.4
2	A	14	DA	9.5
1	X	214	ASN	7.0
1	X	215	GLY	6.9
3	B	14[A]	DG	6.7
3	B	14[A]	DG	6.7
2	A	1[A]	DC	5.3
2	A	1[A]	DC	5.3
1	X	216	ALA	6.2
3	B	13[A]	DA	5.0
3	B	13[A]	DA	5.0
1	X	258	LEU	4.8
1	X	187	GLY	4.1
3	B	2[A]	DC	3.0
3	B	2[A]	DC	3.0
2	A	13[A]	DG	4.0
2	A	13[A]	DG	4.0
1	X	178	LYS	3.0
3	B	4[A]	DA	2.9
3	B	4[A]	DA	2.9
1	X	139	PHE	2.9
3	B	3[A]	DC	2.0
3	B	3[A]	DC	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	A	2[A]	DT	3.4
2	A	2[A]	DT	3.4
1	X	129	PHE	2.8
1	X	117	ILE	2.8
1	X	218	HIS	2.8
1	X	167	LYS	2.7
1	X	120	TYR	2.7
1	X	203	GLY	2.4
1	X	205	ASP	2.4
3	B	12[A]	DC	1.5
3	B	12[A]	DC	1.5
1	X	128	PHE	2.3
1	X	126	VAL	2.3
1	X	144	PHE	2.3
1	X	188	LEU	2.2
1	X	1	MET	2.1
1	X	204	ASP	2.1
1	X	54	CYS	2.1
1	X	130	ILE	2.1
2	A	12[A]	DG	2.7
2	A	12[A]	DG	2.7

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	EDO	B	1801	4/4	0.72	0.26	54,64,67,67	0
5	EDO	X	802	4/4	0.75	0.14	52,63,65,66	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	PO4	X	801	5/5	0.89	0.13	59,59,60,60	0
5	EDO	X	804	4/4	0.90	0.20	33,39,44,46	0
6	SO4	X	805	5/5	0.92	0.18	33,33,34,35	0
6	SO4	X	803[B]	5/5	0.98	0.09	13,13,14,15	5
6	SO4	X	803[A]	5/5	0.98	0.09	18,18,21,21	5

6.5 Other polymers (i)

There are no such residues in this entry.