

Full wwPDB X-ray Structure Validation Report (i)

Jul 31, 2023 – 05:04 AM EDT

PDB ID : 1UNA

Title : UNASSEMBLED VIRUS COAT PROTEIN DIMER, BACTERIOPHAGE

RNA-BINDING DIMER

Authors : Ni, C.-Z.; Ely, K.R.

Deposited on : 1996-04-25

Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.34

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain				
1	A	129	61%	23%	9% • 5%		
1	В	129	52%	32%	9% • 5%		



2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1830 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called GA UNASSEMBLED COAT PROTEIN DIMER.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
1	Λ	122	Total	С	N	О	0	0	0
1	A	122	915	583	154	178	0	U	
1	D	122	Total	С	N	О	0	0	0
1	Б	122	915	583	154	178	0	U	U

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	5	HIS	ARG	conflict	UNP P07234
A	59	THR	ALA	conflict	UNP P07234
A	79	VAL	GLY	conflict	UNP P07234
A	109	THR	ALA	conflict	UNP P07234
В	5	HIS	ARG	conflict	UNP P07234
В	59	THR	ALA	conflict	UNP P07234
В	79	VAL	GLY	conflict	UNP P07234
В	109	THR	ALA	conflict	UNP P07234





3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: GA UNASSEMBLED COAT PROTEIN DIMER







4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	P 21 21 21	Depositor	
Cell constants	59.30Å 60.50Å 67.10Å	Depositor	
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor	
Resolution (Å)	6.00 - 2.80	Depositor	
% Data completeness	50.0 (6.00-2.80)	Depositor	
(in resolution range)	90.0 (0.00 2.00)	Depositor	
R_{merge}	0.07	Depositor	
R_{sym}	(Not available)	Depositor	
Refinement program	PROLSQ, X-PLOR	Depositor	
R, R_{free}	0.204 , 0.284	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	1830	wwPDB-VP	
Average B, all atoms (Å ²)	15.0	wwPDB-VP	



5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain		Bond lengths		ond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.66	0/932	1.32	4/1269 (0.3%)
1	В	0.68	0/932	1.42	4/1269 (0.3%)
All	All	0.67	0/1864	1.37	8/2538 (0.3%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	В	77	LEU	CA-CB-CG	9.16	136.36	115.30
1	В	82	TRP	CA-CB-CG	6.82	126.67	113.70
1	В	5	HIS	CA-CB-CG	6.63	124.88	113.60
1	A	58	TYR	CA-CB-CG	5.77	124.36	113.40
1	В	128	PHE	CB-CA-C	5.14	120.68	110.40
1	A	90	LEU	CA-CB-CG	5.06	126.93	115.30
1	A	82	TRP	N-CA-C	-5.05	97.38	111.00
1	A	101	VAL	CA-CB-CG1	5.04	118.46	110.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	915	0	920	32	0
1	В	915	0	920	43	1
All	All	1830	0	1840	59	1



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap (Å)
1:B:101:VAL:HG13	1:B:104:ILE:HD12	1.73	0.71
1:A:124:SER:HB2	1:A:126:SER:HB2	1.72	0.70
1:A:57:LYS:HE2	1:A:89:ASP:HB3	1.72	0.70
1:A:77:LEU:HA	1:A:80:SER:HB3	1.72	0.69
1:B:20:VAL:HG22	1:B:32:TRP:HB3	1.75	0.69
1:A:66:LYS:HE3	1:B:101:VAL:HG22	1.77	0.67
1:B:66:LYS:HB3	1:B:68:VAL:HG22	1.78	0.66
1:A:117:PRO:HB3	1:B:3:THR:O	2.03	0.59
1:B:66:LYS:O	1:B:68:VAL:N	2.39	0.56
1:A:58:TYR:HB2	1:A:90:LEU:HD23	1.87	0.55
1:A:90:LEU:HA	1:B:85:TYR:O	2.07	0.54
1:A:95:PHE:H	1:B:125:GLN:HE22	1.57	0.53
1:A:3:THR:CG2	1:B:121:ALA:HB2	2.39	0.52
1:B:55:LYS:HA	1:B:93:PRO:HA	1.92	0.51
1:A:113:LYS:O	1:A:116:ASN:HB2	2.12	0.50
1:B:66:LYS:C	1:B:68:VAL:H	2.14	0.50
1:B:25:ASN:O	1:B:27:ASN:N	2.45	0.50
1:A:12:ASN:ND2	1:A:16:GLY:O	2.39	0.49
1:A:112:PHE:CD1	1:B:108:LEU:HD13	2.47	0.49
1:A:3:THR:O	1:B:117:PRO:HB3	2.12	0.49
1:B:76:GLU:O	1:B:78:PRO:HD3	2.11	0.49
1:A:77:LEU:N	1:A:78:PRO:HD3	2.28	0.49
1:B:4:LEU:HD22	1:B:22:PRO:HB3	1.94	0.48
1:B:66:LYS:NZ	1:B:66:LYS:HB2	2.28	0.48
1:A:87:SER:O	1:B:88:ILE:HA	2.14	0.48
1:B:113:LYS:O	1:B:116:ASN:HB2	2.14	0.48
1:A:90:LEU:HD11	1:A:92:ILE:HD11	1.96	0.48
1:B:18:VAL:HG11	1:B:42:TYR:CD1	2.49	0.48
1:B:78:PRO:HD2	1:B:79:VAL:H	1.79	0.47
1:B:55:LYS:HD3	1:B:91:THR:CG2	2.45	0.46
1:A:110:GLY:HA3	1:B:9:LEU:O	2.16	0.46
1:A:129:TYR:CE1	1:B:2:ALA:HA	2.50	0.46
1:A:37:SER:O	1:A:38:ARG:C	2.54	0.46
1:A:2:ALA:O	1:A:5:HIS:HD2	1.99	0.46
1:A:58:TYR:O	1:A:89:ASP:HA	2.16	0.45
1:A:101:VAL:HG22	1:B:66:LYS:HD3	1.99	0.45
1:A:101:VAL:HG22	1:B:66:LYS:CD	2.47	0.45

Continued on next page...



Continued from previous page...

A Land		Interatomic	Clash	
Atom-1	Atom-2	${\rm distance}\ ({\rm \AA})$	overlap (Å)	
1:A:90:LEU:CD1	1:A:92:ILE:HD11	2.47	0.45	
1:A:33:LEU:CD1	1:A:43:ARG:HG2	2.47	0.44	
1:B:49:ARG:HG3	1:B:50:ALA:N	2.32	0.44	
1:B:76:GLU:C	1:B:78:PRO:HD3	2.38	0.44	
1:A:94:ILE:HA	1:B:125:GLN:NE2	2.33	0.43	
1:B:55:LYS:HB3	1:B:93:PRO:HA	2.01	0.43	
1:B:43:ARG:O	1:B:62:LEU:HD12	2.19	0.43	
1:A:58:TYR:CB	1:A:90:LEU:HD23	2.49	0.42	
1:B:63:GLU:HG2	1:B:85:TYR:CD1	2.54	0.42	
1:A:90:LEU:HD13	1:A:104:ILE:HD11	2.02	0.42	
1:B:66:LYS:C	1:B:68:VAL:N	2.72	0.42	
1:A:85:TYR:O	1:B:90:LEU:HA	2.20	0.42	
1:B:37:SER:O	1:B:38:ARG:C	2.57	0.42	
1:B:21:VAL:HG23	1:B:22:PRO:HD2	2.02	0.42	
1:B:25:ASN:HA	1:B:29:VAL:O	2.19	0.42	
1:B:64:VAL:HA	1:B:65:PRO:HD2	1.90	0.41	
1:A:9:LEU:O	1:B:110:GLY:HA3	2.20	0.41	
1:B:68:VAL:HG23	1:B:82:TRP:CD1	2.55	0.41	
1:B:77:LEU:HB2	1:B:80:SER:O	2.21	0.41	
1:A:3:THR:HG22	1:B:128:PHE:CD1	2.57	0.40	
1:B:55:LYS:HD3	1:B:91:THR:HG22	2.03	0.40	
1:A:111:LEU:HD23	1:A:112:PHE:CZ	2.57	0.40	

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{aligned} ext{Interatomic} \ ext{distance} & (ext{Å}) \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:B:38:ARG:NH2	1:B:125:GLN:OE1[4_4511]	2.00	0.20

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	118/129 (92%)	107 (91%)	9 (8%)	2 (2%)	9 29
1	В	118/129 (92%)	108 (92%)	4 (3%)	6 (5%)	2 6
All	All	$236/258 \; (92\%)$	215 (91%)	13 (6%)	8 (3%)	3 13

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	67	ILE
1	В	26	ALA
1	В	67	ILE
1	В	27	ASN
1	A	38	ARG
1	В	38	ARG
1	В	78	PRO
1	В	28	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		les	
1	A	99/105 (94%)	82 (83%)	17 (17%)		2	6	
1	В	99/105 (94%)	83 (84%)	16 (16%)		2	7	
All	All	198/210 (94%)	165 (83%)	33 (17%)		2	6	

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	SER
1	A	25	ASN
1	A	43	ARG
1	A	49	ARG
1	A	57	LYS
1	A	59	THR
1	A	62	LEU
1	A	64	VAL

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	
1	A	66	LYS	
1	A	77	LEU	
1	A	82	TRP	
1	A	87	SER	
1	A	90	LEU	
1	A	104	ILE	
1	A	109	THR	
1	A	124	SER	
1	A	126	SER	
1	В	3	THR	
1	В	4	LEU	
1	В	6	SER	
1	В	25	ASN	
1	В	38	ARG	
1	В	43	ARG	
1	В	49	ARG	
1	В	66	LYS	
1	В	77	LEU	
1	В	82	TRP	
1	В	101	VAL	
1	В	109	THR	
1	В	113	LYS	
1	В	122	ILE	
1	В	124	SER	
1	В	126	SER	

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	ASN
1	A	40	GLN
1	В	5	HIS
1	В	25	ASN
1	В	125	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

