

# Full wwPDB X-ray Structure Validation Report (i)

#### Jan 3, 2024 - 05:03 pm GMT

PDB ID	:	4UO7
Title	:	Structure of the A_Canine_Colorado_17864_06 H3 haemagglutinin in
		complex with 6SO4 Sialyl Lewis X
Authors	:	Vachieri, S.G.; Collins, P.J.; Haire, L.F.; Ogrodowicz, R.W.; Martin, S.R.;
		Walker, P.A.; Xiong, X.; Gamblin, S.J.; Skehel, J.J.
Deposited on	:	2014-05-31
Resolution	:	3.00  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber $(2001)$
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R <sub>free</sub>	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	А	328	% 	11%		
-		020		1170	•	
1	С	328	88%	9%	•	•
1	Ε	328	86%	10%	•	•
2	В	175	.% 97%		•	
2	D	175	94%		5%	



Mol	Chain	Length	Quality of chai	n
2	F	175	94%	6% ·
3	G	2	100%	
3	L	2	100%	
3	М	2	100%	
3	Ο	2	100%	
3	Q	2	100%	
3	R	2	100%	
3	Т	2	50%	50%
4	Н	3	100%	
4	J	3	67%	33%
5	Ι	5	40%	60%
5	N	5	20% 80%	
5	S	5	20% 80%	
6	K	4	100%	
6	Р	4	100%	
6	U	4	75%	25%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	G	2	Х	-	-	-
3	NAG	L	2	-	-	-	Х
3	NAG	Q	2	-	-	-	Х
3	NAG	Т	2	-	-	-	Х
5	MAN	S	4	-	-	-	Х
6	GAL	K	2	Х	-	-	-
6	FUC	K	4	Х	-	-	-
6	FUC	Р	4	Х	-	-	-
6	FUC	U	4	Х	-	-	-



# 2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 12430 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	1 1	210	Total	С	Ν	0	$\mathbf{S}$	0	0	0
I A	519	2458	1534	435	473	16	0	0	0	
1	1 C	319	Total	С	Ν	0	S	0	0	0
			2469	1541	437	475	16	0		0
1	1 E	210	Total	С	Ν	0	S	0	0	0
	519	2471	1542	436	477	16	0		0	

• Molecule 1 is a protein called HAEMAGGLUTININ HA1.

• Molecule 2 is a protein called HAEMAGGLUTININ HA2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	В	174	Total	С	Ν	0	S	0	0	0
2 B	174	1404	874	242	282	6	0	0	0	
0	р	175	Total	С	Ν	0	S	0	0	0
	2 D	175	1405	874	242	283	6	0	0	0
0	F	174	Total	С	Ν	0	S	0	0	0
	2 F	1/4	1404	875	243	280	6	0		0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	173	SER	-	expression tag	UNP E0UVR5
В	174	GLY	-	expression tag	UNP E0UVR5
В	175	ARG	-	expression tag	UNP E0UVR5
В	131	GLU	ASP	conflict	UNP E0UVR5
D	173	SER	-	expression tag	UNP E0UVR5
D	174	GLY	-	expression tag	UNP E0UVR5
D	175	ARG	-	expression tag	UNP E0UVR5
D	131	GLU	ASP	conflict	UNP E0UVR5
F	173	SER	-	expression tag	UNP E0UVR5
F	174	GLY	-	expression tag	UNP E0UVR5
F	175	ARG	-	expression tag	UNP E0UVR5
F	131	GLU	ASP	conflict	UNP E0UVR5





• Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	G	2	Total         C         N         O           28         16         2         10	0	0	0
3	L	2	Total         C         N         O           28         16         2         10	0	0	0
3	М	2	Total         C         N         O           28         16         2         10	0	0	0
3	0	2	Total         C         N         O           28         16         2         10	0	0	0
3	Q	2	Total         C         N         O           28         16         2         10	0	0	0
3	R	2	Total         C         N         O           28         16         2         10	0	0	0
3	Т	2	Total         C         N         O           28         16         2         10	0	0	0

• Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-b eta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	Н	3	Total 39	C 22	N 2	O 15	0	0	0
4	J	3	Total 39	С 22	N 2	O 15	0	0	0

• Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyran ose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	Ι	5	Total 61	С 34	N 2	O 25	0	0	0
5	N	5	Total 61	C 34	N 2	O 25	0	0	0
5	S	5	Total 61	C 34	N 2	O 25	0	0	0

• Molecule 6 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galacto pyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-6-O-sulfo-beta-D-glucopyr anose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
6	K	4	Total	С	Ν	Ο	S	0	0	0
0		4	60	31	2	26	1	0	0	0
6	D	4	Total	С	Ν	Ο	$\mathbf{S}$	0	0	0
0	1	4	60	31	2	26	1	0	0	0
6	U		Total	С	Ν	Ο	S	0	0	0
U		4	60	31	2	26	1	0	0	0

• Molecule 7 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
7	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
7	С	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
7	С	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
7	D	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
7	F	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Α	tor	ns		ZeroOcc	AltConf
8	С	1	Total 14	C 8	N 1	O 5	0	0

• Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	А	43	Total O 43 43	0	0
9	В	12	Total         O           12         12	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	С	36	Total O 36 36	0	0
9	D	14	Total         O           14         14	0	0
9	Е	30	Total         O           30         30	0	0
9	F	3	Total O 3 3	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: HAEMAGGLUTININ HA1



• Molecule 2: HAEM	AGGLUTININ HA2	
Chain D:	94%	5% •
1 16 R.25 R.25 29 434 434 K51 K51 L110 L110	<b>6128</b> <b>8173</b> <b>8173</b> <b>8173</b>	
• Molecule 2: HAEM	AGGLUTININ HA2	
Chain F:	94%	6% •
61 647 K51 L98 L98 L102 L110 L110 V141	C144 H159 Q172 G174 ARG	
• Molecule 3: 2-acetar opyranose	nido-2-deoxy-beta-D-glucopyranose	e-(1-4)-2-acetamido-2-deoxy-beta-D-gluc
Chain G:	100%	

#### NAG1 NAG2

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluc opyranose

$\alpha$		т
(Ch	am	· ·
UII	am	ц.

100%

#### NAG1 NAG2

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluc opyranose

Chain	М	:
CHOIN	<b>T</b> . <b>T</b>	٠

100%

#### NAG1 NAG2

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

$\alpha_1$ ·	$\sim$	
( 'hain	().	
Unam	v.	
	-	

100%

#### NAG1 NAG2

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain Q:

100%



#### NAG1 NAG2

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

100%

#### NAG1 NAG2

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain T:	50%	50%

#### NAG1 NAG2

• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:	100%
MAG1 MAC2 BMA3	

• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I.	670/	220/
Unann J.	67%	33%

#### NAG1 NAG2 BMA3

 • Molecule 5: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)] beta-D-mannopyrano<br/> se-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:	40%	60%

NAG1 NAG2 BMA3 MAN4 MAN5

 • Molecule 5: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)] beta-D-mannopyrano<br/> se-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain N:	20%	80%
NAG1 NAG2 BMA3 MAN4 MAN5		



 $\bullet$  Molecule 5: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)] beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose nose

80%

Chain S: 20%

#### NAG1 NAG2 BMA3 MAN4 MAN5 MAN5

 $\bullet \ Molecule \ 6: \ N-acetyl-alpha-neuraminic \ acid-(2-3)-beta-D-galactopyranose-(1-4)-[alpha-L-fucopy ranose-(1-3)] 2-acetamido-2-deoxy-6-O-sulfo-beta-D-glucopyranose$ 

Chain K:

100%

NGS1 GAL2 SIA3 FUC4

 $\bullet \ Molecule \ 6: \ N-acetyl-alpha-neuraminic \ acid-(2-3)-beta-D-galactopyranose-(1-4)-[alpha-L-fucopy ranose-(1-3)]2-acetamido-2-deoxy-6-O-sulfo-beta-D-glucopyranose$ 

Chain P:

100%

75%

NGS1 GAL2 SIA3 FUC4

 $\bullet \ Molecule \ 6: \ N-acetyl-alpha-neuraminic \ acid-(2-3)-beta-D-galactopyranose-(1-4)-[alpha-L-fucopy ranose-(1-3)] 2-acetamido-2-deoxy-6-O-sulfo-beta-D-glucopyranose$ 

25%

Chain U:

NGS1 GAL2 SIA3 FUC4



# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants	167.31Å 348.49Å 95.81Å	Deperitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
$\mathbf{Posolution} \left( \overset{\circ}{\mathbf{A}} \right)$	174.24 - 3.00	Depositor
Resolution (A)	47.72 - 3.00	EDS
% Data completeness	94.9 (174.24-3.00)	Depositor
(in resolution range)	94.9 (47.72-3.00)	EDS
R <sub>merge</sub>	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.07 (at 3.01 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0069	Depositor
B B.	0.233 , $0.276$	Depositor
II, II, <i>free</i>	0.234 , $0.279$	DCC
$R_{free}$ test set	2728 reflections $(5.09\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	58.1	Xtriage
Anisotropy	0.709	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.30 , $37.7$	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	12430	wwPDB-VP
Average B, all atoms $(Å^2)$	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.57% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NGS, SO4, SIA, GAL, NAG, MAN, FUC, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bo	ond angles
IVIOI	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.33	0/2509	0.56	0/3407
1	С	0.34	0/2520	0.56	0/3419
1	Е	0.34	0/2522	0.57	0/3422
2	В	0.35	0/1429	0.53	1/1924~(0.1%)
2	D	0.35	0/1430	0.54	0/1927
2	F	0.33	0/1429	0.53	0/1923
All	All	0.34	0/11839	0.55	1/16022~(0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	В	110	LEU	CA-CB-CG	5.19	127.24	115.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2458	0	2395	14	0
1	С	2469	0	2417	11	0
1	Е	2471	0	2415	13	0
2	В	1404	0	1309	3	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1405	0	1300	4	0
2	F	1404	0	1316	4	0
3	G	28	0	25	0	0
3	L	28	0	25	0	0
3	М	28	0	25	0	0
3	0	28	0	25	0	0
3	Q	28	0	25	0	0
3	R	28	0	25	0	0
3	Т	28	0	25	0	0
4	Н	39	0	34	0	0
4	J	39	0	34	3	0
5	Ι	61	0	52	0	0
5	Ν	61	0	52	0	0
5	S	61	0	52	0	0
6	Κ	60	0	44	0	0
6	Р	60	0	44	0	0
6	U	60	0	44	2	0
7	А	5	0	0	0	0
7	В	5	0	0	0	0
7	С	10	0	0	0	0
7	D	5	0	0	0	0
7	F	5	0	0	0	0
8	С	14	0	13	0	0
9	А	43	0	0	0	1
9	В	12	0	0	0	0
9	С	36	0	0	0	0
9	D	14	0	0	0	0
9	Е	30	0	0	0	0
9	F	3	0	0	0	0
All	All	12430	0	11696	46	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:230:ILE:HD11	1:C:252:VAL:HG21	1.44	0.99
1:A:97:CYS:O	1:A:224:ARG:NH1	2.14	0.80
1:A:285:ASN:HD21	4:J:1:NAG:C1	2.09	0.65
1:A:285:ASN:ND2	4:J:1:NAG:C1	2.62	0.63



		Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:A:180:TRP:CE2	1:A:204:VAL:HG21	2.35	0.60
1:C:301:THR:HB	1:C:305:CYS:SG	2.43	0.59
2:D:25:ARG:HG2	2:D:34:GLN:HG3	1.84	0.58
2:D:110:LEU:C	2:D:110:LEU:HD22	2.24	0.57
1:E:141:ARG:NH1	1:E:146:SER:OG	2.38	0.56
2:F:98:LEU:HD11	2:F:102:LEU:HD23	1.88	0.56
2:F:47:GLN:HB2	2:F:110:LEU:HD11	1.89	0.55
1:C:57:ARG:O	1:C:85:ASP:HB2	2.07	0.54
1:E:97:CYS:O	1:E:224:ARG:NH1	2.41	0.53
2:F:47:GLN:CB	2:F:110:LEU:HD11	2.38	0.53
1:A:8:ASN:C	1:A:9:ASN:HD22	2.12	0.53
1:A:237:VAL:HG21	1:A:243:LEU:HB2	1.92	0.52
1:E:228:GLY:O	1:E:229:ARG:HG2	2.09	0.52
1:E:218:GLY:O	1:E:220:ARG:NH1	2.45	0.50
2:B:54:ARG:NH1	1:E:29:MET:O	2.45	0.50
2:D:51:LYS:HE2	2:D:107:THR:OG1	2.12	0.49
1:A:77:ASP:OD1	1:A:141:ARG:NH1	2.47	0.47
1:C:195:TYR:O	1:C:197:GLN:N	2.47	0.47
1:E:183:HIS:NE2	6:U:3:SIA:O9	2.47	0.47
1:A:126:THR:HG22	1:A:128:THR:HG23	1.96	0.46
1:E:74:PRO:HG2	1:E:139:CYS:HB3	1.97	0.46
1:A:127:TRP:CZ3	1:A:166:VAL:HG21	2.50	0.45
1:C:277:CYS:SG	1:C:278:VAL:N	2.89	0.45
1:C:17:HIS:CD2	2:D:6:ILE:HG12	2.52	0.45
1:C:146:SER:OG	1:C:147:PHE:N	2.47	0.45
2:B:110:LEU:HD22	2:B:110:LEU:C	2.37	0.45
1:E:50:LYS:HG3	1:E:275:ASP:CB	2.47	0.44
1:A:216:ASN:HB3	1:C:212:THR:HG21	1.99	0.44
2:B:107:THR:HA	2:B:110:LEU:CD1	2.48	0.44
1:E:98:TYR:N	1:E:139:CYS:SG	2.91	0.44
1:E:183:HIS:ND1	1:E:195:TYR:OH	2.46	0.44
1:A:297:VAL:HA	4:J:1:NAG:H82	2.01	0.43
1:E:40:THR:HG22	1:E:316:LEU:HB3	2.01	0.43
1:A:127:TRP:CE3	1:A:166:VAL:HG21	2.54	0.42
1:C:50:LYS:HD2	1:C:275:ASP:HB3	2.02	0.41
1:A:201:ARG:HG2	1:A:202:VAL:N	2.34	0.41
1:C:230:ILE:CD1	1:C:252:VAL:HG21	2.33	0.41
6:U:3:SIA:O1B	6:U:3:SIA:H6	2.21	0.41
1:E:10:THR:HG22	2:F:141:TYR:HA	2.03	0.41
1:E:283:THR:HG22	1:E:301:THR:HG22	2.02	0.41
1:C:191:GLN:HG2	1:C:217:ILE:HD11	2.02	0.41



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42:LEU:O	1:A:292:LYS:HB3	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:2011:HOH:O	9:A:2011:HOH:O[3_554]	2.17	0.03

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	317/328~(97%)	304 (96%)	12 (4%)	1 (0%)	41	76
1	С	317/328~(97%)	292 (92%)	24 (8%)	1 (0%)	41	76
1	Е	317/328~(97%)	297~(94%)	18 (6%)	2 (1%)	25	64
2	В	172/175~(98%)	164 (95%)	8 (5%)	0	100	100
2	D	173/175~(99%)	163 (94%)	8 (5%)	2 (1%)	13	48
2	F	172/175~(98%)	158 (92%)	13 (8%)	1 (1%)	25	64
All	All	1468/1509~(97%)	1378 (94%)	83 (6%)	7 (0%)	29	68

All (7) Ramachandran outliers are listed below:

Mol	Chain	$\mathbf{Res}$	Type
1	С	196	ILE
1	А	55	SER
1	Ε	277	CYS
1	Е	304	LYS
2	F	172	GLN
2	D	29	SER
2	D	174	GLY



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	in Analysed Rotameric Outliers		Percer	ntiles	
1	А	276/289~(96%)	258~(94%)	18 (6%)	17	50
1	С	279/289~(96%)	263~(94%)	16 (6%)	20	56
1	Ε	279/289~(96%)	260~(93%)	19 (7%)	16	48
2	В	145/148~(98%)	142 (98%)	3 (2%)	53	82
2	D	144/148~(97%)	139~(96%)	5 (4%)	36	71
2	F	145/148~(98%)	141 (97%)	4 (3%)	43	77
All	All	$1268/1311\ (97\%)$	1203~(95%)	65~(5%)	24	60

All (65) residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	А	14	CYS
1	А	41	GLU
1	А	42	LEU
1	А	57	ARG
1	А	58	ILE
1	А	59	LEU
1	А	70	MET
1	А	97	CYS
1	А	156	LYS
1	А	157	SER
1	А	167	THR
1	А	175	ASP
1	А	205	SER
1	А	222	LEU
1	А	244	MET
1	А	281	CYS
1	А	295	GLN
1	А	301	THR
2	В	110	LEU
2	В	135	ASP
2	В	159	HIS
1	С	14	CYS



Mol	Chain	Res	Type
1	С	18	HIS
1	С	41	GLU
1	С	54	LYS
1	С	59	LEU
1	С	70	MET
1	С	104	ASP
1	С	155	THR
1	С	157	SER
1	С	195	TYR
1	С	219	SER
1	С	230	ILE
1	С	277	CYS
1	С	290	ASN
1	С	295	GLN
1	С	315	LYS
2	D	34	GLN
2	D	110	LEU
2	D	128	GLU
2	D	137	CYS
2	D	159	HIS
1	Е	14	CYS
1	Е	22	ASN
1	Е	40	THR
1	Е	50	LYS
1	Е	54	LYS
1	Е	59	LEU
1	Е	70	MET
1	Е	97	CYS
1	Е	126	THR
1	E	131	THR
1	Е	156	LYS
1	Е	194	LEU
1	E	195	TYR
1	Е	202	VAL
1	Е	211	GLN
1	E	277	CYS
1	E	290	ASN
1	Е	295	GLN
1	E	313	THR
2	F	51	LYS
2	F	126	LEU
2	F	144	CYS



 $Continued \ from \ previous \ page...$ 

Mol	Chain	Res	Type
2	F	159	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	9	ASN
1	А	285	ASN
1	А	290	ASN
1	С	33	GLN
1	С	80	GLN
1	С	296	ASN
1	Е	8	ASN
1	Е	246	ASN
1	Е	290	ASN
1	Е	296	ASN
2	F	116	ASN
2	F	168	ASN
2	F	172	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

47 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mal	Tuno	Chain	Dog	Tink	Bo	ond leng	ths	В	ond ang	les
	туре	Chan	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
3	NAG	G	1	3,1	14,14,15	0.50	0	$17,\!19,\!21$	2.51	5 (29%)
3	NAG	G	2	3	14,14,15	0.93	1 (7%)	$17,\!19,\!21$	1.87	<mark>5 (29%)</mark>
4	NAG	Н	1	4,1	14,14,15	0.53	0	17,19,21	1.29	2 (11%)
4	NAG	Н	2	4	14,14,15	0.69	0	17,19,21	1.25	1 (5%)
4	BMA	Н	3	4	11,11,12	0.39	0	$15,\!15,\!17$	1.04	1 (6%)
5	NAG	Ι	1	5,1	14,14,15	0.61	0	17,19,21	1.38	3 (17%)
5	NAG	Ι	2	5	14,14,15	0.70	0	17,19,21	0.88	0
5	BMA	Ι	3	5	11,11,12	0.30	0	$15,\!15,\!17$	0.80	0
5	MAN	Ι	4	5	11,11,12	0.56	0	$15,\!15,\!17$	0.99	1 (6%)
5	MAN	Ι	5	5	11,11,12	0.74	0	$15,\!15,\!17$	1.71	3 (20%)
4	NAG	J	1	4	14,14,15	0.45	0	17,19,21	1.16	2 (11%)
4	NAG	J	2	4	14,14,15	0.77	0	17,19,21	2.39	6 (35%)
4	BMA	J	3	4	11,11,12	0.42	0	$15,\!15,\!17$	1.12	3 (20%)
6	NGS	K	1	6	19,19,19	0.86	1 (5%)	26,28,28	1.56	5 (19%)
6	GAL	K	2	6	11,11,12	0.77	0	$15,\!15,\!17$	2.46	3 (20%)
6	SIA	K	3	6	20,20,21	0.66	0	24,28,31	1.27	4 (16%)
6	FUC	K	4	6	10,10,11	0.81	0	14,14,16	1.79	2 (14%)
3	NAG	L	1	3,1	14,14,15	0.63	0	17,19,21	1.10	2 (11%)
3	NAG	L	2	3	14,14,15	0.68	0	$17,\!19,\!21$	1.12	1 (5%)
3	NAG	М	1	3,1	14,14,15	0.54	0	17,19,21	1.03	1 (5%)
3	NAG	М	2	3	14,14,15	0.70	0	$17,\!19,\!21$	1.98	5 (29%)
5	NAG	Ν	1	5,1	14,14,15	0.52	0	17,19,21	1.32	3 (17%)
5	NAG	N	2	5	14,14,15	0.56	0	17,19,21	0.86	0
5	BMA	N	3	5	11,11,12	0.44	0	$15,\!15,\!17$	1.05	1 (6%)
5	MAN	Ν	4	5	11,11,12	0.61	0	$15,\!15,\!17$	1.01	1 (6%)
5	MAN	N	5	5	11,11,12	0.53	0	$15,\!15,\!17$	0.93	1 (6%)
3	NAG	0	1	3,1	14,14,15	0.73	0	17,19,21	1.31	2 (11%)
3	NAG	0	2	3	14,14,15	0.51	0	17,19,21	1.06	1 (5%)
6	NGS	Р	1	6	19,19,19	0.84	1 (5%)	26,28,28	1.43	4 (15%)
6	GAL	Р	2	6	11,11,12	0.58	0	15,15,17	1.43	3 (20%)
6	SIA	Р	3	6	20,20,21	0.66	0	24,28,31	1.29	3 (12%)
6	FUC	Р	4	6	10,10,11	0.89	0	14,14,16	1.31	2 (14%)
3	NAG	Q	1	3,1	14,14,15	0.48	0	17,19,21	1.21	1 (5%)
3	NAG	Q	2	3	14,14,15	0.54	0	17,19,21	1.03	1 (5%)
3	NAG	R	1	3,1	14,14,15	0.64	0	17,19,21	0.96	2 (11%)



Mol	Type	Chain	Bos	Link	Bo	Bond lengths		B	ond ang	les	
WIOI	Type	Ullalli	1005	TICS		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
3	NAG	R	2	3	$14,\!14,\!15$	0.69	0	17,19,21	1.54	3 (17%)	
5	NAG	S	1	5,1	$14,\!14,\!15$	0.66	0	17,19,21	1.50	3 (17%)	
5	NAG	S	2	5	$14,\!14,\!15$	0.60	0	17,19,21	0.89	0	
5	BMA	S	3	5	11,11,12	0.31	0	15,15,17	1.89	3 (20%)	
5	MAN	S	4	5	11,11,12	0.64	0	15,15,17	1.50	2 (13%)	
5	MAN	S	5	5	11,11,12	0.61	0	$15,\!15,\!17$	0.87	1 (6%)	
3	NAG	Т	1	3,1	$14,\!14,\!15$	0.65	0	17,19,21	1.11	0	
3	NAG	Т	2	3	$14,\!14,\!15$	0.81	0	17,19,21	1.58	2 (11%)	
6	NGS	U	1	6	19,19,19	0.86	1 (5%)	26,28,28	1.65	4 (15%)	
6	GAL	U	2	6	11,11,12	0.64	0	15,15,17	2.04	2 (13%)	
6	SIA	U	3	6	20,20,21	0.54	0	24,28,31	1.42	4 (16%)	
6	FUC	U	4	6	10,10,11	0.80	0	14,14,16	1.71	2 (14%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	G	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	G	2	3	1/1/5/7	2/6/23/26	0/1/1/1
4	NAG	Н	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	Н	2	4	-	0/6/23/26	0/1/1/1
4	BMA	Н	3	4	-	2/2/19/22	0/1/1/1
5	NAG	Ι	1	5,1	-	2/6/23/26	0/1/1/1
5	NAG	Ι	2	5	-	0/6/23/26	0/1/1/1
5	BMA	Ι	3	5	-	0/2/19/22	0/1/1/1
5	MAN	Ι	4	5	-	1/2/19/22	0/1/1/1
5	MAN	Ι	5	5	-	2/2/19/22	0/1/1/1
4	NAG	J	1	4	-	1/6/23/26	0/1/1/1
4	NAG	J	2	4	-	4/6/23/26	0/1/1/1
4	BMA	J	3	4	-	2/2/19/22	0/1/1/1
6	NGS	K	1	6	-	5/10/30/30	0/1/1/1
6	GAL	К	2	6	1/1/4/5	0/2/19/22	0/1/1/1
6	SIA	К	3	6	-	2/18/34/38	0/1/1/1
6	FUC	Κ	4	6	1/1/4/5	-	0/1/1/1
3	NAG	L	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	L	2	3	-	2/6/23/26	0/1/1/1



4	U	O	7

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	М	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	М	2	3	-	4/6/23/26	0/1/1/1
5	NAG	N	1	5,1	-	2/6/23/26	0/1/1/1
5	NAG	N	2	5	-	0/6/23/26	0/1/1/1
5	BMA	N	3	5	-	2/2/19/22	0/1/1/1
5	MAN	N	4	5	-	1/2/19/22	1/1/1/1
5	MAN	N	5	5	-	2/2/19/22	1/1/1/1
3	NAG	0	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	0	2	3	-	1/6/23/26	0/1/1/1
6	NGS	Р	1	6	-	1/10/30/30	0/1/1/1
6	GAL	Р	2	6	-	1/2/19/22	0/1/1/1
6	SIA	Р	3	6	-	2/18/34/38	0/1/1/1
6	FUC	Р	4	6	1/1/4/5	-	0/1/1/1
3	NAG	Q	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	Q	2	3	-	0/6/23/26	0/1/1/1
3	NAG	R	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	R	2	3	-	1/6/23/26	0/1/1/1
5	NAG	S	1	5,1	-	4/6/23/26	0/1/1/1
5	NAG	S	2	5	-	0/6/23/26	0/1/1/1
5	BMA	S	3	5	-	2/2/19/22	1/1/1/1
5	MAN	S	4	5	-	2/2/19/22	0/1/1/1
5	MAN	S	5	5	-	1/2/19/22	0/1/1/1
3	NAG	Т	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	Т	2	3	-	1/6/23/26	0/1/1/1
6	NGS	U	1	6	-	4/10/30/30	0/1/1/1
6	GAL	U	2	6	-	2/2/19/22	0/1/1/1
6	SIA	U	3	6	-	4/18/34/38	0/1/1/1
6	FUC	U	4	6	1/1/4/5	-	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	2	NAG	O5-C1	-2.99	1.38	1.43
6	U	1	NGS	08-S	2.27	1.64	1.50
6	Р	1	NGS	08-S	2.26	1.64	1.50
6	Κ	1	NGS	08-S	2.25	1.64	1.50

All (106) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	G	1	NAG	C1-O5-C5	6.62	121.16	112.19
6	Κ	2	GAL	O5-C1-C2	6.20	120.34	110.77
6	U	2	GAL	C1-C2-C3	6.18	117.26	109.67
4	J	2	NAG	C2-N2-C7	5.73	131.06	122.90
6	Κ	2	GAL	C1-C2-C3	5.57	116.51	109.67
5	S	3	BMA	C1-O5-C5	5.56	119.72	112.19
6	U	4	FUC	C1-C2-C3	-5.02	103.49	109.67
6	U	1	NGS	C1-C2-C3	4.96	117.31	110.54
5	Ι	5	MAN	C1-C2-C3	4.87	115.65	109.67
3	Т	2	NAG	C4-C3-C2	4.76	118.00	111.02
3	G	2	NAG	C1-O5-C5	-4.68	105.85	112.19
6	Κ	4	FUC	C1-C2-C3	-4.43	104.22	109.67
4	J	2	NAG	C8-C7-N2	4.41	123.56	116.10
3	G	1	NAG	O4-C4-C3	4.39	120.49	110.35
3	G	1	NAG	O5-C1-C2	4.37	118.19	111.29
5	S	4	MAN	C1-C2-C3	4.30	114.96	109.67
6	Κ	1	NGS	C4-C3-C2	4.15	116.42	110.34
3	Q	1	NAG	C1-O5-C5	3.97	117.57	112.19
6	Κ	4	FUC	O5-C1-C2	3.93	116.83	110.77
3	G	1	NAG	C4-C3-C2	-3.87	105.34	111.02
3	М	2	NAG	C8-C7-N2	3.82	122.57	116.10
3	М	2	NAG	C1-O5-C5	3.78	117.31	112.19
4	J	2	NAG	C4-C3-C2	3.69	116.42	111.02
6	U	2	GAL	O5-C5-C6	3.61	112.86	107.20
6	U	1	NGS	C4-C3-C2	3.58	115.59	110.34
6	U	3	SIA	O6-C2-C3	-3.57	105.54	110.46
3	0	1	NAG	C4-C3-C2	3.57	116.25	111.02
6	Р	2	GAL	C1-C2-C3	3.56	114.04	109.67
4	Н	2	NAG	C4-C3-C2	3.51	116.16	111.02
4	Н	3	BMA	C1-O5-C5	3.49	116.92	112.19
6	Р	3	SIA	C6-O6-C2	3.44	118.69	111.34
5	Ν	1	NAG	C8-C7-N2	3.43	121.90	116.10
6	Κ	1	NGS	O5-C5-C4	3.41	115.89	109.69
6	Р	1	NGS	C4-C3-C2	3.40	115.32	110.34
3	М	2	NAG	C2-N2-C7	3.38	127.72	122.90
5	S	1	NAG	O5-C1-C2	-3.32	106.05	111.29
4	J	1	NAG	C1-O5-C5	3.25	116.59	112.19
4	J	2	NAG	C1-C2-N2	3.25	116.03	110.49
5	S	1	NAG	C8-C7-N2	3.24	121.58	116.10
6	Р	1	NGS	O6-C6-C5	3.20	113.60	107.62
3	R	2	NAG	C2-N2-C7	3.20	127.46	122.90
3	G	2	NAG	C4-C3-C2	-3.19	106.35	111.02
3	М	2	NAG	C4-C3-C2	3.19	115.69	111.02



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	Ι	1	NAG	C8-C7-N2	3.18	121.49	116.10
6	Κ	2	GAL	O5-C5-C6	3.13	112.10	107.20
5	Ι	5	MAN	C2-C3-C4	3.08	116.23	110.89
3	L	2	NAG	C4-C3-C2	3.07	115.52	111.02
3	L	1	NAG	C1-C2-N2	-3.07	105.24	110.49
4	J	2	NAG	O7-C7-C8	-3.03	116.42	122.06
5	Ν	4	MAN	C1-O5-C5	2.92	116.15	112.19
3	R	2	NAG	C4-C3-C2	2.87	115.22	111.02
3	R	2	NAG	C1-C2-N2	2.81	115.28	110.49
6	Κ	1	NGS	O5-C1-C2	-2.80	106.70	109.52
6	U	1	NGS	C1-C2-N2	-2.75	107.54	110.73
3	0	1	NAG	C3-C4-C5	2.73	115.10	110.24
5	Ι	4	MAN	C1-O5-C5	2.69	115.84	112.19
3	G	2	NAG	C1-C2-N2	2.67	115.06	110.49
3	G	2	NAG	O5-C5-C6	2.65	111.36	107.20
5	S	3	BMA	C1-C2-C3	2.64	112.92	109.67
6	U	4	FUC	O5-C1-C2	2.62	114.82	110.77
3	М	1	NAG	C4-C3-C2	2.62	114.86	111.02
6	Р	2	GAL	O5-C5-C6	2.61	111.30	107.20
4	Н	1	NAG	C1-C2-N2	-2.61	106.03	110.49
5	Ν	3	BMA	O5-C1-C2	-2.54	106.85	110.77
5	Ν	5	MAN	O5-C5-C6	2.53	111.18	107.20
5	Ι	1	NAG	C1-O5-C5	2.53	115.61	112.19
6	Р	1	NGS	C3-C4-C5	2.51	114.72	110.24
3	R	1	NAG	C4-C3-C2	2.50	114.68	111.02
6	U	1	NGS	O5-C1-C2	2.46	111.98	109.52
4	J	2	NAG	C1-O5-C5	2.41	115.46	112.19
6	Р	1	NGS	O5-C5-C4	2.41	114.06	109.69
6	Р	3	SIA	O1B-C1-C2	2.40	119.89	113.03
4	J	3	BMA	C1-C2-C3	2.39	112.60	109.67
3	Q	2	NAG	C4-C3-C2	2.38	114.51	111.02
6	Κ	1	NGS	O5-C5-C6	-2.35	101.91	106.67
6	U	3	SIA	O1B-C1-C2	2.35	119.73	113.03
5	S	1	NAG	O7-C7-C8	-2.34	117.70	122.06
6	Κ	1	NGS	O3-C3-C2	-2.34	104.93	109.66
5	Ν	1	NAG	O7-C7-C8	-2.33	117.72	122.06
3	М	2	NAG	O7-C7-C8	-2.33	117.72	122.06
6	Р	4	FUC	C3-C4-C5	2.32	113.39	109.77
5	S	4	MAN	C1-O5-C5	2.32	115.34	112.19
6	Κ	3	SIA	C11-C10-N5	2.32	120.03	116.10
6	Р	2	GAL	O3-C3-C2	-2.31	105.57	109.99
6	Κ	3	SIA	O1B-C1-C2	2.31	119.61	113.03



Mol	Chain	$\mathbf{Res}$	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
6	Κ	3	SIA	C6-O6-C2	2.28	116.22	111.34
6	U	3	SIA	C11-C10-N5	2.23	119.87	116.10
6	Р	4	FUC	C1-O5-C5	-2.21	107.77	112.78
5	S	3	BMA	C3-C4-C5	2.16	114.08	110.24
4	J	3	BMA	O5-C5-C6	2.14	110.56	107.20
3	0	2	NAG	C4-C3-C2	2.12	114.12	111.02
3	Т	2	NAG	C3-C4-C5	2.10	113.99	110.24
3	L	1	NAG	C4-C3-C2	2.10	114.09	111.02
6	Р	3	SIA	C3-C4-C5	-2.08	108.94	111.46
3	G	1	NAG	C1-C2-N2	-2.05	106.98	110.49
4	Н	1	NAG	O5-C5-C4	-2.05	105.84	110.83
5	N	1	NAG	O5-C1-C2	-2.05	108.05	111.29
3	R	1	NAG	O5-C5-C4	-2.05	105.84	110.83
4	J	1	NAG	O5-C5-C6	2.04	110.40	107.20
6	U	3	SIA	C6-O6-C2	2.03	115.69	111.34
5	Ι	5	MAN	C3-C4-C5	2.03	113.86	110.24
5	Ι	1	NAG	O7-C7-C8	-2.03	118.29	122.06
6	К	3	SIA	C6-C5-N5	-2.02	107.55	110.91
3	G	2	NAG	O5-C1-C2	2.02	114.48	111.29
5	S	5	MAN	C3-C4-C5	2.01	113.83	110.24
4	J	3	BMA	C1-O5-C5	2.01	114.91	112.19

All (5) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	G	2	NAG	C1
6	Κ	2	GAL	C1
6	Κ	4	FUC	C1
6	Р	4	FUC	C1
6	U	4	FUC	C1

All (66) torsion outliers are listed below:

Mol	Chain	$\operatorname{Res}$	Type	Atoms
6	U	1	NGS	C6-O6-S-O7A
6	U	1	NGS	C6-O6-S-O8
6	U	1	NGS	C6-O6-S-O9
6	U	3	SIA	C7-C8-C9-O9
6	U	3	SIA	O8-C8-C9-O9
6	U	2	GAL	O5-C5-C6-O6
5	S	3	BMA	O5-C5-C6-O6
5	S	4	MAN	C4-C5-C6-O6



Mol	Chain	Res	Type	Atoms
4	Н	3	BMA	O5-C5-C6-O6
5	Ν	5	MAN	O5-C5-C6-O6
6	U	2	GAL	C4-C5-C6-O6
3	G	2	NAG	O5-C5-C6-O6
4	J	3	BMA	O5-C5-C6-O6
5	S	3	BMA	C4-C5-C6-O6
3	R	2	NAG	C1-C2-N2-C7
5	Ι	5	MAN	C4-C5-C6-O6
4	J	3	BMA	C4-C5-C6-O6
3	М	2	NAG	C8-C7-N2-C2
3	М	2	NAG	O7-C7-N2-C2
4	J	2	NAG	C8-C7-N2-C2
4	J	2	NAG	O7-C7-N2-C2
5	Ι	1	NAG	C8-C7-N2-C2
5	Ι	1	NAG	O7-C7-N2-C2
5	N	1	NAG	C8-C7-N2-C2
5	N	1	NAG	O7-C7-N2-C2
5	S	1	NAG	C8-C7-N2-C2
5	S	1	NAG	O7-C7-N2-C2
6	Κ	3	SIA	C11-C10-N5-C5
6	Κ	3	SIA	O10-C10-N5-C5
6	Р	3	SIA	C11-C10-N5-C5
6	Р	3	SIA	O10-C10-N5-C5
6	U	3	SIA	C11-C10-N5-C5
6	U	3	SIA	O10-C10-N5-C5
3	G	2	NAG	C4-C5-C6-O6
5	N	3	BMA	O5-C5-C6-O6
5	S	4	MAN	O5-C5-C6-O6
5	N	3	BMA	C4-C5-C6-O6
5	I	5	MAN	O5-C5-C6-O6
5	N	5	MAN	C4-C5-C6-O6
4	Н	3	BMA	C4-C5-C6-O6
3	Q	1	NAG	O5-C5-C6-O6
3	Q	1	NAG	C4-C5-C6-O6
6	K	1	NGS	C4-C5-C6-O6
3	0	2	NAG	O5-C5-C6-O6
5	N	4	MAN	O5-C5-C6-O6
3	Т	2	NAG	O5-C5-C6-O6
4	J	2	NAG	O5-C5-C6-O6
3	М	2	NAG	C4-C5-C6-O6
4	J	1	NAG	O5-C5-C6-O6
4	J	2	NAG	C1-C2-N2-C7



4	U	Ο	7

Mol	Chain	Res	Type	Atoms
6	K	1	NGS	C6-O6-S-O7A
6	K	1	NGS	C6-O6-S-O9
6	K	1	NGS	O5-C5-C6-O6
5	Ι	4	MAN	C4-C5-C6-O6
3	L	2	NAG	C4-C5-C6-O6
5	S	5	MAN	O5-C5-C6-O6
3	L	2	NAG	O5-C5-C6-O6
3	М	2	NAG	O5-C5-C6-O6
6	Р	2	GAL	C4-C5-C6-O6
6	K	1	NGS	C6-O6-S-O8
6	Р	1	NGS	C6-O6-S-O8
5	S	1	NAG	C4-C5-C6-O6
3	L	1	NAG	O5-C5-C6-O6
3	L	1	NAG	C4-C5-C6-O6
5	S	1	NAG	O5-C5-C6-O6
6	U	1	NGS	O5-C5-C6-O6

All (3) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	S	3	BMA	C1-C2-C3-C4-C5-O5
5	Ν	4	MAN	C1-C2-C3-C4-C5-O5
5	Ν	5	MAN	C1-C2-C3-C4-C5-O5

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	U	3	SIA	2	0
4	J	1	NAG	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

























































## 5.6 Ligand geometry (i)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mal Tura		Chain	Chain	Bos	Tink	Bo	Bond lengths			Bond angles		
1VIOI	Moi Type Ci	Unam	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2		
7	SO4	В	1175	-	$4,\!4,\!4$	0.35	0	$6,\!6,\!6$	0.22	0		
7	SO4	С	1328	-	4,4,4	0.33	0	$6,\!6,\!6$	0.07	0		
7	SO4	F	1175	-	$4,\!4,\!4$	0.35	0	$6,\!6,\!6$	0.14	0		
8	NAG	С	601	1	$14,\!14,\!15$	0.42	0	$17,\!19,\!21$	2.08	1 (5%)		
7	SO4	D	1176	-	4,4,4	0.37	0	6,6,6	0.14	0		
7	SO4	С	1327	-	4,4,4	0.34	0	6,6,6	0.13	0		
7	SO4	А	1327	-	4,4,4	0.36	0	$6,\!6,\!6$	0.09	0		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	С	601	1	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
8	С	601	NAG	C1-O5-C5	7.69	122.61	112.19

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	$\operatorname{Res}$	Type	Atoms
8	С	601	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	319/328~(97%)	-0.28	2 (0%) 89 72	41, 63, 84, 129	0
1	С	319/328~(97%)	-0.28	1 (0%) 94 84	40, 63, 87, 126	0
1	Ε	319/328~(97%)	-0.28	0 100 100	41, 63, 85, 127	0
2	В	174/175~(99%)	-0.11	2 (1%) 80 56	43, 79, 97, 116	0
2	D	175/175~(100%)	-0.18	2 (1%) 80 56	40, 74, 96, 130	0
2	F	174/175~(99%)	-0.22	0 100 100	40, 75, 99, 108	0
All	All	1480/1509~(98%)	-0.24	7 (0%) 91 75	40, 67, 95, 130	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	11	ALA	2.6
2	В	174	GLY	2.4
2	D	173	SER	2.3
2	В	173	SER	2.3
2	D	175	ARG	2.1
1	А	326	LYS	2.1
1	С	326	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
3	NAG	Т	2	14/15	0.56	0.51	90,109,112,113	0
5	MAN	N	4	11/12	0.63	0.32	104,112,119,122	0
3	NAG	Q	2	14/15	0.64	0.62	125,130,133,134	0
4	BMA	Н	3	11/12	0.67	0.36	118,126,129,130	0
4	BMA	J	3	11/12	0.73	0.39	127,133,138,139	0
3	NAG	L	2	14/15	0.73	0.41	105,113,120,123	0
5	MAN	S	4	11/12	0.74	0.47	123,128,133,135	0
5	MAN	S	5	11/12	0.79	0.22	113,118,123,123	0
5	MAN	N	5	11/12	0.80	0.33	115,117,123,124	0
3	NAG	Т	1	14/15	0.81	0.35	84,91,96,102	0
4	NAG	Н	2	14/15	0.82	0.34	112,122,128,129	0
3	NAG	G	2	14/15	0.82	0.26	110,118,127,127	0
5	BMA	S	3	11/12	0.82	0.27	102,110,115,118	0
4	NAG	J	2	14/15	0.82	0.43	110,121,129,130	0
3	NAG	R	1	14/15	0.82	0.22	92,96,102,104	0
3	NAG	0	2	14/15	0.83	0.38	89,100,106,108	0
4	NAG	J	1	14/15	0.83	0.33	93,98,104,111	0
3	NAG	G	1	14/15	0.83	0.23	85,92,101,109	0
3	NAG	L	1	14/15	0.84	0.28	88,94,101,106	0
5	BMA	Ι	3	11/12	0.85	0.17	94,99,106,106	0
3	NAG	М	2	14/15	0.85	0.22	90,97,99,100	0
3	NAG	0	1	14/15	0.85	0.31	76,82,89,95	0
4	NAG	Н	1	14/15	0.86	0.19	89,94,101,109	0
3	NAG	Q	1	14/15	0.86	0.33	95,101,110,119	0
3	NAG	М	1	14/15	0.87	0.22	84,89,92,93	0
3	NAG	R	2	14/15	0.87	0.31	95,106,110,110	0
6	FUC	P	4	10/11	0.87	0.34	95,99,100,101	0
6	NGS	Р	1	19/19	0.88	0.29	97,102,111,113	0
5	MAN	I	4	11/12	0.88	0.28	105,110,116,116	0
6	NGS	K	1	19/19	0.89	0.23	82,89,99,102	0
6	FUC	K	4	10/11	0.89	0.30	89,90,91,92	0
5	BMA	N	3	11/12	0.89	0.19	95,101,106,108	0
5	MAN	I	5	11/12	0.89	0.19	100,107,113,116	0
6	GAL	K	2	11/12	0.91	0.24	77,80,84,87	0
6	NGS	U	1	19/19	0.91	0.19	89,96,106,107	0
5	NAG	Ι	2	14/15	0.92	0.20	73,79,86,89	0
6	GAL	P	2	11/12	0.92	0.24	83,86,91,92	0
6	SIA	U	3	20/21	0.92	0.24	67,72,81,84	0
6	GAL	U	2	11/12	0.93	0.18	80,83,87,87	0
5	NAG	N	2	14/15	0.93	0.18	69,79,86,90	0
5	NAG	S	1	14/15	0.94	0.21	67,70,74,74	0
6	SIA	K	3	20/21	0.94	0.30	$66,\!68,\!77,\!79$	0

labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	$Q{<}0.9$				
5	NAG	S	2	14/15	0.94	0.16	72,79,86,93	0				
6	FUC	U	4	10/11	0.94	0.24	84,88,92,93	0				
6	SIA	Р	3	20/21	0.95	0.24	68,72,78,79	0				
5	NAG	Ι	1	14/15	0.95	0.15	69,73,75,77	0				
5	NAG	N	1	14/15	0.96	0.14	71,75,76,77	0				

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







































## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
7	SO4	С	1328	5/5	0.66	0.40	137,138,140,141	0
7	SO4	С	1327	5/5	0.81	0.27	120,121,124,125	0
7	SO4	А	1327	5/5	0.85	0.22	130,131,134,135	0
8	NAG	С	601	14/15	0.86	0.21	76,86,91,92	0
7	SO4	В	1175	5/5	0.90	0.21	85,89,90,96	0
7	SO4	D	1176	5/5	0.94	0.18	78,79,80,83	0
7	SO4	F	1175	5/5	0.95	0.15	85,85,89,90	0

## 6.5 Other polymers (i)

There are no such residues in this entry.

