



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 15, 2023 – 02:39 AM EDT

PDB ID : 7US3
Title : Structure of Putrescine N-hydroxylase Involved Complexed with NADP+
Authors : Tanner, J.J.; Bogner, A.N.
Deposited on : 2022-04-22
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	466	85% 7% 8%
1	B	466	82% 9% 8%
1	C	466	82% 9% 8%
1	D	466	80% 12% 8%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 14794 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putrescine N-hydroxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	428	3443	2224	563	644	12	0	0	0
1	B	428	3437	2222	564	639	12	0	0	0
1	C	428	3444	2223	567	642	12	0	0	0
1	D	428	3430	2217	558	643	12	0	0	0

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP A0A1E3MAZ6
A	-18	GLY	-	expression tag	UNP A0A1E3MAZ6
A	-17	SER	-	expression tag	UNP A0A1E3MAZ6
A	-16	SER	-	expression tag	UNP A0A1E3MAZ6
A	-15	HIS	-	expression tag	UNP A0A1E3MAZ6
A	-14	HIS	-	expression tag	UNP A0A1E3MAZ6
A	-13	HIS	-	expression tag	UNP A0A1E3MAZ6
A	-12	HIS	-	expression tag	UNP A0A1E3MAZ6
A	-11	HIS	-	expression tag	UNP A0A1E3MAZ6
A	-10	HIS	-	expression tag	UNP A0A1E3MAZ6
A	-9	SER	-	expression tag	UNP A0A1E3MAZ6
A	-8	SER	-	expression tag	UNP A0A1E3MAZ6
A	-7	GLY	-	expression tag	UNP A0A1E3MAZ6
A	-6	LEU	-	expression tag	UNP A0A1E3MAZ6
A	-5	VAL	-	expression tag	UNP A0A1E3MAZ6
A	-4	PRO	-	expression tag	UNP A0A1E3MAZ6
A	-3	ARG	-	expression tag	UNP A0A1E3MAZ6
A	-2	GLY	-	expression tag	UNP A0A1E3MAZ6
A	-1	SER	-	expression tag	UNP A0A1E3MAZ6
A	0	HIS	-	expression tag	UNP A0A1E3MAZ6
B	-19	MET	-	initiating methionine	UNP A0A1E3MAZ6

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-18	GLY	-	expression tag	UNP A0A1E3MAZ6
B	-17	SER	-	expression tag	UNP A0A1E3MAZ6
B	-16	SER	-	expression tag	UNP A0A1E3MAZ6
B	-15	HIS	-	expression tag	UNP A0A1E3MAZ6
B	-14	HIS	-	expression tag	UNP A0A1E3MAZ6
B	-13	HIS	-	expression tag	UNP A0A1E3MAZ6
B	-12	HIS	-	expression tag	UNP A0A1E3MAZ6
B	-11	HIS	-	expression tag	UNP A0A1E3MAZ6
B	-10	HIS	-	expression tag	UNP A0A1E3MAZ6
B	-9	SER	-	expression tag	UNP A0A1E3MAZ6
B	-8	SER	-	expression tag	UNP A0A1E3MAZ6
B	-7	GLY	-	expression tag	UNP A0A1E3MAZ6
B	-6	LEU	-	expression tag	UNP A0A1E3MAZ6
B	-5	VAL	-	expression tag	UNP A0A1E3MAZ6
B	-4	PRO	-	expression tag	UNP A0A1E3MAZ6
B	-3	ARG	-	expression tag	UNP A0A1E3MAZ6
B	-2	GLY	-	expression tag	UNP A0A1E3MAZ6
B	-1	SER	-	expression tag	UNP A0A1E3MAZ6
B	0	HIS	-	expression tag	UNP A0A1E3MAZ6
C	-19	MET	-	initiating methionine	UNP A0A1E3MAZ6
C	-18	GLY	-	expression tag	UNP A0A1E3MAZ6
C	-17	SER	-	expression tag	UNP A0A1E3MAZ6
C	-16	SER	-	expression tag	UNP A0A1E3MAZ6
C	-15	HIS	-	expression tag	UNP A0A1E3MAZ6
C	-14	HIS	-	expression tag	UNP A0A1E3MAZ6
C	-13	HIS	-	expression tag	UNP A0A1E3MAZ6
C	-12	HIS	-	expression tag	UNP A0A1E3MAZ6
C	-11	HIS	-	expression tag	UNP A0A1E3MAZ6
C	-10	HIS	-	expression tag	UNP A0A1E3MAZ6
C	-9	SER	-	expression tag	UNP A0A1E3MAZ6
C	-8	SER	-	expression tag	UNP A0A1E3MAZ6
C	-7	GLY	-	expression tag	UNP A0A1E3MAZ6
C	-6	LEU	-	expression tag	UNP A0A1E3MAZ6
C	-5	VAL	-	expression tag	UNP A0A1E3MAZ6
C	-4	PRO	-	expression tag	UNP A0A1E3MAZ6
C	-3	ARG	-	expression tag	UNP A0A1E3MAZ6
C	-2	GLY	-	expression tag	UNP A0A1E3MAZ6
C	-1	SER	-	expression tag	UNP A0A1E3MAZ6
C	0	HIS	-	expression tag	UNP A0A1E3MAZ6
D	-19	MET	-	initiating methionine	UNP A0A1E3MAZ6
D	-18	GLY	-	expression tag	UNP A0A1E3MAZ6
D	-17	SER	-	expression tag	UNP A0A1E3MAZ6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-16	SER	-	expression tag	UNP A0A1E3MAZ6
D	-15	HIS	-	expression tag	UNP A0A1E3MAZ6
D	-14	HIS	-	expression tag	UNP A0A1E3MAZ6
D	-13	HIS	-	expression tag	UNP A0A1E3MAZ6
D	-12	HIS	-	expression tag	UNP A0A1E3MAZ6
D	-11	HIS	-	expression tag	UNP A0A1E3MAZ6
D	-10	HIS	-	expression tag	UNP A0A1E3MAZ6
D	-9	SER	-	expression tag	UNP A0A1E3MAZ6
D	-8	SER	-	expression tag	UNP A0A1E3MAZ6
D	-7	GLY	-	expression tag	UNP A0A1E3MAZ6
D	-6	LEU	-	expression tag	UNP A0A1E3MAZ6
D	-5	VAL	-	expression tag	UNP A0A1E3MAZ6
D	-4	PRO	-	expression tag	UNP A0A1E3MAZ6
D	-3	ARG	-	expression tag	UNP A0A1E3MAZ6
D	-2	GLY	-	expression tag	UNP A0A1E3MAZ6
D	-1	SER	-	expression tag	UNP A0A1E3MAZ6
D	0	HIS	-	expression tag	UNP A0A1E3MAZ6

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

Continued on next page...



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		
4	D	1	Total	C	O	0	0
			10	6	4		


- Molecule 5 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			13	8	5		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	160	Total 160	O 160	0	0
6	B	143	Total 143	O 143	0	0
6	C	152	Total 152	O 152	0	0
6	D	148	Total 148	O 148	0	0



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	124.83Å 126.31Å 140.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	88.79 – 2.20 93.86 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.2 (88.79-2.20) 99.2 (93.86-2.20)	Depositor EDS
R_{merge}	0.25	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.38 (at 2.20Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.182 , 0.223 0.182 , 0.223	Depositor DCC
R_{free} test set	5591 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	39.1	Xtrriage
Anisotropy	0.319	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 28.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.000 for k,h,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14794	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, FAD, PG4, NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/3532	0.57	0/4802
1	B	0.40	0/3526	0.57	0/4795
1	C	0.40	0/3533	0.57	0/4804
1	D	0.39	0/3519	0.55	0/4788
All	All	0.40	0/14110	0.57	0/19189

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3443	0	3326	19	0
1	B	3437	0	3324	21	0
1	C	3444	0	3328	24	0
1	D	3430	0	3300	28	0
2	A	53	0	31	3	0
2	B	53	0	31	0	0
2	C	53	0	31	1	0
2	D	53	0	31	0	0
3	A	48	0	23	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	48	0	24	1	0
3	C	48	0	24	0	0
3	D	48	0	24	2	0
4	A	10	0	14	0	0
4	D	10	0	14	0	0
5	C	13	0	18	3	0
6	A	160	0	0	1	0
6	B	143	0	0	0	0
6	C	152	0	0	1	0
6	D	148	0	0	0	0
All	All	14794	0	13543	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 86 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:39:LYS:NZ	2:A:501:FAD:O3B	2.20	0.74
1:B:308:SER:OG	1:B:309:ASN:N	2.25	0.68
1:C:56:GLN:HE21	1:C:271:GLU:HB2	1.58	0.67
1:B:235:LEU:HD13	1:D:295:MET:HE1	1.82	0.62
1:C:371:ASP:HB3	1:C:373:THR:HG23	1.84	0.59

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	426/466 (91%)	413 (97%)	12 (3%)	1 (0%)	47	55
1	B	426/466 (91%)	412 (97%)	14 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	426/466 (91%)	412 (97%)	14 (3%)	0	100	100
1	D	426/466 (91%)	410 (96%)	15 (4%)	1 (0%)	47	55
All	All	1704/1864 (91%)	1647 (97%)	55 (3%)	2 (0%)	51	60

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	308	SER
1	A	308	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	372/419 (89%)	362 (97%)	10 (3%)	44	57
1	B	370/419 (88%)	361 (98%)	9 (2%)	49	62
1	C	372/419 (89%)	360 (97%)	12 (3%)	39	50
1	D	369/419 (88%)	355 (96%)	14 (4%)	33	42
All	All	1483/1676 (88%)	1438 (97%)	45 (3%)	41	53

5 of 45 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	383	SER
1	D	321	ASN
1	C	409	GLU
1	D	223	LYS
1	D	351	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	GLN
1	C	56	GLN
1	C	141	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PGE	A	503	-	9,9,9	0.53	0	8,8,8	0.29	0
2	FAD	B	502	-	53,58,58	2.07	14 (26%)	68,89,89	1.46	11 (16%)
3	NAP	A	502	-	45,52,52	4.32	15 (33%)	56,80,80	1.94	9 (16%)
3	NAP	C	502	-	45,52,52	4.33	14 (31%)	56,80,80	1.98	9 (16%)
4	PGE	D	501	-	9,9,9	0.53	0	8,8,8	0.45	0
2	FAD	D	503	-	53,58,58	2.06	16 (30%)	68,89,89	1.43	7 (10%)
2	FAD	C	503	-	53,58,58	2.04	13 (24%)	68,89,89	1.41	8 (11%)
2	FAD	A	501	-	53,58,58	2.14	15 (28%)	68,89,89	1.36	9 (13%)
5	PG4	C	501	-	12,12,12	0.53	0	11,11,11	0.30	0
3	NAP	B	501	-	45,52,52	4.36	15 (33%)	56,80,80	1.89	9 (16%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAP	D	502	-	45,52,52	4.39	15 (33%)	56,80,80	1.94	7 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PGE	A	503	-	-	5/7/7/7	-
2	FAD	B	502	-	-	2/30/50/50	0/6/6/6
3	NAP	A	502	-	-	9/31/67/67	0/5/5/5
3	NAP	C	502	-	-	9/31/67/67	0/5/5/5
4	PGE	D	501	-	-	4/7/7/7	-
2	FAD	D	503	-	-	2/30/50/50	0/6/6/6
2	FAD	C	503	-	-	3/30/50/50	0/6/6/6
2	FAD	A	501	-	-	2/30/50/50	0/6/6/6
5	PG4	C	501	-	-	4/10/10/10	-
3	NAP	B	501	-	-	8/31/67/67	0/5/5/5
3	NAP	D	502	-	-	9/31/67/67	0/5/5/5

The worst 5 of 117 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	502	NAP	O4D-C1D	15.48	1.62	1.41
3	B	501	NAP	O4D-C1D	15.38	1.62	1.41
3	A	502	NAP	O4D-C1D	15.24	1.62	1.41
3	D	502	NAP	O4B-C1B	14.71	1.61	1.41
3	C	502	NAP	O4D-C1D	14.70	1.61	1.41

The worst 5 of 69 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	502	NAP	C5A-C6A-N6A	9.32	134.51	120.35
3	D	502	NAP	C5A-C6A-N6A	9.03	134.07	120.35
3	A	502	NAP	C5A-C6A-N6A	8.55	133.35	120.35
3	B	501	NAP	C5A-C6A-N6A	8.47	133.22	120.35
3	D	502	NAP	N6A-C6A-N1A	-6.33	105.44	118.57

There are no chirality outliers.

5 of 57 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	FAD	O4B-C4B-C5B-O5B
2	C	503	FAD	O4B-C4B-C5B-O5B
3	A	502	NAP	C2B-O2B-P2B-O1X
3	B	501	NAP	C2B-O2B-P2B-O1X
3	B	501	NAP	C2N-C3N-C7N-O7N


There are no ring outliers.


5 monomers are involved in 10 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	503	FAD	1	0
2	A	501	FAD	3	0
5	C	501	PG4	3	0
3	B	501	NAP	1	0
3	D	502	NAP	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	428/466 (91%)	-0.33	1 (0%) 95 94	29, 35, 50, 79	0
1	B	428/466 (91%)	-0.34	0 100 100	29, 36, 50, 74	0
1	C	428/466 (91%)	-0.32	0 100 100	29, 37, 51, 73	0
1	D	428/466 (91%)	-0.32	0 100 100	29, 39, 53, 71	0
All	All	1712/1864 (91%)	-0.33	1 (0%) 95 95	29, 37, 51, 79	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	3	SER	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PGE	D	501	10/10	0.77	0.21	52,59,61,63	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	PG4	C	501	13/13	0.86	0.19	53,55,60,61	0
4	PGE	A	503	10/10	0.89	0.22	49,54,55,55	0
3	NAP	B	501	48/48	0.95	0.12	35,40,46,55	0
3	NAP	D	502	48/48	0.96	0.11	38,43,48,58	0
3	NAP	A	502	48/48	0.96	0.11	33,38,44,52	0
2	FAD	C	503	53/53	0.96	0.13	31,37,41,42	0
3	NAP	C	502	48/48	0.96	0.13	33,41,48,59	0
2	FAD	A	501	53/53	0.97	0.12	29,36,39,43	0
2	FAD	D	503	53/53	0.97	0.12	29,38,42,47	0
2	FAD	B	502	53/53	0.97	0.12	30,35,37,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around NAP D 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAP A 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD C 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAP C 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FAD A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD D 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.