



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 13, 2020 – 12:58 am BST

PDB ID : 3UTS  
Title : 1E6-A\*0201-ALWGPDPA AAA Complex, Monoclinic  
Authors : Rizkallah, P.J.; Cole, D.K.; Sewell, A.K.; Bulek, A.M.  
Deposited on : 2011-11-26  
Resolution : 2.71 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11

# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:  
*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3359 (2.74-2.70)
Clashscore	141614	3686 (2.74-2.70)
Ramachandran outliers	138981	3622 (2.74-2.70)
Sidechain outliers	138945	3623 (2.74-2.70)
RSRZ outliers	127900	3276 (2.74-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	
1	F	276	
2	B	100	
2	G	100	
3	C	10	
3	H	10	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	D	201	 <p>69% 27%</p>
4	I	201	 <p>69% 28%</p>
5	E	246	 <p>69% 29%</p>
5	J	246	 <p>76% 21%</p>

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 13552 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, A-2 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	276	Total	C	N	O	S	0	0	0
			2253	1408	410	426	9			
1	F	276	Total	C	N	O	S	0	1	0
			2264	1414	414	427	9			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			
2	G	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	INITIATING METHIONINE	UNP P61769
G	0	MET	-	INITIATING METHIONINE	UNP P61769

- Molecule 3 is a protein called Insulin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	10	Total	C	N	O	0	0	0
			69	45	11	13			
3	H	10	Total	C	N	O	0	0	0
			69	45	11	13			

- Molecule 4 is a protein called 1E6 TCR Alpha Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	200	Total	C	N	O	S	0	0	0
			1579	989	260	320	10			

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	I	201	1586	994	261	321	10	0	0	0

- Molecule 5 is a protein called 1E6 TCR Beta Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	246	1974	1249	341	374	10	0	0	0
5	J	246	1973	1249	341	373	10	0	0	0

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
6	A	1	5	4	1	0	0
6	D	1	5	4	1	0	0
6	F	1	5	4	1	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	J	1	Total	C O	0	0
			6	3 3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	18	Total	O	0	0
			18	18		
8	B	10	Total	O	0	0
			10	10		
8	D	7	Total	O	0	0
			7	7		
8	E	14	Total	O	0	0
			14	14		
8	F	15	Total	O	0	0
			15	15		
8	G	5	Total	O	0	0
			5	5		
8	H	3	Total	O	0	0
			3	3		
8	I	4	Total	O	0	0
			4	4		
8	J	14	Total	O	0	0
			14	14		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class I histocompatibility antigen, A-2 alpha chain




- Molecule 1: HLA class I histocompatibility antigen, A-2 alpha chain





- Molecule 2: Beta-2-microglobulin




- Molecule 2: Beta-2-microglobulin




• Molecule 3: Insulin




• Molecule 3: Insulin




• Molecule 4: 1E6 TCR Alpha Chain



• Molecule 4: 1E6 TCR Alpha Chain





• Molecule 5: 1E6 TCR Beta Chain



• Molecule 5: 1E6 TCR Beta Chain





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.68Å 84.59Å 126.28Å 90.00° 90.03° 90.00°	Depositor
Resolution (Å)	63.14 – 2.71 63.14 – 2.71	Depositor EDS
% Data completeness (in resolution range)	100.0 (63.14-2.71) 99.2 (63.14-2.71)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.25 (at 2.73Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.201 , 0.269 0.201 , 0.263	Depositor DCC
$R_{free}$ test set	2715 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	55.5	Xtrriage
Anisotropy	0.549	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 18.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	0.165 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	13552	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	60.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.11% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.65	0/2319	0.59	0/3149
1	F	0.59	0/2330	0.57	0/3163
2	B	0.69	0/860	0.59	0/1162
2	G	0.64	0/860	0.55	0/1162
3	C	0.73	0/72	0.63	0/99
3	H	0.70	0/72	0.58	0/99
4	D	0.72	1/1615 (0.1%)	0.65	0/2185
4	I	0.73	1/1623 (0.1%)	0.66	0/2197
5	E	0.64	0/2029	0.60	0/2759
5	J	0.73	0/2028	0.67	1/2759 (0.0%)
All	All	0.67	2/13808 (0.0%)	0.61	1/18734 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	23	CYS	CB-SG	-8.22	1.68	1.82
4	I	134	CYS	CB-SG	5.55	1.91	1.82

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	J	63	ASP	N-CA-C	-6.38	93.78	111.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2253	0	2103	64	0
1	F	2264	0	2115	80	0
2	B	837	0	803	12	0
2	G	837	0	803	17	0
3	C	69	0	64	2	0
3	H	69	0	64	3	0
4	D	1579	0	1494	68	0
4	I	1586	0	1501	78	0
5	E	1974	0	1887	72	0
5	J	1973	0	1887	64	0
6	A	5	0	0	1	0
6	D	5	0	0	0	0
6	F	5	0	0	1	0
7	J	6	0	8	0	0
8	A	18	0	0	3	0
8	B	10	0	0	0	0
8	D	7	0	0	2	0
8	E	14	0	0	2	0
8	F	15	0	0	0	0
8	G	5	0	0	0	0
8	H	3	0	0	0	0
8	I	4	0	0	0	0
8	J	14	0	0	0	0
All	All	13552	0	12729	429	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 429 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:266:LEU:HD22	1:F:270:LEU:CD1	1.62	1.28
1:F:266:LEU:CD2	1:F:270:LEU:HD12	1.66	1.24
1:F:266:LEU:HD23	1:F:270:LEU:HD12	1.31	1.11
4:D:33:PHE:CD1	4:D:72:ILE:HD11	1.90	1.06

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:266:LEU:CD2	1:F:270:LEU:CD1	2.27	1.05

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/276 (99%)	260 (95%)	14 (5%)	0	100	100
1	F	275/276 (100%)	255 (93%)	18 (6%)	2 (1%)	22	45
2	B	98/100 (98%)	94 (96%)	4 (4%)	0	100	100
2	G	98/100 (98%)	91 (93%)	5 (5%)	2 (2%)	7	17
3	C	8/10 (80%)	7 (88%)	1 (12%)	0	100	100
3	H	8/10 (80%)	7 (88%)	1 (12%)	0	100	100
4	D	198/201 (98%)	185 (93%)	11 (6%)	2 (1%)	15	35
4	I	199/201 (99%)	187 (94%)	10 (5%)	2 (1%)	15	35
5	E	244/246 (99%)	228 (93%)	15 (6%)	1 (0%)	34	58
5	J	244/246 (99%)	236 (97%)	6 (2%)	2 (1%)	19	41
All	All	1646/1666 (99%)	1550 (94%)	85 (5%)	11 (1%)	22	45

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	130	ASP
1	F	227	ASP
5	J	64	ARG
5	E	85	ARG
4	I	151	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/232 (100%)	225 (97%)	7 (3%)	41	69
1	F	233/232 (100%)	226 (97%)	7 (3%)	41	69
2	B	95/95 (100%)	92 (97%)	3 (3%)	39	67
2	G	95/95 (100%)	94 (99%)	1 (1%)	73	89
3	C	5/5 (100%)	5 (100%)	0	100	100
3	H	5/5 (100%)	5 (100%)	0	100	100
4	D	181/182 (100%)	173 (96%)	8 (4%)	28	54
4	I	182/182 (100%)	174 (96%)	8 (4%)	28	54
5	E	216/216 (100%)	206 (95%)	10 (5%)	27	52
5	J	216/216 (100%)	209 (97%)	7 (3%)	39	67
All	All	1460/1460 (100%)	1409 (96%)	51 (4%)	36	63

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	E	80	GLN
1	F	12	VAL
5	J	60	MET
5	E	195	ARG
1	F	35	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 44 such sidechains are listed below:

Mol	Chain	Res	Type
5	E	186	ASN
1	F	180	GLN
5	J	121	ASN
5	E	215	GLN
1	F	70	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	SO4	A	277	-	4,4,4	0.16	0	6,6,6	0.10	0
7	GOL	J	247	-	5,5,5	0.67	0	5,5,5	0.36	0
6	SO4	F	277	-	4,4,4	0.17	0	6,6,6	0.10	0
6	SO4	D	203	-	4,4,4	0.18	0	6,6,6	0.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	J	247	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	J	247	GOL	O1-C1-C2-C3
7	J	247	GOL	C1-C2-C3-O3
7	J	247	GOL	O1-C1-C2-O2
7	J	247	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	277	SO4	1	0
6	F	277	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	276/276 (100%)	-0.17	1 (0%) 92 93	25, 56, 98, 118	0
1	F	276/276 (100%)	0.02	5 (1%) 68 70	28, 69, 141, 173	0
2	B	100/100 (100%)	-0.05	4 (4%) 38 37	30, 55, 98, 119	0
2	G	100/100 (100%)	0.03	4 (4%) 38 37	35, 62, 99, 128	0
3	C	10/10 (100%)	-0.15	0 100 100	31, 32, 38, 40	0
3	H	10/10 (100%)	0.17	1 (10%) 7 5	30, 35, 44, 47	0
4	D	200/201 (99%)	-0.06	3 (1%) 73 76	28, 53, 97, 131	0
4	I	201/201 (100%)	-0.08	0 100 100	26, 49, 88, 111	0
5	E	246/246 (100%)	-0.05	6 (2%) 59 60	26, 55, 103, 139	0
5	J	246/246 (100%)	-0.20	4 (1%) 72 74	25, 45, 89, 108	0
All	All	1665/1666 (99%)	-0.08	28 (1%) 70 72	25, 55, 106, 173	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	G	0	MET	7.8
2	G	1	ILE	6.9
5	E	40	MET	5.1
1	F	220	ASP	4.6
5	E	222	ASN	4.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
7	GOL	J	247	6/6	0.57	0.31	42,69,71,74	0
6	SO4	F	277	5/5	0.83	0.39	100,100,100,100	0
6	SO4	A	277	5/5	0.84	0.41	99,100,100,100	0
6	SO4	D	203	5/5	0.85	0.38	100,100,100,100	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.