wwPDB EM Validation Summary Report

Dec 10, 2022 – 12:18 pm GMT

PDB ID	:	4UUK
EMDB ID	:	EMD-2701
Title	:	Human dynamin 1 K44A superconstricted polymer stabilized with GTP strand 2
Authors	:	Sundborger, A.C.; Fang, S.; Heymann, J.A.; Ray, P.; Chappie, J.S.; Hinshaw, J.E.
Deposited on	:	2014-07-29
Resolution	:	12.50 Å(reported)
Based on initial models	:	3SNH, 1DYN, 3ZYC

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev43
MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.9
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.31.3

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 12.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	864	24% 12% • 62%
1	D	864	24% 12% • 61%
1	G	864	24% 12% • 62%
1	K	864	24% 12% • 61%
2	B	864	10% 7% • • 76%
2	C	864	7% 5% •• 87%
2	E	864	8% 11% • • 76%
2	F	864	7% • •• 87%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	864	
2	I	864	
2	J	864	
2	L	864	

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 21116 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DYNAMIN-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	329	Total 2567	C 1615	N 453	O 489	S 10	0	0
1	D	337	Total 2643	C 1664	N 466	O 503	S 10	0	0
1	G	329	Total 2567	C 1615	N 453	O 489	S 10	0	0
1	K	337	Total 2643	C 1664	N 466	O 503	S 10	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	744	ASN	ASP	variant	UNP Q05193
D	744	ASN	ASP	variant	UNP Q05193
G	744	ASN	ASP	variant	UNP Q05193
K	744	ASN	ASP	variant	UNP Q05193

- Molecule 2 is a protein called DYNAMIN-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	208	Total 1728	C 1097	N 304	O 313	S 14	0	0
2	C	113	Total 946	C 609	N 158	O 175	S 4	0	0
2	E	208	Total 1728	C 1097	N 304	O 313	S 14	0	0
2	F	113	Total 946	C 609	N 158	O 175	S 4	0	0
2	H	113	Total 946	C 609	N 158	O 175	S 4	0	0
2	I	208	Total 1728	C 1097	N 304	O 313	S 14	0	0
2	J	208	Total 1728	C 1097	N 304	O 313	S 14	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
2	L	113	Total	C	N	O	S	0	0
			946	609	158	175	4		

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DYNAMIN-1

Chain A:



• Molecule 1: DYNAMIN-1

Chain D: 



• Molecule 1: DYNAMIN-1

Chain G: 





• Molecule 1: DYNAMIN-1

Chain K:





• Molecule 2: DYNAMIN-1

Chain B:





• Molecule 2: DYNAMIN-1

Chain C: 7% 5% 87%



• Molecule 2: DYNAMIN-1

Chain E: 8% 11% 76%



• Molecule 2: DYNAMIN-1

Chain F: 7% 87%



• Molecule 2: DYNAMIN-1

Chain H: 7% 5% •• 87%





• Molecule 2: DYNAMIN-1

Chain I:





• Molecule 2: DYNAMIN-1

Chain J: 8% 11% 76%





- Molecule 2: DYNAMIN-1

Chain L: 7% 87%



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=Not provided°, rise=Not provided Å, axial sym=Not provided	Depositor
Number of segments used	7525	Depositor
Resolution determination method	Not provided	
CTF correction method	INDIVIDUAL IMAGES	Depositor
Microscope	FEI/PHILIPS CM300FEG/HE	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/Å^2$)	10	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	49000	Depositor
Image detector	KODAK SO-163 FILM	Depositor
Maximum map value	18.352	Depositor
Minimum map value	-18.065	Depositor
Average map value	0.361	Depositor
Map value standard deviation	3.624	Depositor
Recommended contour level	0.6	Depositor
Map size (Å)	510.0, 510.0, 510.0	wwPDB
Map dimensions	200, 200, 200	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	2.55, 2.55, 2.55	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.72	2/2604 (0.1%)	1.19	19/3524 (0.5%)
1	D	0.88	6/2683 (0.2%)	1.44	37/3630 (1.0%)
1	G	0.72	2/2604 (0.1%)	1.19	19/3524 (0.5%)
1	K	0.88	6/2683 (0.2%)	1.44	37/3630 (1.0%)
2	B	1.32	14/1748 (0.8%)	2.41	83/2331 (3.6%)
2	C	0.86	2/966 (0.2%)	1.42	20/1298 (1.5%)
2	E	1.39	20/1748 (1.1%)	2.56	104/2331 (4.5%)
2	F	1.15	7/966 (0.7%)	1.82	37/1298 (2.9%)
2	H	0.86	2/966 (0.2%)	1.42	20/1298 (1.5%)
2	I	1.32	14/1748 (0.8%)	2.41	83/2331 (3.6%)
2	J	1.39	20/1748 (1.1%)	2.56	104/2331 (4.5%)
2	L	1.15	7/966 (0.7%)	1.82	37/1298 (2.9%)
All	All	1.05	102/21430 (0.5%)	1.83	600/28824 (2.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	11
1	D	0	14
1	G	1	12
1	K	0	14
2	B	5	40
2	C	4	9
2	E	10	36
2	F	7	10
2	H	4	9
2	I	5	39
2	J	10	36
2	L	7	10
All	All	54	240

The worst 5 of 102 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	490	HIS	C-O	-16.99	0.91	1.23
2	I	490	HIS	C-O	-16.99	0.91	1.23
2	B	699	SER	CB-OG	-16.13	1.21	1.42
2	I	699	SER	CB-OG	-16.07	1.21	1.42
2	I	492	ASP	C-O	-14.57	0.95	1.23

The worst 5 of 600 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	I	492	ASP	CB-CG-OD2	34.21	149.09	118.30
2	B	492	ASP	CB-CG-OD2	34.19	149.07	118.30
2	J	492	ASP	CB-CG-OD2	32.74	147.76	118.30
2	E	492	ASP	CB-CG-OD2	32.72	147.75	118.30
2	B	489	ASN	O-C-N	-30.67	73.63	122.70

5 of 54 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	726	ASP	CA
2	B	339	ASP	CA
2	B	442	CYS	CA
2	B	489	ASN	CA
2	B	490	HIS	CA

5 of 240 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	105	THR	Mainchain
1	A	123	ARG	Sidechain
1	A	28	ASP	Mainchain
1	A	41	SER	Mainchain
1	A	50	ASN	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2567	0	2629	78	0
1	D	2643	0	2693	164	0
1	G	2567	0	2629	75	0
1	K	2643	0	2693	167	0
2	B	1728	0	1777	198	0
2	C	946	0	937	29	0
2	E	1728	0	1777	285	0
2	F	946	0	935	28	0
2	H	946	0	937	29	0
2	I	1728	0	1777	98	0
2	J	1728	0	1777	193	0
2	L	946	0	935	26	0
All	All	21116	0	21496	1076	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 1076 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:J:491:GLU:CB	2:J:491:GLU:CA	1.74	1.58
2:E:491:GLU:CB	2:E:491:GLU:CA	1.74	1.57
2:B:338:VAL:HG13	2:E:687:HIS:CE1	1.40	1.54
2:J:453:ARG:HG2	1:K:318:ARG:N	1.17	1.44
1:D:318:ARG:N	2:E:453:ARG:HG2	1.17	1.44

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	325/864 (38%)	313 (96%)	10 (3%)	2 (1%)	25 66

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	333/864 (38%)	317 (95%)	14 (4%)	2 (1%)	25	66
1	G	325/864 (38%)	313 (96%)	10 (3%)	2 (1%)	25	66
1	K	333/864 (38%)	317 (95%)	14 (4%)	2 (1%)	25	66
2	B	196/864 (23%)	165 (84%)	22 (11%)	9 (5%)	2	21
2	C	111/864 (13%)	93 (84%)	13 (12%)	5 (4%)	2	22
2	E	196/864 (23%)	161 (82%)	27 (14%)	8 (4%)	3	23
2	F	111/864 (13%)	93 (84%)	12 (11%)	6 (5%)	2	19
2	H	111/864 (13%)	93 (84%)	13 (12%)	5 (4%)	2	22
2	I	196/864 (23%)	165 (84%)	22 (11%)	9 (5%)	2	21
2	J	196/864 (23%)	161 (82%)	27 (14%)	8 (4%)	3	23
2	L	111/864 (13%)	93 (84%)	12 (11%)	6 (5%)	2	19
All	All	2544/10368 (24%)	2284 (90%)	196 (8%)	64 (2%)	9	32

5 of 64 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	88	GLY
2	B	489	ASN
2	B	490	HIS
2	B	492	ASP
2	C	534	MET

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/761 (38%)	275 (96%)	12 (4%)	30	54
1	D	295/761 (39%)	281 (95%)	14 (5%)	26	51
1	G	287/761 (38%)	275 (96%)	12 (4%)	30	54
1	K	295/761 (39%)	281 (95%)	14 (5%)	26	51
2	B	194/761 (26%)	155 (80%)	39 (20%)	1	7

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	C	102/761 (13%)	94 (92%)	8 (8%)	12 36
2	E	194/761 (26%)	153 (79%)	41 (21%)	1 6
2	F	102/761 (13%)	89 (87%)	13 (13%)	4 18
2	H	102/761 (13%)	94 (92%)	8 (8%)	12 36
2	I	194/761 (26%)	155 (80%)	39 (20%)	1 7
2	J	194/761 (26%)	152 (78%)	42 (22%)	1 6
2	L	102/761 (13%)	89 (87%)	13 (13%)	4 18
All	All	2348/9132 (26%)	2093 (89%)	255 (11%)	10 23

5 of 255 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	534	MET
2	J	679	ASP
2	H	579	PHE
2	J	676	THR
1	K	171	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 60 such sidechains are listed below:

Mol	Chain	Res	Type
2	E	704	ASN
1	K	236	ASN
2	H	529	ASN
1	K	155	GLN
2	L	602	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	D	1
1	K	1
2	B	1
2	I	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	D	316:ASN	C	317:PHE	N	1.18
1	K	316:ASN	C	317:PHE	N	1.18
1	B	489:ASN	C	490:HIS	N	1.17
1	I	489:ASN	C	490:HIS	N	1.17

6 Map visualisation

This section contains visualisations of the EMDB entry EMD-2701. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections

6.1.1 Primary map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices

6.2.1 Primary map



X Index: 100



Y Index: 100

Z Index: 100

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 145



Y Index: 55



Z Index: 197

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views (i)

6.4.1 Primary map

X



Y

Z

The images above show the 3D surface view of the map at the recommended contour level 0.6. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate (i)



The volume at the recommended contour level is 33840 nm^3; this corresponds to an approximate mass of 30569 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.080 Å^{-1}

8 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-2701 and PDB model 4UUK. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.6 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model (i)

X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.6).

9.4 Atom inclusion (i)



At the recommended contour level, 100% of all backbone atoms, 99% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.6) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.9909	0.0570
A	0.9877	0.0550
B	0.9893	0.0420
C	0.9805	0.0730
D	0.9958	0.0570
E	0.9941	0.0620
F	0.9913	0.0590
G	0.9873	0.0550
H	0.9816	0.0730
I	0.9905	0.0410
J	0.9941	0.0640
K	0.9958	0.0570
L	0.9924	0.0570