



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 10, 2022 – 04:41 PM EST

PDB ID : 3UXH
Title : Design, Synthesis and Biological Evaluation of Potent Quinoline and Pyrrolo-quinoline Ammosamide Analogues as Inhibitors of Quinone Reductase 2
Authors : Cushman, M.; Mesecar, A.D.; Fanwick, P.E.; Narasimha, R.; Jensen, K.C.
Deposited on : 2011-12-05
Resolution : 1.53 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.31.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2556 (1.56-1.52)
Clashscore	141614	2634 (1.56-1.52)
Ramachandran outliers	138981	2580 (1.56-1.52)
Sidechain outliers	138945	2577 (1.56-1.52)
RSRZ outliers	127900	2524 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	230	 9% 87% 11%
1	B	230	 5% 86% 13%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4190 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribosyldihyronicotinamide dehydrogenase [quinone].

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	230	1829	1178	304	338	9	0	1	0
1	B	230	1834	1181	304	340	9	0	2	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	46	PHE	LEU	variant	UNP P16083
B	46	PHE	LEU	variant	UNP P16083

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
2	A	1	1	1	0	0
2	B	1	1	1	0	0

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
3	A	1	53	27	9	15	2	0	0
3	B	1	53	27	9	15	2	0	0

- Molecule 4 is 6,8-diamino-7-chloro-1-methyl-2-oxo-1,2-dihydropyrrolo[4,3,2-de]quinoline-4-carboxamide (three-letter code: UXH) (formula: C₁₂H₁₀ClN₅O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Cl	N			O
4	A	1	20	12	1	5	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Cl	N			O
4	B	1	20	12	1	5	2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	170	Total	O	0	0
			170	170		
5	B	209	Total	O	0	0
			209	209		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ribosyldihyronicotinamide dehydrogenase [quinone]



- Molecule 1: Ribosyldihyronicotinamide dehydrogenase [quinone]



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.45Å 84.06Å 106.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.11 – 1.53 39.11 – 1.53	Depositor EDS
% Data completeness (in resolution range)	97.0 (39.11-1.53) 97.0 (39.11-1.53)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.58 (at 1.53Å)	Xtrriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.162 , 0.184 0.172 , 0.192	Depositor DCC
R_{free} test set	3780 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	17.2	Xtrriage
Anisotropy	0.087	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 38.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4190	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: UXH, ZN, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.34	4/1882 (0.2%)	1.15	10/2552 (0.4%)
1	B	1.25	2/1890 (0.1%)	1.26	12/2563 (0.5%)
All	All	1.30	6/3772 (0.2%)	1.21	22/5115 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	193	GLU	CD-OE1	6.59	1.32	1.25
1	A	33	GLY	N-CA	5.62	1.54	1.46
1	B	182	LYS	CE-NZ	5.40	1.62	1.49
1	B	193	GLU	CD-OE1	5.29	1.31	1.25
1	A	13	GLU	CD-OE2	5.12	1.31	1.25
1	A	153	GLU	CG-CD	5.04	1.59	1.51

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	78	ARG	NE-CZ-NH1	15.18	127.89	120.30
1	B	78	ARG	NE-CZ-NH2	-12.47	114.06	120.30
1	B	49	ARG	NE-CZ-NH1	9.33	124.96	120.30
1	B	118	ARG	NE-CZ-NH2	-8.98	115.81	120.30
1	B	127[A]	ASP	CB-CG-OD1	8.15	125.64	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	127[B]	ASP	CB-CG-OD1	8.15	125.64	118.30
1	A	104	TYR	CB-CG-CD2	-7.54	116.48	121.00
1	B	52	ASP	CB-CG-OD1	7.34	124.91	118.30
1	B	127[A]	ASP	CB-CG-OD2	-7.33	111.71	118.30
1	B	127[B]	ASP	CB-CG-OD2	-7.33	111.71	118.30
1	A	143	LEU	CB-CG-CD1	-7.07	98.98	111.00
1	A	31	ARG	NE-CZ-NH1	6.80	123.70	120.30
1	A	27	ASP	CB-CG-OD2	-6.44	112.50	118.30
1	A	31	ARG	NE-CZ-NH2	-6.22	117.19	120.30
1	A	65	PHE	CB-CG-CD2	6.19	125.14	120.80
1	B	49	ARG	CG-CD-NE	6.17	124.75	111.80
1	A	184	LEU	CB-CG-CD1	-6.04	100.73	111.00
1	A	101	PHE	CB-CG-CD1	-5.98	116.62	120.80
1	B	86	ASP	CB-CG-OD1	5.20	122.98	118.30
1	A	52	ASP	CB-CG-OD1	5.14	122.93	118.30
1	B	117	ASP	CB-CG-OD1	5.09	122.88	118.30
1	A	1	ALA	C-N-CA	5.06	132.92	122.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1829	0	1788	23	0
1	B	1834	0	1792	18	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	53	0	31	6	0
3	B	53	0	31	2	0
4	A	20	0	10	1	0
4	B	20	0	10	0	0
5	A	170	0	0	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	209	0	0	2	0
All	All	4190	0	3662	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:ALA:CB	1:A:34:CYS:SG	2.61	0.89
1:A:161:ASN:HD21	4:A:303:UXH:HN1C	1.20	0.86
1:A:200:ARG:HD3	3:A:302:FAD:O4B	1.81	0.81
1:A:1:ALA:HB3	1:A:34:CYS:SG	2.26	0.75
3:A:302:FAD:H2B	5:A:561:HOH:O	1.88	0.73
1:B:24:VAL:HG13	5:B:493:HOH:O	1.90	0.71
1:A:197:GLU:O	1:A:201:LYS:HG2	1.92	0.70
1:A:193:GLU:CD	5:A:566:HOH:O	2.31	0.69
3:A:302:FAD:O1A	5:A:566:HOH:O	2.09	0.69
1:B:24:VAL:HG11	1:B:204:VAL:CG1	2.23	0.69
1:A:1:ALA:HB2	1:A:34:CYS:SG	2.37	0.65
1:B:187:GLN:HE21	1:B:210:ARG:HH11	1.48	0.62
1:A:187:GLN:HE21	1:A:210:ARG:HH11	1.47	0.61
1:A:197:GLU:OE2	1:A:200:ARG:NH1	2.33	0.61
1:B:52:ASP:OD1	1:B:53:LYS:HE2	2.01	0.60
1:A:200:ARG:HB3	3:A:302:FAD:C2A	2.32	0.60
1:B:200:ARG:CD	3:B:303:FAD:O4B	2.51	0.59
1:A:196:SER:O	1:A:198:GLU:O	2.25	0.54
1:B:200:ARG:HD2	3:B:303:FAD:O4B	2.09	0.53
1:A:193:GLU:CG	5:A:566:HOH:O	2.57	0.53
1:A:230:GLN:O	1:A:230:GLN:HG3	2.10	0.52
1:A:15:LYS:HD2	1:B:64:VAL:HG22	1.93	0.51
1:A:1:ALA:HA	1:A:3:LYS:HG2	1.92	0.50
1:B:45:ASN:OD1	5:B:596:HOH:O	2.19	0.50
1:B:101:PHE:CZ	1:B:146:VAL:HG22	2.47	0.50
1:B:229:GLY:O	1:B:230:GLN:HG3	2.13	0.48
1:A:200:ARG:HD3	3:A:302:FAD:C4B	2.44	0.47
1:A:172:GLN:HE22	1:A:186:PRO:HD3	1.79	0.46
1:B:4:LYS:HG2	1:B:35:THR:HB	1.97	0.46
1:A:187:GLN:NE2	1:A:210:ARG:HH11	2.14	0.46
1:B:157:LYS:HB3	1:B:157:LYS:HE3	1.60	0.46
1:B:24:VAL:HG11	1:B:204:VAL:HG12	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:197:GLU:C	1:A:198:GLU:O	2.55	0.44
1:A:153:GLU:CG	5:A:505:HOH:O	2.64	0.44
1:A:67:TYR:CZ	1:A:71:THR:HG21	2.53	0.43
1:B:1:ALA:HA	1:B:3:LYS:HG2	2.01	0.42
1:A:153:GLU:HG2	5:A:505:HOH:O	2.20	0.42
1:A:200:ARG:CB	3:A:302:FAD:C2A	2.98	0.42
1:B:187:GLN:NE2	1:B:210:ARG:HH11	2.14	0.41
1:B:165:ARG:HA	1:B:168:LEU:HD12	2.03	0.41
1:B:195:ALA:HB1	1:B:199:GLU:HB2	2.02	0.40
1:B:172:GLN:HE22	1:B:186:PRO:HD3	1.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	229/230 (100%)	218 (95%)	10 (4%)	1 (0%)	34 13
1	B	230/230 (100%)	222 (96%)	8 (4%)	0	100 100
All	All	459/460 (100%)	440 (96%)	18 (4%)	1 (0%)	47 24

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	198	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/194 (100%)	191 (98%)	4 (2%)	53	22
1	B	196/194 (101%)	192 (98%)	4 (2%)	55	24
All	All	391/388 (101%)	383 (98%)	8 (2%)	53	24

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	LYS
1	A	157	LYS
1	A	201	LYS
1	A	230	GLN
1	B	53	LYS
1	B	157	LYS
1	B	197	GLU
1	B	230	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	161	ASN
1	A	172	GLN
1	A	187	GLN
1	A	212	GLN
1	B	172	GLN
1	B	187	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	UXH	A	303	-	19,22,22	3.42	9 (47%)	22,35,35	3.04	11 (50%)
3	FAD	A	302	-	53,58,58	1.85	12 (22%)	68,89,89	2.00	17 (25%)
3	FAD	B	303	-	53,58,58	1.79	14 (26%)	68,89,89	1.96	19 (27%)
4	UXH	B	302	-	19,22,22	3.41	11 (57%)	22,35,35	4.30	11 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	UXH	A	303	-	-	4/4/4/4	0/3/3/3
3	FAD	A	302	-	-	2/30/50/50	0/6/6/6
3	FAD	B	303	-	-	5/30/50/50	0/6/6/6
4	UXH	B	302	-	-	0/4/4/4	0/3/3/3

All (46) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	302	UXH	C10-C9	8.58	1.49	1.39
4	A	303	UXH	C9-C8	7.54	1.48	1.41
4	A	303	UXH	C10-C9	6.93	1.47	1.39
3	A	302	FAD	C10-N1	6.04	1.45	1.33
4	B	302	UXH	C9-C8	5.71	1.46	1.41
3	B	303	FAD	C9A-C5X	5.24	1.50	1.41
4	A	303	UXH	O19-C17	5.20	1.32	1.22
4	A	303	UXH	C7-C8	5.17	1.46	1.40
4	B	302	UXH	C7-C8	5.15	1.46	1.40
4	B	302	UXH	O19-C17	5.09	1.32	1.22
3	A	302	FAD	C1'-C2'	4.97	1.59	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	303	FAD	C4X-N5	4.91	1.40	1.30
3	A	302	FAD	C4X-N5	4.57	1.39	1.30
4	A	303	UXH	C6-C14	-4.09	1.45	1.50
4	A	303	UXH	C4-C3	-3.90	1.36	1.42
3	A	302	FAD	C5X-N5	-3.63	1.32	1.39
4	B	302	UXH	C4-C3	-3.56	1.37	1.42
3	B	303	FAD	C9A-N10	3.22	1.46	1.41
3	B	303	FAD	C1'-C2'	3.09	1.57	1.52
4	B	302	UXH	C10-N18	-3.09	1.36	1.40
3	B	303	FAD	C10-N1	3.07	1.39	1.33
4	A	303	UXH	C2-N1	-3.04	1.31	1.37
3	B	303	FAD	O2-C2	2.94	1.29	1.24
3	B	303	FAD	C5X-N5	-2.70	1.34	1.39
4	A	303	UXH	C10-N18	-2.69	1.36	1.40
3	A	302	FAD	C2-N1	-2.66	1.30	1.36
3	A	302	FAD	C6-C5X	2.61	1.44	1.40
4	B	302	UXH	C6-C14	-2.61	1.47	1.50
3	A	302	FAD	C5A-C4A	2.56	1.47	1.40
3	A	302	FAD	O4B-C1B	2.54	1.44	1.41
3	B	303	FAD	C5A-C4A	2.50	1.47	1.40
3	B	303	FAD	O3'-C3'	2.49	1.48	1.43
4	B	302	UXH	C8-CL1	2.45	1.78	1.72
3	B	303	FAD	C4-N3	2.42	1.43	1.38
4	B	302	UXH	C4-C17	-2.40	1.45	1.49
3	B	303	FAD	C8-C7	2.37	1.46	1.40
3	A	302	FAD	O2-C2	2.33	1.28	1.24
4	B	302	UXH	C2-N1	-2.30	1.32	1.37
4	A	303	UXH	C4-C17	-2.29	1.46	1.49
3	A	302	FAD	O4'-C4'	-2.28	1.38	1.43
3	A	302	FAD	C10-N10	-2.20	1.32	1.37
3	B	303	FAD	C8M-C8	-2.18	1.46	1.51
4	B	302	UXH	O16-C14	2.12	1.28	1.24
3	B	303	FAD	C5'-C4'	2.09	1.54	1.51
3	A	302	FAD	C6A-C5A	2.05	1.50	1.43
3	B	303	FAD	P-O5'	-2.00	1.51	1.59

All (58) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	302	UXH	C10-N18-C17	-11.97	103.96	110.97
4	A	303	UXH	C10-N18-C17	-8.85	105.79	110.97
4	B	302	UXH	O19-C17-C4	-7.67	122.38	129.78

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	302	FAD	N3-C2-N1	6.81	132.74	119.38
4	B	302	UXH	C3-C10-N18	6.51	112.32	106.63
4	B	302	UXH	C6-C14-N15	5.61	121.80	116.25
3	A	302	FAD	O2-C2-N1	-5.46	112.78	121.83
3	B	303	FAD	C5X-C9A-N10	-5.38	112.40	117.95
4	B	302	UXH	C5-C4-C17	5.37	138.90	129.41
4	B	302	UXH	C9-C8-C7	5.19	131.14	122.58
3	B	303	FAD	C4-N3-C2	-5.07	116.28	125.64
3	B	303	FAD	N3-C2-N1	5.04	129.28	119.38
3	A	302	FAD	C4-N3-C2	-4.96	116.47	125.64
4	B	302	UXH	C8-C7-N11	4.82	128.38	121.92
4	A	303	UXH	O16-C14-C6	4.72	123.56	119.61
4	B	302	UXH	C7-C8-CL1	-4.29	112.80	118.82
3	A	302	FAD	C1B-N9A-C4A	-4.27	119.14	126.64
3	B	303	FAD	C4X-C10-N10	4.04	122.39	116.48
4	A	303	UXH	C5-C4-C17	4.01	136.49	129.41
4	A	303	UXH	C9-C8-C7	3.90	129.02	122.58
4	A	303	UXH	O19-C17-C4	-3.85	126.07	129.78
3	A	302	FAD	C4A-C5A-N7A	-3.74	105.50	109.40
3	A	302	FAD	N3A-C2A-N1A	-3.65	122.97	128.68
3	A	302	FAD	C4X-C10-N1	-3.53	116.53	124.73
3	B	303	FAD	O2-C2-N3	-3.39	112.05	118.65
4	A	303	UXH	C3-C10-N18	3.35	109.56	106.63
3	A	302	FAD	C9A-C5X-N5	3.26	125.97	122.43
3	B	303	FAD	C4-C4X-N5	3.26	122.87	118.23
3	B	303	FAD	C4A-C5A-N7A	-3.26	106.00	109.40
3	B	303	FAD	C9-C9A-N10	3.24	126.22	121.84
4	A	303	UXH	C4-C5-C6	-3.21	115.25	120.33
3	A	302	FAD	O2'-C2'-C1'	3.18	117.48	109.80
3	B	303	FAD	C10-C4X-N5	-3.15	118.16	124.86
3	B	303	FAD	N3A-C2A-N1A	-3.05	123.91	128.68
4	B	302	UXH	C5-C6-C14	2.84	127.47	118.59
3	A	302	FAD	C4-C4X-C10	2.75	121.42	116.79
3	B	303	FAD	C1'-N10-C9A	-2.69	116.02	120.51
3	B	303	FAD	C1B-N9A-C4A	-2.60	122.08	126.64
4	A	303	UXH	C5-C6-C14	2.59	126.69	118.59
4	A	303	UXH	C7-C8-CL1	-2.57	115.22	118.82
4	B	302	UXH	O16-C14-C6	-2.52	117.50	119.61
3	B	303	FAD	C5'-C4'-C3'	-2.51	107.35	112.20
3	A	302	FAD	O4'-C4'-C3'	-2.45	103.15	109.10
3	A	302	FAD	C5X-C9A-N10	-2.42	115.46	117.95
3	A	302	FAD	C1'-N10-C9A	-2.42	116.48	120.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	303	FAD	O2A-PA-O5B	-2.40	96.60	107.75
3	B	303	FAD	O4'-C4'-C5'	-2.38	104.56	109.92
3	A	302	FAD	C5A-C6A-N6A	2.31	123.86	120.35
3	B	303	FAD	C9A-C5X-N5	2.29	124.92	122.43
4	A	303	UXH	C9-C8-CL1	-2.22	115.76	118.83
3	A	302	FAD	O2-C2-N3	-2.18	114.40	118.65
3	A	302	FAD	C4X-C10-N10	2.17	119.65	116.48
3	A	302	FAD	C2A-N1A-C6A	2.14	122.41	118.75
4	B	302	UXH	C9-C8-CL1	-2.13	115.89	118.83
4	A	303	UXH	C3-C4-C17	-2.11	102.65	105.80
3	B	303	FAD	C2A-N1A-C6A	2.05	122.26	118.75
3	B	303	FAD	C4X-C10-N1	-2.03	120.02	124.73
3	B	303	FAD	C4X-C4-N3	2.01	118.30	113.19

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	303	FAD	C5B-O5B-PA-O1A
3	B	303	FAD	C5B-O5B-PA-O2A
4	A	303	UXH	N15-C14-C6-C5
4	A	303	UXH	O16-C14-C6-C5
4	A	303	UXH	O16-C14-C6-N1
3	B	303	FAD	C5B-O5B-PA-O3P
3	B	303	FAD	C4'-C5'-O5'-P
4	A	303	UXH	N15-C14-C6-N1
3	A	302	FAD	C4'-C5'-O5'-P
3	A	302	FAD	P-O3P-PA-O1A
3	B	303	FAD	C4B-C5B-O5B-PA


There are no ring outliers.


3 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	303	UXH	1	0
3	A	302	FAD	6	0
3	B	303	FAD	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	230/230 (100%)	0.44	21 (9%) 9 10	10, 22, 46, 72	0
1	B	230/230 (100%)	0.16	11 (4%) 30 34	10, 18, 38, 59	0
All	All	460/460 (100%)	0.30	32 (6%) 16 18	10, 20, 42, 72	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	230	GLN	9.1
1	A	1	ALA	6.5
1	B	1	ALA	6.4
1	B	2	GLY	5.4
1	B	230	GLN	4.7
1	A	2	GLY	4.0
1	B	197	GLU	3.6
1	A	58	THR	3.6
1	A	229	GLY	3.6
1	A	197	GLU	3.5
1	A	57	GLY	3.4
1	A	63	GLU	3.3
1	B	198	GLU	3.2
1	B	112	LEU	3.1
1	A	158	THR	3.1
1	B	201	LYS	3.0
1	A	112	LEU	3.0
1	A	108	VAL	2.9
1	A	153	GLU	2.9
1	A	201	LYS	2.9
1	A	59	LEU	2.4
1	B	196	SER	2.4
1	B	63	GLU	2.3
1	A	78	ARG	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	108	VAL	2.2
1	A	101	PHE	2.1
1	A	103	LEU	2.1
1	A	171	LEU	2.1
1	B	111	ILE	2.1
1	A	198	GLU	2.1
1	A	62	PRO	2.0
1	A	196	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	FAD	A	302	53/53	0.91	0.14	11,18,62,64	0
3	FAD	B	303	53/53	0.91	0.13	12,17,47,49	0
4	UXH	B	302	20/20	0.95	0.08	15,19,26,26	0
4	UXH	A	303	20/20	0.96	0.07	15,18,23,25	0
2	ZN	A	301	1/1	0.99	0.03	26,26,26,26	0
2	ZN	B	301	1/1	1.00	0.03	21,21,21,21	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD B 303:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around UXH B 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.