



# wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 8, 2023 – 05:54 am GMT

PDB ID : 2V5H  
Title : Controlling the storage of nitrogen as arginine: the complex of PII and acetylglutamate kinase from *Synechococcus elongatus* PCC 7942  
Authors : Llacer, J.L.; Marco-Marin, C.; Gil-Ortiz, F.; Fita, I.; Rubio, V.  
Deposited on : 2007-07-04  
Resolution : 2.75 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	321	 2% 71% 13% • 12%
1	B	321	 % 73% 14% • 10%
1	C	321	 72% 13% • 12%
1	D	321	 2% 73% 12% • • 11%
1	E	321	 2% 72% 13% • 11%
1	F	321	 2% 71% 13% • 11%

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	112	 7% 77% 20% ..
2	H	112	 3% 73% 22% ..
2	I	112	 11% 74% 21% ..
2	J	112	 % 77% 21% ..
2	K	112	 78% 17% ..
2	L	112	 2% 81% 15% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	CL	K	1109	-	-	X	-

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 18185 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ACETYLGLUTAMATE KINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	284	Total 2126	C 1333	N 386	O 396	S 11	0	1	0
1	B	289	Total 2165	C 1357	N 390	O 407	S 11	0	2	0
1	C	284	Total 2128	C 1334	N 383	O 400	S 11	0	1	0
1	D	286	Total 2137	C 1339	N 385	O 402	S 11	0	1	0
1	E	286	Total 2142	C 1341	N 385	O 405	S 11	0	2	0
1	F	287	Total 2142	C 1342	N 386	O 403	S 11	0	1	0

- Molecule 2 is a protein called NITROGEN REGULATORY PROTEIN P-II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	G	111	Total 853	C 540	N 149	O 162	S 2	0	0	0
2	H	108	Total 840	C 531	N 148	O 159	S 2	0	0	0
2	I	109	Total 838	C 531	N 147	O 158	S 2	0	0	0
2	J	110	Total 850	C 537	N 150	O 161	S 2	0	0	0
2	K	108	Total 840	C 531	N 148	O 159	S 2	0	0	0
2	L	109	Total 841	C 532	N 148	O 159	S 2	0	0	0

- Molecule 3 is N-ACETYL-L-GLUTAMATE (three-letter code: NLG) (formula: C<sub>7</sub>H<sub>11</sub>NO<sub>5</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	13	7	1	5	0	0
3	B	1	13	7	1	5	0	0
3	C	1	13	7	1	5	0	0
3	D	1	13	7	1	5	0	0
3	E	1	13	7	1	5	0	0
3	F	1	13	7	1	5	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Na		
4	A	1	1	1	0	0
4	B	1	1	1	0	0
4	C	1	1	1	0	0
4	D	1	1	1	0	0
4	E	1	1	1	0	0
4	F	1	1	1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	1	Total	C O	0	0
			6	3 3		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	G	1	Total	Cl	0	0
			1	1		
6	H	2	Total	Cl	0	0
			2	2		
6	J	2	Total	Cl	0	0
			2	2		
6	K	1	Total	Cl	0	0
			1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	29	Total	O	0	0
			29	29		
7	B	22	Total	O	0	0
			22	22		
7	C	16	Total	O	0	0
			16	16		
7	D	16	Total	O	0	0
			16	16		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
7	E	22	Total 22	O 22	0	0
7	F	23	Total 23	O 23	0	0
7	G	9	Total 9	O 9	0	0
7	H	11	Total 11	O 11	0	0
7	I	8	Total 8	O 8	0	0
7	J	10	Total 10	O 10	0	0
7	K	7	Total 7	O 7	0	0
7	L	14	Total 14	O 14	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: ACETYLGLUTAMATE KINASE



#### • Molecule 1: ACETYLGLUTAMATE KINASE



#### • Molecule 1: ACETYLGLUTAMATE KINASE






HIS  
GLU  
ALA  
HIS  
GLN  
PRO  
TRP  
GLN


● Molecule 1: ACETYLGLUTAMATE KINASE

Chain D: 73% 12% 11%




● Molecule 1: ACETYLGLUTAMATE KINASE

Chain E: 72% 13% 11%



● Molecule 1: ACETYLGLUTAMATE KINASE

Chain F: 71% 13% 11%




● Molecule 2: NITROGEN REGULATORY PROTEIN P-II



Chain G: 77% 20% 3%





- Molecule 2: NITROGEN REGULATORY PROTEIN P-II




- Molecule 2: NITROGEN REGULATORY PROTEIN P-II




- Molecule 2: NITROGEN REGULATORY PROTEIN P-II



- Molecule 2: NITROGEN REGULATORY PROTEIN P-II



- Molecule 2: NITROGEN REGULATORY PROTEIN P-II



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.38Å 161.03Å 91.56Å 90.00° 106.53° 90.00°	Depositor
Resolution (Å)	50.00 – 2.75 45.79 – 2.75	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.00-2.75) 100.0 (45.79-2.75)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.16 (at 2.77Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.200 , 0.236 0.203 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.7	Xtrriage
Anisotropy	0.089	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 50.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.022 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	18185	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, NLG, GOL, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.60	0/2155	0.71	2/2920 (0.1%)
1	B	0.56	0/2196	0.70	2/2979 (0.1%)
1	C	0.52	0/2157	0.69	2/2923 (0.1%)
1	D	0.53	1/2166 (0.0%)	0.65	0/2935
1	E	0.55	0/2171	0.65	0/2943
1	F	0.50	1/2171 (0.0%)	0.61	0/2943
2	G	0.50	0/860	0.66	0/1155
2	H	0.52	0/847	0.71	0/1137
2	I	0.51	0/844	0.67	0/1133
2	J	0.52	0/857	0.70	0/1151
2	K	0.55	0/847	0.72	0/1137
2	L	0.57	0/848	0.71	1/1139 (0.1%)
All	All	0.54	2/18119 (0.0%)	0.68	7/24495 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	2
1	E	0	1
1	F	0	2
All	All	0	5

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	218	LYS	CD-CE	6.18	1.66	1.51
1	F	251	CYS	CB-SG	-5.38	1.73	1.81

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	88	HIS	CB-CA-C	9.76	129.91	110.40
1	C	11	ALA	CB-CA-C	-8.28	97.68	110.10
1	C	10	ALA	N-CA-C	6.89	129.59	111.00
1	B	10	ALA	CB-CA-C	-6.07	100.99	110.10
1	B	10	ALA	N-CA-C	5.95	127.06	111.00

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	12	ASP	Peptide
1	D	9	GLY	Peptide
1	E	9	GLY	Peptide
1	F	11	ALA	Peptide
1	F	9	GLY	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2126	0	2176	54	0
1	B	2165	0	2190	53	0
1	C	2128	0	2173	50	0
1	D	2137	0	2178	58	0
1	E	2142	0	2172	54	0
1	F	2142	0	2181	48	1
2	G	853	0	892	12	0
2	H	840	0	880	16	0
2	I	838	0	874	16	0
2	J	850	0	887	14	0
2	K	840	0	880	15	1
2	L	841	0	879	8	0
3	A	13	0	9	1	0
3	B	13	0	9	2	0
3	C	13	0	9	1	0
3	D	13	0	9	2	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	13	0	9	0	0
3	F	13	0	9	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
5	E	6	0	8	1	0
6	G	1	0	0	0	0
6	H	2	0	0	1	0
6	J	2	0	0	1	0
6	K	1	0	0	4	0
7	A	29	0	0	3	0
7	B	22	0	0	0	0
7	C	16	0	0	2	0
7	D	16	0	0	1	0
7	E	22	0	0	5	0
7	F	23	0	0	1	0
7	G	9	0	0	2	0
7	H	11	0	0	2	0
7	I	8	0	0	0	0
7	J	10	0	0	3	0
7	K	7	0	0	1	0
7	L	14	0	0	1	0
All	All	18185	0	18424	357	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 357 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:J:38:ARG:HD2	2:J:52:THR:O	1.69	0.93
1:C:12:ASP:N	1:C:13:ARG:HB2	1.85	0.91
1:A:211:ARG:HG2	7:A:2020:HOH:O	1.71	0.90
1:D:211:ARG:HH11	1:D:211:ARG:HG3	1.35	0.88
1:B:11:ALA:HA	1:B:13:ARG:HG3	1.65	0.79

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:80:ARG:NH2	2:K:22:ASN:O[1_455]	2.02	0.18

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	283/321 (88%)	274 (97%)	5 (2%)	4 (1%)	11	19
1	B	289/321 (90%)	277 (96%)	8 (3%)	4 (1%)	11	19
1	C	283/321 (88%)	274 (97%)	7 (2%)	2 (1%)	22	39
1	D	285/321 (89%)	280 (98%)	2 (1%)	3 (1%)	14	25
1	E	286/321 (89%)	280 (98%)	4 (1%)	2 (1%)	22	39
1	F	286/321 (89%)	273 (96%)	9 (3%)	4 (1%)	11	19
2	G	109/112 (97%)	103 (94%)	5 (5%)	1 (1%)	17	31
2	H	106/112 (95%)	102 (96%)	4 (4%)	0	100	100
2	I	105/112 (94%)	101 (96%)	4 (4%)	0	100	100
2	J	108/112 (96%)	101 (94%)	6 (6%)	1 (1%)	17	31
2	K	106/112 (95%)	101 (95%)	4 (4%)	1 (1%)	17	31
2	L	107/112 (96%)	104 (97%)	3 (3%)	0	100	100
All	All	2353/2598 (91%)	2270 (96%)	61 (3%)	22 (1%)	17	31

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	178	ASP
1	F	178	ASP
1	A	10	ALA
1	A	70	GLY
1	A	89	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/256 (88%)	203 (91%)	21 (9%)	8	15
1	B	226/256 (88%)	205 (91%)	21 (9%)	9	15
1	C	225/256 (88%)	203 (90%)	22 (10%)	8	13
1	D	225/256 (88%)	201 (89%)	24 (11%)	6	11
1	E	225/256 (88%)	204 (91%)	21 (9%)	9	15
1	F	225/256 (88%)	200 (89%)	25 (11%)	6	10
2	G	90/93 (97%)	81 (90%)	9 (10%)	7	13
2	H	90/93 (97%)	82 (91%)	8 (9%)	9	17
2	I	88/93 (95%)	81 (92%)	7 (8%)	12	21
2	J	90/93 (97%)	81 (90%)	9 (10%)	7	13
2	K	90/93 (97%)	83 (92%)	7 (8%)	12	22
2	L	89/93 (96%)	82 (92%)	7 (8%)	12	22
All	All	1887/2094 (90%)	1706 (90%)	181 (10%)	8	14

5 of 181 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	48	GLU
2	H	13	LEU
1	F	131	THR
1	F	261	ARG
2	H	101	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	156	ASN
1	F	200	ASN
2	J	98	GLN
2	I	108	ASN

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	B	89	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 12 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NLG	D	1294	-	12,12,12	1.89	1 (8%)	15,15,15	1.04	0
3	NLG	F	1293	-	12,12,12	1.90	1 (8%)	15,15,15	1.19	1 (6%)
3	NLG	B	1297	-	12,12,12	1.83	1 (8%)	15,15,15	1.08	1 (6%)
3	NLG	E	1294	-	12,12,12	1.97	2 (16%)	15,15,15	1.22	2 (13%)
5	GOL	E	1295	-	5,5,5	0.43	0	5,5,5	0.29	0
3	NLG	A	1292	-	12,12,12	1.85	1 (8%)	15,15,15	1.23	1 (6%)
3	NLG	C	1292	-	12,12,12	1.82	1 (8%)	15,15,15	1.41	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NLG	D	1294	-	-	0/13/13/13	-
3	NLG	F	1293	-	-	1/13/13/13	-
3	NLG	B	1297	-	-	2/13/13/13	-
3	NLG	E	1294	-	-	1/13/13/13	-
5	GOL	E	1295	-	-	4/4/4/4	-
3	NLG	A	1292	-	-	4/13/13/13	-
3	NLG	C	1292	-	-	4/13/13/13	-

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	1294	NLG	C8-C7	-5.57	1.39	1.50
3	D	1294	NLG	C8-C7	-5.56	1.39	1.50
3	F	1293	NLG	C8-C7	-5.55	1.39	1.50
3	C	1292	NLG	C8-C7	-5.52	1.39	1.50
3	A	1292	NLG	C8-C7	-5.28	1.39	1.50

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1292	NLG	O7-C7-C8	-2.75	116.94	122.06
3	A	1292	NLG	O7-C7-C8	-2.44	117.53	122.06
3	E	1294	NLG	O7-C7-C8	-2.28	117.83	122.06
3	C	1292	NLG	OE1-CD-CG	-2.23	115.93	123.08
3	B	1297	NLG	O7-C7-C8	-2.16	118.04	122.06

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	E	1295	GOL	C1-C2-C3-O3
5	E	1295	GOL	O2-C2-C3-O3
5	E	1295	GOL	O1-C1-C2-C3
5	E	1295	GOL	O1-C1-C2-O2
3	A	1292	NLG	CA-CB-CG-CD

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	1294	NLG	2	0
3	B	1297	NLG	2	0
5	E	1295	GOL	1	0
3	A	1292	NLG	1	0
3	C	1292	NLG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	284/321 (88%)	0.21	5 (1%) 68 76	31, 37, 46, 50	0
1	B	289/321 (90%)	0.13	3 (1%) 82 87	31, 37, 46, 50	1 (0%)
1	C	284/321 (88%)	0.15	0 100 100	31, 37, 45, 50	0
1	D	286/321 (89%)	0.21	5 (1%) 70 78	31, 37, 46, 52	0
1	E	286/321 (89%)	0.21	7 (2%) 59 68	31, 37, 46, 54	1 (0%)
1	F	287/321 (89%)	0.25	6 (2%) 63 72	31, 37, 45, 50	0
2	G	111/112 (99%)	0.48	8 (7%) 15 18	31, 38, 49, 53	0
2	H	108/112 (96%)	0.35	3 (2%) 53 62	31, 37, 46, 56	0
2	I	109/112 (97%)	0.85	12 (11%) 5 6	31, 37, 48, 53	0
2	J	110/112 (98%)	0.30	1 (0%) 84 89	31, 38, 46, 52	0
2	K	108/112 (96%)	0.32	0 100 100	31, 38, 46, 55	0
2	L	109/112 (97%)	0.25	2 (1%) 68 76	31, 38, 45, 48	0
All	All	2371/2598 (91%)	0.26	52 (2%) 62 70	31, 37, 46, 56	2 (0%)

The worst 5 of 52 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	G	41	GLY	6.0
1	E	293	TYR	5.9
2	I	109	ALA	5.4
1	F	6	ILE	4.8
1	E	12	ASP	4.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NA	A	1293	1/1	0.51	0.18	57,57,57,57	0
4	NA	F	1294	1/1	0.71	0.20	47,47,47,47	0
4	NA	E	1296	1/1	0.84	0.14	46,46,46,46	0
5	GOL	E	1295	6/6	0.84	0.29	56,56,57,58	0
4	NA	C	1293	1/1	0.86	0.16	36,36,36,36	0
4	NA	D	1295	1/1	0.87	0.11	50,50,50,50	0
3	NLG	F	1293	13/13	0.90	0.25	42,46,47,49	0
6	CL	H	1110	1/1	0.90	0.23	63,63,63,63	0
6	CL	J	1112	1/1	0.90	0.13	62,62,62,62	0
4	NA	B	1298	1/1	0.91	0.10	37,37,37,37	0
3	NLG	A	1292	13/13	0.91	0.24	50,53,54,54	0
3	NLG	C	1292	13/13	0.92	0.20	38,39,40,41	0
3	NLG	E	1294	13/13	0.93	0.20	41,43,44,46	0
3	NLG	B	1297	13/13	0.94	0.19	42,44,47,48	0
6	CL	G	1112	1/1	0.94	0.23	43,43,43,43	0
6	CL	J	1111	1/1	0.95	0.15	30,30,30,30	0
3	NLG	D	1294	13/13	0.95	0.20	41,45,49,49	0
6	CL	H	1109	1/1	0.97	0.12	35,35,35,35	0
6	CL	K	1109	1/1	0.99	0.12	39,39,39,39	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.