



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 12:58 pm BST

PDB ID : 2V78
Title : Crystal structure of Sulfolobus solfataricus 2-keto-3-deoxygluconate kinase
Authors : Potter, J.A.; Theodossis, A.; Kerou, M.; Lamble, H.J.; Bull, S.D.; Hough, D.W.; Danson, M.J.; Taylor, G.L.
Deposited on : 2007-07-27
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	313	 90% 8% ..
1	B	313	 86% 13% .
1	C	313	 87% 11% ..

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8098 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FRUCTOKINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	311	2571	1646	427	494	4	0	19	0
1	B	311	2559	1632	428	494	5	0	18	0
1	C	311	2506	1598	421	483	4	0	9	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	173	Total 173	O 173	0	0
2	B	150	Total 150	O 150	0	0
2	C	139	Total 139	O 139	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: FRUCTOKINASE

Chain A:  90% 8% ..




- Molecule 1: FRUCTOKINASE

Chain B:  86% 13% .



- Molecule 1: FRUCTOKINASE

Chain C:  87% 11% ..



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	104.86Å 104.86Å 422.50Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.98 – 2.00 46.96 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.9 (46.98-2.00) 96.9 (46.96-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.31 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.162 , 0.198 0.162 , 0.196	Depositor DCC
R_{free} test set	4554 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	22.1	Xtrriage
Anisotropy	0.108	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 53.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8098	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.97	0/2650	0.95	6/3580 (0.2%)
1	B	0.94	0/2632	0.91	6/3558 (0.2%)
1	C	0.94	0/2564	0.90	4/3467 (0.1%)
All	All	0.95	0/7846	0.92	16/10605 (0.2%)

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	159	ARG	NE-CZ-NH2	-8.52	116.04	120.30
1	A	159	ARG	NE-CZ-NH2	-8.29	116.16	120.30
1	A	159	ARG	NE-CZ-NH1	7.35	123.98	120.30
1	A	312	LYS	CD-CE-NZ	-7.13	95.29	111.70
1	B	166	ARG	NE-CZ-NH2	-6.94	116.83	120.30
1	B	70	ARG	NE-CZ-NH1	6.73	123.66	120.30
1	C	15	ASN	CB-CA-C	-6.47	97.45	110.40
1	B	70	ARG	NE-CZ-NH2	-6.31	117.14	120.30
1	A	301	LEU	CB-CG-CD1	6.19	121.52	111.00
1	A	301	LEU	CA-CB-CG	5.92	128.93	115.30
1	B	94	ARG	NE-CZ-NH1	5.88	123.24	120.30
1	C	159	ARG	NE-CZ-NH1	5.62	123.11	120.30
1	C	161	LEU	CA-CB-CG	5.53	128.01	115.30
1	B	94	ARG	NE-CZ-NH2	-5.25	117.67	120.30
1	A	70	ARG	NE-CZ-NH2	-5.21	117.70	120.30
1	B	130	ARG	NE-CZ-NH2	-5.09	117.75	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2571	0	2618	12	1
1	B	2559	0	2585	25	1
1	C	2506	0	2525	18	0
2	A	173	0	0	2	0
2	B	150	0	0	3	0
2	C	139	0	0	2	0
All	All	8098	0	7728	54	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:190[B]:GLU:OE1	1:B:221[B]:LYS:HE2	1.69	0.91
1:B:190[B]:GLU:CD	1:B:221[B]:LYS:HE2	1.92	0.90
1:A:296:GLU:HG3	2:A:2164:HOH:O	1.76	0.84
1:C:130[A]:ARG:HH11	1:C:130[A]:ARG:HG2	1.42	0.83
1:B:151:LYS:O	1:B:154[A]:GLU:HG2	1.83	0.79
1:B:292[A]:ARG:O	2:B:2140:HOH:O	2.08	0.71
1:B:13:GLN:OE1	1:B:15[B]:ASN:ND2	2.15	0.70
1:A:186[B]:LYS:HD2	1:A:187:TYR:CZ	2.31	0.66
1:B:151:LYS:O	1:B:154[A]:GLU:CG	2.45	0.64
1:B:187:TYR:O	2:B:2102:HOH:O	2.14	0.64
1:C:151:LYS:O	1:C:155:LEU:HD13	2.01	0.61
1:A:186[B]:LYS:HD2	1:A:187:TYR:CE1	2.38	0.58
1:B:243:LYS:HD3	1:B:279:ALA:HA	1.88	0.56
1:C:80:LYS:HD2	1:C:80:LYS:N	2.20	0.56
1:B:16[A]:SER:HB2	1:B:25:VAL:HG11	1.88	0.55
1:B:190[B]:GLU:OE1	1:B:221[B]:LYS:CE	2.50	0.55
1:B:123:GLU:O	1:B:127:ARG:HG3	2.10	0.52
1:A:123:GLU:HA	1:A:155:LEU:HD13	1.95	0.48
1:B:184:LEU:O	2:B:2102:HOH:O	2.20	0.48
1:C:159:ARG:HD2	1:C:188:ASP:O	2.13	0.48
1:C:16[A]:SER:HB2	1:C:25:VAL:HG11	1.96	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:29:GLU:CD	1:B:292[B]:ARG:HE	2.16	0.47
1:B:250:VAL:HG22	1:B:289:ILE:HD12	1.96	0.47
1:C:140:ALA:HB2	1:C:170:TRP:CE2	2.51	0.46
1:A:117:SER:OG	1:A:119[A]:GLU:HG2	2.16	0.46
1:C:130[A]:ARG:HH11	1:C:130[A]:ARG:CG	2.19	0.46
1:B:40:CYS:O	1:B:44[B]:VAL:HG12	2.15	0.46
1:A:129:SER:O	1:A:157:LYS:HE3	2.17	0.45
1:C:195:ASP:HB2	1:C:196:PRO:CD	2.47	0.45
1:C:40:CYS:HG	1:C:50:CYS:HG	1.66	0.44
1:B:12:ILE:HB	1:B:89:ILE:HG22	1.99	0.44
1:C:127:ARG:O	1:C:157:LYS:HE3	2.19	0.43
1:C:154[A]:GLU:HG3	1:C:187:TYR:OH	2.18	0.43
1:B:146:LYS:HD2	1:B:179:THR:HG23	2.01	0.43
1:A:186[B]:LYS:HB3	1:A:187:TYR:CD1	2.54	0.42
1:A:80[B]:LYS:HD3	2:A:2079:HOH:O	2.18	0.42
1:B:154[A]:GLU:HG3	1:B:155:LEU:HG	2.00	0.42
1:B:129:SER:O	1:B:157:LYS:HE2	2.19	0.42
1:C:94[B]:ARG:HG3	2:C:2113:HOH:O	2.19	0.42
1:C:285[A]:SER:O	1:C:289:ILE:HG23	2.19	0.42
1:C:92:ILE:HG12	1:C:104:LEU:HD22	2.00	0.42
1:B:140:ALA:HB2	1:B:170:TRP:CE2	2.55	0.42
1:B:154[A]:GLU:HG3	1:B:155:LEU:N	2.35	0.42
1:C:247:LYS:HB3	1:C:247:LYS:HE2	1.77	0.42
1:C:247:LYS:HE3	2:C:2042:HOH:O	2.19	0.41
1:A:137:ILE:O	1:A:141:ILE:HG23	2.20	0.41
1:A:159:ARG:HD2	1:A:188:ASP:O	2.21	0.41
1:B:71:ALA:HB2	1:C:71:ALA:HB2	2.03	0.40
1:A:203:LEU:HD13	1:A:214:LYS:HB2	2.03	0.40
1:B:123:GLU:HA	1:B:155:LEU:HD13	2.03	0.40
1:B:39:PHE:CE1	1:B:262:GLY:HA2	2.56	0.40
1:C:18:ASN:HB3	1:C:19:PRO:HD2	2.01	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:178[B]:GLU:OE1	1:B:186:LYS:NZ[10_555]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	328/313 (105%)	322 (98%)	5 (2%)	1 (0%)	41	37
1	B	327/313 (104%)	317 (97%)	10 (3%)	0	100	100
1	C	318/313 (102%)	310 (98%)	8 (2%)	0	100	100
All	All	973/939 (104%)	949 (98%)	23 (2%)	1 (0%)	51	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	164	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	286/269 (106%)	275 (96%)	11 (4%)	33	31
1	B	284/269 (106%)	276 (97%)	8 (3%)	43	44
1	C	276/269 (103%)	260 (94%)	16 (6%)	20	15
All	All	846/807 (105%)	811 (96%)	35 (4%)	32	28

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18[A]	ASN
1	A	18[B]	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	64	ASN
1	A	161[A]	LEU
1	A	161[B]	LEU
1	A	167	PRO
1	A	173	LEU
1	A	218	LEU
1	A	269	LEU
1	A	301	LEU
1	A	312	LYS
1	B	64	ASN
1	B	161[A]	LEU
1	B	161[B]	LEU
1	B	166	ARG
1	B	173	LEU
1	B	183	ILE
1	B	200	LYS
1	B	218	LEU
1	C	17	PHE
1	C	64	ASN
1	C	80	LYS
1	C	130[A]	ARG
1	C	130[B]	ARG
1	C	144	ASN
1	C	155	LEU
1	C	161	LEU
1	C	173	LEU
1	C	186	LYS
1	C	206	THR
1	C	210	GLU
1	C	218	LEU
1	C	247	LYS
1	C	305	GLU
1	C	312	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	ASN
1	B	13	GLN
1	B	58	ASN
1	B	83	ASN
1	C	144	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	311/313 (99%)	-0.57	1 (0%) 94 93	13, 19, 28, 37	10 (3%)
1	B	311/313 (99%)	-0.63	0 100 100	12, 19, 29, 39	12 (3%)
1	C	311/313 (99%)	-0.43	0 100 100	13, 21, 30, 41	13 (4%)
All	All	933/939 (99%)	-0.54	1 (0%) 95 95	12, 20, 29, 41	35 (3%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	18[A]	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.