

# Full wwPDB X-ray Structure Validation Report (i)

#### Oct 5, 2023 – 05:06 AM EDT

PDB ID : 6V7E

Title : Human Arginase1 Complexed with Bicyclic Inhibitor Compound 12

Authors : Palte, R.L. Deposited on : 2019-12-08

Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : FAILED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\ DIFFRACTION$ 

The reported resolution of this entry is 1.99 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



## 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 15697 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Arginase-1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	317	Total	С	N	О	S	0	0	0
1	A	317	2409	1535	411	457	6	0	U	U
1	В	317	Total	С	N	О	S	0	0	0
1	Ъ	317	2409	1535	411	457	6	0	0	0
1	1 C	317	Total	С	N	О	S	0	0	0
1		317	2409	1535	411	457	6			0
1	D	317	Total	С	N	О	S	0	0	0
1	D	317	2409	1535	411	457	6	0	0	
1	Е	317	Total	С	N	О	S	0	0	0
1		317	2409	1535	411	457	6	0	U	U
1	1 E	217	Total	С	N	О	S	0	0	0
1 F	317	2409	1535	411	457	6	0	U	U	

• Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Mn 2 2	0	0
2	В	2	Total Mn 2 2	0	0
2	С	2	Total Mn 2 2	0	0
2	D	2	Total Mn 2 2	0	0
2	E	2	Total Mn 2 2	0	0
2	F	2	Total Mn 2 2	0	0

• Molecule 3 is 3-[(5 {S},7 {S},8 {S})-8-azanyl-8-carboxy-1-azaspiro[4.4]nonan-7-yl]propyl-\$  $1^{3}$ -oxidanyl-bis(oxidanyl)boranuide (three-letter code: QRA) (formula:  $C_{12}H_{24}BN_2O_5$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf		
3	A	1	Total	В	С	N	О	0	0		
3	A	1	20	1	12	2	5	U	0		
3	В	1	Total	В	С	N	О	0	0		
3	Б	1	20	1	12	2	5	U	0		
3	C	C	C	1	Total	В	С	N	О	0	0
3		1	20	1	12	2	5	0	0		
3	D	1	Total	В	С	N	О	0	0		
3	ע		20	1	12	2	5	U	0		
3	Е	1	Total	В	С	N	О	0	0		
3	Ŀ	1	20	1	12	2	5	0			
3	F	1	Total	В	С	N	О	0	0		
3	I'	1	20	1	12	2	5	U	U		

#### • Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	196	Total O 196 196	0	0
4	В	204	Total O 204 204	0	0
4	С	184	Total O 184 184	0	0
4	D	198	Total O 198 198	0	0
4	Е	166	Total O 166 166	0	0
4	F	163	Total O 163 163	0	0



MolProbity and EDS failed to run properly - this section is therefore empty.



## 3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	$53.72\text{\AA}  286.49\text{Å}  67.12\text{Å}$	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.03^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	71.62 - 1.99	Depositor
% Data completeness	98.5 (71.62-1.99)	Depositor
(in resolution range)	,	-
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.09  (at  1.98Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
$R, R_{free}$	0.214 , $0.249$	Depositor
Wilson B-factor $(A^2)$	20.5	Xtriage
Anisotropy	0.766	Xtriage
L-test for twinning <sup>2</sup>	$< L > = 0.44, < L^2> = 0.27$	Xtriage
Estimated twinning fraction	0.450 for h,-k,-l	Xtriage
Total number of atoms	15697	wwPDB-VP
Average B, all atoms $(\mathring{A}^2)$	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $< L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

### 4 Model quality (i)

### 4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3 Torsion angles (i)

#### 4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

### 4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 4.6 Ligand geometry (i)

Of 18 ligands modelled in this entry, 12 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type Chain Res Link				Вс	ond leng	ths	Bond angles		
MIOI	Mol   Type	Chain	nes	Res Link	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	QRA	Е	1003	2	13,21,21	1.12	2 (15%)	13,33,33	1.91	3 (23%)
3	QRA	A	1003	2	13,21,21	1.06	2 (15%)	13,33,33	1.85	3 (23%)
3	QRA	С	1003	2	13,21,21	1.03	1 (7%)	13,33,33	1.92	3 (23%)
3	QRA	F	1003	2	13,21,21	1.02	2 (15%)	13,33,33	1.88	3 (23%)
3	QRA	В	1003	2	13,21,21	1.10	2 (15%)	13,33,33	1.82	3 (23%)
3	QRA	D	1003	2	13,21,21	1.10	2 (15%)	13,33,33	1.92	3 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	QRA	Е	1003	2	-	0/8/39/39	0/2/2/2
3	QRA	A	1003	2	-	0/8/39/39	0/2/2/2
3	QRA	С	1003	2	-	0/8/39/39	0/2/2/2
3	QRA	F	1003	2	-	0/8/39/39	0/2/2/2
3	QRA	В	1003	2	-	0/8/39/39	0/2/2/2
3	QRA	D	1003	2	-	0/8/39/39	0/2/2/2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(\AA)$	Ideal(Å)
3	Е	1003	QRA	O1-C1	3.09	1.32	1.22
3	D	1003	QRA	O1-C1	2.95	1.31	1.22
3	A	1003	QRA	O1-C1	2.87	1.31	1.22
3	С	1003	QRA	O1-C1	2.86	1.31	1.22
3	В	1003	QRA	O1-C1	2.82	1.31	1.22
3	F	1003	QRA	O1-C1	2.62	1.30	1.22
3	D	1003	QRA	O-C1	-2.19	1.22	1.30
3	В	1003	QRA	O-C1	-2.17	1.22	1.30
3	Е	1003	QRA	O-C1	-2.11	1.22	1.30
3	F	1003	QRA	O-C1	-2.07	1.22	1.30
3	A	1003	QRA	O-C1	-2.01	1.22	1.30

All (18) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	F	1003	QRA	O-C1-C	4.10	124.67	113.60
3	С	1003	QRA	O-C1-C	4.08	124.63	113.60
3	D	1003	QRA	O-C1-C	3.98	124.34	113.60
3	A	1003	QRA	O-C1-C	3.94	124.26	113.60
3	В	1003	QRA	O-C1-C	3.84	123.98	113.60
3	${ m E}$	1003	QRA	O-C1-C	3.84	123.97	113.60
3	D	1003	QRA	O1-C1-C	-3.69	112.60	122.50
3	Ε	1003	QRA	O1-C1-C	-3.63	112.75	122.50
3	С	1003	QRA	O1-C1-C	-3.62	112.78	122.50
3	В	1003	QRA	O1-C1-C	-3.61	112.81	122.50
3	A	1003	QRA	O1-C1-C	-3.60	112.83	122.50
3	F	1003	QRA	O1-C1-C	-3.57	112.91	122.50
3	${ m E}$	1003	QRA	C5-C4-C3	3.33	108.82	105.15
3	D	1003	QRA	C5-C4-C3	3.27	108.75	105.15
3	С	1003	QRA	C5-C4-C3	3.07	108.53	105.15
3	F	1003	QRA	C5-C4-C3	2.99	108.44	105.15
3	A	1003	QRA	C5-C4-C3	2.74	108.17	105.15
3	В	1003	QRA	C5-C4-C3	2.67	108.09	105.15

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















## 4.7 Other polymers (i)

There are no such residues in this entry.

### 4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



### 5 Fit of model and data (i)

#### 5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

#### 5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

### 5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

#### 5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

### 5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

