

# wwPDB X-ray Structure Validation Summary Report (i)

Dec 8, 2023 – 04:21 am GMT

PDB ID : 2V9X

Title: E138D variant of Escherichia coli dCTP deaminase in complex with dUTP

Authors: Thymark, M.; Johansson, E.; Larsen, S.; Willemoes, M.

Deposited on : 2007-08-28

Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS: 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
			5%	
1	A	193	92%	7% •
			9%	
1	В	193	92%	8%
			6%	
1	С	193	88%	11% •
			6%	
1	D	193	93%	7%
			9%	
1	Е	193	93%	7%



Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	F	193	9%	8% •
1	G	193	93%	7%
1	Н	193	94%	5% •
1	Ι	193	90%	10% •
1	J	193	64% 17% • •	17%
1	K	193	90%	10%
1	L	193	69% 15% 69	6 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	K	1197	-	-	-	X



# 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 18359 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called DEOXYCYTIDINE TRIPHOSPHATE DEAMINASE.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Λ	102	Total	С	N	О	S	0	0	0
1	A	193	1495	940	271	278	6	U	0	0
1	D	D 102	Total	С	N	О	S	0	0	0
1	В	193	1495	940	271	278	6	U	0	0
1	С	102	Total	С	N	О	S	0	0	0
1		193	1495	940	271	278	6	U	U	U
1	D	102	Total	С	N	О	S	0	0	0
1	D	193	1495	940	271	278	6	0	0	0
1	E	193	Total	С	N	О	S	0	0	0
1		190	1495	940	271	278	6	U	0	0
1	F	193	Total	С	N	О	S	0	0	0
1	Г	193	1495	940	271	278	6	U	U	
1	G	102	Total	С	N	О	S	0	0	0
1	G	193	1495	940	271	278	6	U	0	0
1	Н	193	Total	С	N	О	S	0	0	0
1	Π	195	1495	940	271	278	6	U	U	0
1	I	193	Total	С	N	О	S	0	0	0
1	1	190	1495	940	271	278	6	U	0	0
1	J	161	Total	С	N	О	S	0	0	0
1	J	101	1247	796	220	225	6	U	0	0
1	K	102	Total	С	N	О	S	0	0	0
1	117	X 193	1495	940	271	278	6	U		U
1	L	174	Total	С	N	О	S	0	0	1
1	ь	1/4	1334	846	238	244	6	U	U	1

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	138	ASP	GLU	engineered mutation	UNP P28248
В	138	ASP	GLU	engineered mutation	UNP P28248
С	138	ASP	GLU	engineered mutation	UNP P28248
D	138	ASP	GLU	engineered mutation	UNP P28248
Е	138	ASP	GLU	engineered mutation	UNP P28248



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
F	138	ASP	GLU	engineered mutation	UNP P28248
G	138	ASP	GLU	engineered mutation	UNP P28248
Н	138	ASP	GLU	engineered mutation	UNP P28248
I	138	ASP	GLU	engineered mutation	UNP P28248
J	138	ASP	GLU	engineered mutation	UNP P28248
K	138	ASP	GLU	engineered mutation	UNP P28248
L	138	ASP	GLU	engineered mutation	UNP P28248

 $\bullet$  Molecule 2 is DEOXYURIDINE-5'-TRIPHOSPHATE (three-letter code: DUT) (formula:  $C_9H_{15}N_2O_{14}P_3).$ 



Mol	Chain	Residues		At	oms	}		ZeroOcc	AltConf	
2	A	1	Total	С	N	О	Р	0	0	
	Λ	1	28	9	2	14	3	U	U	
2	В	1	Total	С	N	Ο	Р	0	0	
	Ъ	1	28	9	2	14	3	0	U	
2	C	1	Total	С	N	Ο	Р	0	0	
		1	28	9	2	14	3		U	
2	D	D	1	Total	С	N	О	Р	0	0
2	D	1	28	9	2	14	3		U	
2	Е	1	Total	С	N	О	Р	0	0	
2	<u> 1</u> 2	1	28	9	2	14	3	0	0	
2	F	1	Total	С	N	О	Р	0	0	
2	I'	1	28	9	2	14	3		U	
2	С	G 1	Total	С	N	О	Р	0	0	
	G		28	9	2	14	3		U	



Continued from previous page...


Mol	Chain	Residues		At	oms	}		ZeroOcc	AltConf								
2	Н	1	Total	С	N	О	Р	0	0								
2	11	1	28	9	2	14	3	0	O								
2	Т	1	Total	С	N	О	Р	0	0								
2	1	1	28	9	2	14	3		U								
2	Ţ	Ţ	т	T	Т	T	Ţ	Ţ	Ţ	1	Total	С	N	О	Р	0	0
2	J	1	28	9	2	14	3		0								
2	K	1	Total	С	N	О	Р	0	0								
2	IX	1	28	9	2	14	3	0	0								
2	Т	1	Total	С	N	О	Р	0	0								
	П		28	9	2	14	3										

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	В	1	Total Mg 1 1	0	0
3	С	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0
3	Е	1	Total Mg 1 1	0	0
3	F	1	Total Mg 1 1	0	0
3	G	1	Total Mg 1 1	0	0
3	Н	1	Total Mg 1 1	0	0
3	I	1	Total Mg 1 1	0	0
3	J	1	Total Mg 1 1	0	0
3	К	1	Total Mg 1 1	0	0
3	L	1	Total Mg 1 1	0	0

 $\bullet$  Molecule 4 is SULFATE ION (three-letter code: SO4) (formula:  $\mathrm{O_4S}).$ 





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
1	С	1	Total O S	0	0
4	4   0	1	5   4   1		U
1	D	1	Total O S	0	0
4	D	1	5   4   1	0	
1	G	1	Total O S	0	0
4	G	1	5 4 1	0	0
4	I/	1	Total O S	0	0
4	N	1	5 4 1	0	0

#### • Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	44	Total O 44 44	0	0
5	В	48	Total O 48 48	0	0
5	С	54	Total O 54 54	0	0
5	D	36	Total O 36 36	0	0
5	Е	40	Total O 40 40	0	0
5	F	33	Total O 33 33	0	0
5	G	25	Total O 25 25	0	0
5	Н	45	Total O 45 45	0	0



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	I	41	Total O 41 41	0	0
5	J	32	Total O 32 32	0	0
5	K	37	Total O 37 37	0	0
5	L	25	Total O 25 25	0	0



#### 3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DEOXYCYTIDINE TRIPHOSPHATE DEAMINASE




• Molecule 1: DEOXYCYTIDINE TRIPHOSPHATE DEAMINASE











# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	94.19Å 71.30Å 319.92Å	Donogitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 - 2.20	Depositor
resolution (A)	19.80 - 2.20	EDS
% Data completeness	96.7 (20.00-2.20)	Depositor
(in resolution range)	96.7 (19.80-2.20)	EDS
$R_{merge}$	0.01	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	3.53  (at  2.19Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
P. P.	0.203 , $0.239$	Depositor
$R, R_{free}$	0.210 , 0.246	DCC
$R_{free}$ test set	5341 reflections (5.01%)	wwPDB-VP
Wilson B-factor $(\mathring{A}^2)$	25.5	Xtriage
Anisotropy	0.041	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.39, 47.6	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	18359	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.74% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, SO4, DUT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain		nd lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z >5	
1	A	0.86	0/1526	0.85	2/2069~(0.1%)	
1	В	0.89	1/1526 (0.1%)	0.86	$2/2069 \ (0.1\%)$	
1	С	0.88	0/1526	0.88	$2/2069 \ (0.1\%)$	
1	D	0.84	0/1526	0.85	1/2069~(0.0%)	
1	Е	0.88	0/1526	0.85	1/2069~(0.0%)	
1	F	0.88	0/1526	0.86	2/2069~(0.1%)	
1	G	0.82	0/1526	0.83	1/2069~(0.0%)	
1	Н	0.86	0/1526	0.84	1/2069~(0.0%)	
1	I	0.87	0/1526	0.85	1/2069~(0.0%)	
1	J	1.04	4/1275~(0.3%)	0.98	2/1733~(0.1%)	
1	K	0.85	0/1526	0.86	$2/2069 \ (0.1\%)$	
1	L	0.97	0/1364	0.97	4/1855~(0.2%)	
All	All	0.89	5/17899 (0.0%)	0.87	21/24278 (0.1%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	J	0	1
1	L	0	1
All	All	0	2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$\operatorname{Ideal}( ext{\AA})$
1	J	108	ASP	CB-CG	8.17	1.69	1.51
1	J	79	GLU	CG-CD	5.45	1.60	1.51
1	J	95	GLU	CG-CD	5.31	1.59	1.51
1	В	108	ASP	CB-CG	5.29	1.62	1.51



Continued from previous page...

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	Observed(A)	$\operatorname{Ideal}( ext{\AA})$
1	J	77	GLU	CG-CD	5.12	1.59	1.51

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$Ideal(^{o})$
1	С	174	ARG	NE-CZ-NH2	-6.24	117.18	120.30
1	F	174	ARG	NE-CZ-NH2	-6.05	117.27	120.30
1	J	126	ARG	NE-CZ-NH1	5.94	123.27	120.30
1	A	174	ARG	NE-CZ-NH2	-5.80	117.40	120.30
1	G	126	ARG	NE-CZ-NH1	5.55	123.07	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	J	66	ASP	Peptide
1	L	56	GLY	Peptide

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1495	0	1493	11	1
1	В	1495	0	1493	16	0
1	С	1495	0	1493	19	0
1	D	1495	0	1493	17	0
1	Е	1495	0	1493	7	2
1	F	1495	0	1493	13	6
1	G	1495	0	1493	10	1
1	Н	1495	0	1493	12	1
1	I	1495	0	1493	17	0
1	J	1247	0	1254	25	0
1	K	1495	0	1493	12	5
1	L	1334	0	1337	31	0
2	A	28	0	11	0	0
2	В	28	0	11	0	0
2	С	28	0	11	0	0



Continued from previous page...

Mol	Chain	Non-H		H(added)	Clashes	Symm-Clashes
2	D	28	0	11	0	0
2	Е	28	0	11	0	0
2	F	28	0	11	0	0
2	G	28	0	11	0	0
2	Н	28	0	11	0	0
2	I	28	0	11	0	0
2	J	28	0	11	0	0
2	K	28	0	11	0	0
2	L	28	0	11	0	0
3	A	1	0	0	0	0
3	В	1	0	0	0	0
3	С	1	0	0	0	0
3	D	1	0	0	0	0
3	Ε	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	Н	1	0	0	0	0
3	I	1	0	0	0	0
3	J	1	0	0	0	0
3	K	1	0	0	0	0
3	L	1	0	0	0	0
4	С	5	0	0	0	0
4	D	5	0	0	0	0
4	G	5	0	0	0	0
4	K	5	0	0	1	0
5	A	44	0	0	4	0
5	В	48	0	0	4	0
5	С	54	0	0	12	0
5	D	36	0	0	2	0
5	${ m E}$	40	0	0	3	0
5	F	33	0	0	6	0
5	G	25	0	0	1	0
5	Н	45	0	0	5	0
5	I	41	0	0	8	0
5	J	32	0	0	3	0
5	K	37	0	0	0	0
5	L	25	0	0	7	0
All	All	18359	0	17653	144	8

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 4.

The worst 5 of 144 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance}  ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:183:GLN:OE1	1:D:74:VAL:CG2	1.72	1.37
1:I:134:CYS:HB2	5:I:2037:HOH:O	1.25	1.27
1:B:183:GLN:OE1	1:D:74:VAL:HG23	1.08	1.21
1:L:55:SER:C	5:L:2009:HOH:O	1.74	1.19
1:B:183:GLN:CD	1:D:74:VAL:HG21	1.73	1.08

The worst 5 of 8 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{aligned} &  ext{Interatomic} \ &  ext{distance} \ &  ext{(Å)} \end{aligned}$	Clash overlap (Å)
1:A:183:GLN:OE1	1:F:74:VAL:CG1[1_655]	1.56	0.64
1:E:77:GLU:OE2	1:H:175:GLU:OE1[1_545]	1.91	0.29
1:F:57:PRO:CG	1:K:36:ARG:CZ[1_445]	2.00	0.20
1:F:57:PRO:CG	1:K:36:ARG:NH1[1_445]	2.06	0.14
1:E:181:ASN:OD1	1:G:77:GLU:CG[1_545]	2.12	0.08

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$191/193\ (99\%)$	188 (98%)	3 (2%)	0	100	100
1	В	$191/193\ (99\%)$	188 (98%)	3 (2%)	0	100	100
1	С	$191/193\ (99\%)$	189 (99%)	2 (1%)	0	100	100
1	D	$191/193\ (99\%)$	188 (98%)	3 (2%)	0	100	100
1	E	$191/193\ (99\%)$	188 (98%)	3 (2%)	0	100	100
1	F	$191/193\ (99\%)$	189 (99%)	2 (1%)	0	100	100
1	G	$191/193\ (99\%)$	189 (99%)	2 (1%)	0	100	100
1	Н	$191/193\ (99\%)$	188 (98%)	3 (2%)	0	100	100
1	I	$191/193\ (99\%)$	189 (99%)	2 (1%)	0	100	100
1	J	$157/193\ (81\%)$	151 (96%)	4 (2%)	2 (1%)	12	9



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	$\mathbf{s}$
1	K	191/193 (99%)	188 (98%)	3 (2%)	0	100 100	
1	L	172/193 (89%)	159 (92%)	9 (5%)	4 (2%)	6 3	
All	All	2239/2316 (97%)	2194 (98%)	39 (2%)	6 (0%)	41 46	

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	58	LYS
1	L	61	VAL
1	J	66	ASP
1	L	63	ALA
1	L	57	PRO

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	158/158 (100%)	153 (97%)	5 (3%)	39 50
1	В	158/158 (100%)	154 (98%)	4 (2%)	47 60
1	С	158/158 (100%)	152 (96%)	6 (4%)	33 42
1	D	158/158 (100%)	154 (98%)	4 (2%)	47 60
1	E	158/158 (100%)	154 (98%)	4 (2%)	47 60
1	F	158/158 (100%)	153 (97%)	5 (3%)	39 50
1	G	158/158 (100%)	154 (98%)	4 (2%)	47 60
1	Н	158/158 (100%)	153 (97%)	5 (3%)	39 50
1	I	158/158 (100%)	152 (96%)	6 (4%)	33 42
1	J	133/158 (84%)	124 (93%)	9 (7%)	16 17
1	K	158/158 (100%)	152 (96%)	6 (4%)	33 42
1	L	142/158 (90%)	129 (91%)	13 (9%)	9 9
All	All	1855/1896 (98%)	1784 (96%)	71 (4%)	33 42



5 of 71 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	K	175	GLU
1	L	6	ARG
1	L	69	MET
1	F	59	ASP
1	F	6	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	L	39	ASN
1	L	29	ASN
1	I	181	ASN
1	I	125	HIS
1	J	39	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

Of 28 ligands modelled in this entry, 12 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Во	Bond lengths			ond ang	gles
WIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	DUT	G	1194	3	25,29,29	1.01	1 (4%)	37,45,45	1.95	12 (32%)
2	DUT	J	1194	3	25,29,29	1.09	4 (16%)	37,45,45	1.44	7 (18%)
2	DUT	Е	1194	3	25,29,29	1.10	2 (8%)	37,45,45	2.44	11 (29%)
4	SO4	С	1197	-	4,4,4	0.23	0	6,6,6	0.37	0
2	DUT	D	1194	3	25,29,29	1.15	2 (8%)	37,45,45	1.77	11 (29%)
2	DUT	F	1194	3	25,29,29	1.03	1 (4%)	37,45,45	1.55	6 (16%)
2	DUT	I	1194	3	25,29,29	1.33	5 (20%)	37,45,45	1.92	9 (24%)
2	DUT	В	1194	3	25,29,29	0.89	1 (4%)	37,45,45	1.70	7 (18%)
2	DUT	A	1194	3	25,29,29	1.14	1 (4%)	37,45,45	2.30	9 (24%)
2	DUT	K	1194	3	25,29,29	1.13	2 (8%)	37,45,45	2.03	10 (27%)
2	DUT	С	1194	3	25,29,29	1.26	3 (12%)	37,45,45	1.46	10 (27%)
4	SO4	K	1197	-	4,4,4	0.17	0	6,6,6	0.12	0
4	SO4	G	1197	-	4,4,4	0.13	0	6,6,6	0.32	0
2	DUT	Н	1194	3	25,29,29	1.21	4 (16%)	37,45,45	2.07	12 (32%)
4	SO4	D	1197	-	4,4,4	0.15	0	6,6,6	0.22	0
2	DUT	L	1194	3	25,29,29	1.20	3 (12%)	37,45,45	1.77	8 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DUT	G	1194	3	-	3/22/34/34	0/2/2/2
2	DUT	J	1194	3	-	4/22/34/34	0/2/2/2
2	DUT	Е	1194	3	-	3/22/34/34	0/2/2/2
2	DUT	D	1194	3	-	4/22/34/34	0/2/2/2
2	DUT	F	1194	3	-	4/22/34/34	0/2/2/2
2	DUT	I	1194	3	-	4/22/34/34	0/2/2/2
2	DUT	В	1194	3	-	4/22/34/34	0/2/2/2
2	DUT	A	1194	3	-	4/22/34/34	0/2/2/2
2	DUT	K	1194	3	-	4/22/34/34	0/2/2/2
2	DUT	С	1194	3	-	4/22/34/34	0/2/2/2
2	DUT	Н	1194	3	-	4/22/34/34	0/2/2/2
2	DUT	L	1194	3	-	2/22/34/34	0/2/2/2

The worst 5 of 29 bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	Observed(A)	$\operatorname{Ideal}(\text{\AA})$
2	I	1194	DUT	C2-N1	3.47	1.44	1.38
2	A	1194	DUT	C2-N1	3.19	1.43	1.38
2	F	1194	DUT	C4-N3	-3.08	1.33	1.38
2	L	1194	DUT	C2-N1	3.06	1.43	1.38
2	С	1194	DUT	O2-C2	2.95	1.28	1.23

The worst 5 of 112 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\mathrm{Ideal}(^{o})$
2	A	1194	DUT	C4-N3-C2	-7.54	116.63	126.58
2	Е	1194	DUT	C4-N3-C2	-6.95	117.41	126.58
2	A	1194	DUT	N3-C2-N1	5.74	122.50	114.89
2	Е	1194	DUT	N3-C2-N1	5.62	122.35	114.89
2	K	1194	DUT	O4-C4-C5	-5.45	115.58	125.16

There are no chirality outliers.

5 of 44 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	J	1194	DUT	PB-O3B-PG-O3G
2	A	1194	DUT	PG-O3B-PB-O1B
2	В	1194	DUT	PG-O3B-PB-O1B
2	D	1194	DUT	PA-O3A-PB-O1B
2	Е	1194	DUT	PG-O3B-PB-O1B


There are no ring outliers.

1 monomer is involved in 1 short contact:


$\mathbf{Mol}$	Chain	Res	Type	Clashes	Symm-Clashes
4	K	1197	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

















Rings


Torsions






Rings

Torsions







## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	193/193 (100%)	0.14	9 (4%) 31 30	13, 25, 46, 55	0
1	В	193/193 (100%)	0.27	18 (9%) 8 7	13, 25, 46, 55	0
1	С	193/193 (100%)	0.23	11 (5%) 23 22	13, 25, 45, 54	0
1	D	193/193 (100%)	0.14	12 (6%) 20 19	13, 25, 46, 54	0
1	E	193/193 (100%)	0.24	17 (8%) 10 8	13, 25, 46, 55	0
1	F	193/193 (100%)	0.18	17 (8%) 10 8	14, 25, 46, 54	0
1	G	193/193 (100%)	0.09	19 (9%) 7 6	13, 25, 46, 55	0
1	Н	193/193 (100%)	0.10	12 (6%) 20 19	13, 25, 45, 54	0
1	I	193/193 (100%)	0.05	11 (5%) 23 22	13, 25, 46, 55	0
1	J	161/193 (83%)	-0.14	7 (4%) 35 33	13, 24, 41, 56	0
1	K	193/193 (100%)	0.11	15 (7%) 13 11	13, 25, 46, 54	0
1	L	174/193 (90%)	0.33	16 (9%) 9 7	13, 30, 54, 62	0
All	All	2265/2316 (97%)	0.15	164 (7%) 15 14	13, 25, 47, 62	0

The worst 5 of 164 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	183	GLN	9.4
1	В	181	ASN	8.3
1	A	77	GLU	7.8
1	L	63	ALA	6.9
1	С	183	GLN	6.8

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
4	SO4	K	1197	5/5	0.70	0.90	174,174,174,175	0
3	MG	J	1195	1/1	0.84	0.07	29,29,29,29	0
4	SO4	G	1197	5/5	0.89	0.13	76,77,78,78	0
3	MG	G	1195	1/1	0.89	0.21	33,33,33,33	0
3	MG	I	1195	1/1	0.90	0.17	24,24,24,24	0
3	MG	D	1195	1/1	0.91	0.20	26,26,26,26	0
4	SO4	С	1197	5/5	0.92	0.12	68,69,70,71	0
3	MG	Н	1195	1/1	0.93	0.21	25,25,25,25	0
4	SO4	D	1197	5/5	0.94	0.09	76,77,77,79	0
3	MG	L	1195	1/1	0.94	0.15	32,32,32,32	0
3	MG	В	1195	1/1	0.94	0.14	23,23,23,23	0
3	MG	A	1195	1/1	0.95	0.26	21,21,21,21	0
2	DUT	J	1194	28/28	0.95	0.11	18,24,33,38	0
2	DUT	L	1194	28/28	0.96	0.09	15,21,32,35	0
3	MG	С	1195	1/1	0.96	0.08	15,15,15,15	0
3	MG	Е	1195	1/1	0.97	0.08	20,20,20,20	0
2	DUT	G	1194	28/28	0.97	0.09	13,22,27,28	0
3	MG	K	1195	1/1	0.98	0.02	17,17,17,17	0
3	MG	F	1195	1/1	0.98	0.09	17,17,17,17	0
2	DUT	F	1194	28/28	0.98	0.09	10,16,20,24	0
2	DUT	K	1194	28/28	0.98	0.09	11,16,20,23	0
2	DUT	D	1194	28/28	0.98	0.09	12,18,24,25	0
2	DUT	Н	1194	28/28	0.98	0.09	12,17,20,22	0
2	DUT	I	1194	28/28	0.99	0.08	8,17,18,21	0
2	DUT	Е	1194	28/28	0.99	0.07	10,14,19,22	0
2	DUT	В	1194	28/28	0.99	0.08	8,15,19,22	0
2	DUT	С	1194	28/28	0.99	0.07	8,13,15,18	0
2	DUT	A	1194	28/28	0.99	0.09	11,15,18,21	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different




orientation to approximate a three-dimensional view.







# Electron density around DUT G 1194:

 $2 {
m mF}_o {
m -DF}_c$  (at 0.7 rmsd) in gray  ${
m mF}_o {
m -DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around DUT F 1194:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)














 $2 {
m mF}_o {
m -DF}_c$  (at 0.7 rmsd) in gray  ${
m mF}_o {
m -DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)












#### Electron density around DUT H 1194:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around DUT I 1194:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)












## Electron density around DUT E 1194:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around DUT B 1194:

 $2 \text{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\text{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)












#### Electron density around DUT C 1194:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around DUT A 1194:

 $2 \text{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\text{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)









# 6.5 Other polymers (i)

There are no such residues in this entry.

