



Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 10:43 am BST

PDB ID : 3VAE
Title : Crystal Structure of *M. tuberculosis* LD-transpeptidase type 2 with Modified Catalytic Cysteine (C354)
Authors : Erdemli, S.; Bianchet, M.A.; Gupta, R.; Lamichhane, G.; Amzel, L.M.
Deposited on : 2011-12-29
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	287	 63% 27% • 7%
1	B	287	 71% 17% • 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PEG	A	501	-	-	-	X

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4244 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LD-transpeptidase type 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	266	Total 2071	C 1305	N 360	O 399	S 7	0	4	0
1	B	258	Total 2005	C 1265	N 345	O 387	S 8	0	4	0

- Molecule 2 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 7	C 4	O 3	0	0
2	B	1	Total 7	C 4	O 3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	76	Total 76	O 76	0	0
3	B	78	Total 78	O 78	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	I 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	119.04Å 121.22Å 122.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.84 – 2.80 19.84 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (19.84-2.80) 99.8 (19.84-2.80)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.66 (at 2.79Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.187 , 0.258 0.182 , 0.252	Depositor DCC
R_{free} test set	1127 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	28.8	Xtriage
Anisotropy	0.454	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 35.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.116 for -h,-l,-k 0.014 for l,-k,h 0.015 for -k,-h,-l 0.007 for k,-l,-h 0.007 for -l,h,-k	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4244	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CSO, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.87	0/2128	0.87	6/2908 (0.2%)
1	B	0.89	0/2065	0.84	2/2822 (0.1%)
All	All	0.88	0/4193	0.86	8/5730 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	258	ASP	CB-CG-OD2	-6.03	112.87	118.30
1	A	399[A]	ASP	CB-CA-C	-5.35	99.69	110.40
1	A	399[B]	ASP	CB-CA-C	-5.35	99.69	110.40
1	A	232	ASP	CB-CG-OD1	5.25	123.03	118.30
1	A	130	ALA	N-CA-CB	-5.24	102.77	110.10
1	B	211	ARG	NE-CZ-NH2	-5.23	117.68	120.30
1	B	152	THR	CB-CA-C	-5.21	97.55	111.60
1	A	283	ASP	CB-CG-OD1	5.09	122.89	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2071	0	1969	57	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2005	0	1911	37	0
2	A	7	0	10	0	0
2	B	7	0	10	0	0
3	A	76	0	0	3	0
3	B	78	0	0	5	0
All	All	4244	0	3900	93	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:398:TRP:O	1:A:399[B]:ASP:HB3	1.41	1.15
1:A:152:THR:HG23	1:A:176:GLU:HB2	1.34	1.10
1:B:289:ASN:HD21	1:B:380:THR:H	1.05	1.00
1:A:289:ASN:HD21	1:A:380:THR:H	1.10	0.99
1:B:204:ASN:HD22	1:B:206:ARG:H	1.16	0.92
1:A:398:TRP:O	1:A:399[B]:ASP:CB	2.18	0.91
1:A:398:TRP:O	1:A:399[A]:ASP:HB2	1.61	0.90
1:A:153:MET:HE2	1:A:154:PRO:HD2	1.53	0.88
1:A:213:GLU:HG3	1:A:398:TRP:CG	2.09	0.87
1:A:153:MET:CE	1:A:154:PRO:HD2	2.04	0.86
1:A:181:ARG:HH21	1:A:205:ASN:ND2	1.76	0.84
1:A:152:THR:CG2	1:A:176:GLU:HB2	2.09	0.83
1:B:289:ASN:HD21	1:B:380:THR:N	1.80	0.80
1:B:204:ASN:ND2	1:B:206:ARG:H	1.85	0.74
1:A:152:THR:HG23	1:A:176:GLU:CB	2.16	0.74
1:A:291:ILE:H	1:A:395:ASN:HD21	1.35	0.74
1:A:151:LEU:HA	3:A:636:HOH:O	1.89	0.71
1:A:281:GLY:HA2	1:A:351:SER:O	1.92	0.69
1:B:289:ASN:ND2	1:B:380:THR:H	1.86	0.69
1:B:297:ARG:HG3	1:B:297:ARG:HH11	1.60	0.67
1:B:198:GLY:HA3	1:B:210:TRP:CZ2	2.30	0.66
1:B:203:LEU:HD22	1:B:388:ILE:HG22	1.77	0.64
1:A:181:ARG:HH21	1:A:205:ASN:HD22	1.45	0.63
1:B:291:ILE:H	1:B:395:ASN:HD21	1.47	0.63
1:B:152:THR:HG22	1:B:175:ASP:OD1	1.99	0.62
1:B:297:ARG:NH1	1:B:297:ARG:HG3	2.13	0.61
1:A:125:THR:HG21	1:A:244:GLN:HG3	1.81	0.60
1:A:297[B]:ARG:NH2	1:A:366:TYR:O	2.34	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:153:MET:HE3	1:A:154:PRO:HD2	1.83	0.59
1:B:150:HIS:HB3	3:B:646:HOH:O	2.03	0.59
1:A:406:ALA:O	1:A:407:LYS:HB2	2.01	0.59
1:B:198:GLY:HA3	1:B:210:TRP:CH2	2.38	0.58
1:A:246:HIS:HB2	3:A:635:HOH:O	2.02	0.58
1:B:156:VAL:HG23	1:B:172:ILE:HG13	1.84	0.58
1:B:213:GLU:HB2	1:B:398:TRP:CE2	2.40	0.56
1:A:213:GLU:HG3	1:A:398:TRP:CD1	2.39	0.56
1:B:387:GLY:HA3	1:B:396:ILE:HG13	1.87	0.55
1:B:152:THR:HG23	1:B:176:GLU:HG2	1.88	0.55
1:A:130:ALA:HB2	1:A:159:GLY:HA3	1.89	0.55
1:B:255:ALA:HB1	1:B:264:LEU:HD11	1.89	0.54
1:A:365:PHE:O	1:A:369:VAL:HB	2.08	0.54
1:A:228:THR:O	1:A:239:GLY:HA3	2.08	0.53
1:A:157:MET:HB2	1:A:171:ALA:HB3	1.91	0.53
1:A:398:TRP:O	1:A:399[A]:ASP:CB	2.30	0.53
1:A:399[B]:ASP:OD1	1:A:400:GLN:N	2.42	0.53
1:B:246:HIS:HB2	3:B:630:HOH:O	2.09	0.53
1:B:370:LYS:HE3	3:B:607:HOH:O	2.09	0.52
1:A:340:TRP:HB2	1:B:320:THR:HG23	1.92	0.52
1:A:212:PRO:HB2	1:A:214:HIS:O	2.10	0.51
1:A:189:LYS:HB3	1:A:225:ALA:HB3	1.92	0.51
1:B:203:LEU:CD2	1:B:388:ILE:HG22	2.39	0.51
1:A:196[B]:VAL:HG21	3:A:641:HOH:O	2.09	0.51
1:B:330:TYR:CE1	1:B:391:LEU:HG	2.47	0.50
1:B:224:VAL:HB	1:B:245:THR:HG22	1.93	0.50
1:A:399[B]:ASP:CG	1:A:400:GLN:N	2.58	0.49
1:A:224:VAL:HB	1:A:245:THR:HG22	1.94	0.49
1:B:406:ALA:O	1:B:407:LYS:HB2	2.12	0.49
1:A:181:ARG:NH2	1:A:205:ASN:ND2	2.54	0.49
1:A:153:MET:HE2	1:A:154:PRO:CD	2.36	0.49
1:A:292:TYR:CD1	1:A:385:LEU:HD22	2.47	0.49
1:B:281:GLY:O	1:B:349:ASN:HB3	2.13	0.49
1:A:152:THR:HG22	1:A:175:ASP:OD1	2.12	0.49
1:A:198:GLY:HA3	1:A:210:TRP:CH2	2.47	0.48
1:B:167:GLY:HA3	1:B:374:ILE:HD11	1.95	0.48
1:A:172:ILE:HD13	1:A:188:ILE:HD13	1.96	0.47
1:A:193:ASN:HA	1:A:194:PRO:C	2.34	0.47
1:A:198:GLY:HA3	1:A:210:TRP:CZ2	2.49	0.47
1:A:276:MET:SD	1:A:361:ASN:HB3	2.54	0.47
1:A:324:TRP:CE2	1:A:363:GLN:HG3	2.49	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:330:TYR:CD2	1:A:390:GLY:HA3	2.50	0.47
1:B:302:ILE:HA	1:B:320:THR:O	2.16	0.45
1:B:337:SER:HB3	3:B:631:HOH:O	2.16	0.44
1:B:253:VAL:HB	1:B:373:ASP:CG	2.38	0.44
1:A:211:ARG:HG2	1:A:401:TRP:CZ2	2.53	0.43
1:A:274:LYS:HE3	1:A:364:TRP:CD1	2.52	0.43
1:A:302:ILE:HA	1:A:320:THR:O	2.18	0.43
1:A:158:PRO:HD2	1:A:247:PHE:CE2	2.54	0.43
1:B:402:ARG:HB3	1:B:402:ARG:HE	1.67	0.43
1:A:313:ASN:CB	3:B:653:HOH:O	2.66	0.43
1:A:378:VAL:O	1:A:379:ASN:HB2	2.18	0.42
1:B:267:ARG:HA	1:B:271:GLU:O	2.20	0.42
1:B:152:THR:HG23	1:B:176:GLU:CG	2.49	0.42
1:B:337:SER:OG	1:B:359:PRO:HA	2.20	0.42
1:A:213:GLU:HG3	1:A:398:TRP:CB	2.46	0.42
1:A:292:TYR:CE1	1:A:385:LEU:HD22	2.55	0.42
1:B:297:ARG:CG	1:B:297:ARG:HH11	2.29	0.42
1:B:281:GLY:HA2	1:B:351:SER:O	2.20	0.42
1:A:167:GLY:HA3	1:A:374:ILE:HD11	2.02	0.41
1:A:255:ALA:HB1	1:A:264:LEU:HD11	2.02	0.41
1:A:152:THR:OG1	1:A:237:MET:CE	2.68	0.41
1:A:341:SER:O	1:A:345:GLN:HG3	2.21	0.41
1:B:152:THR:CG2	1:B:175:ASP:OD1	2.68	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	265/287 (92%)	246 (93%)	17 (6%)	2 (1%)	19 49
1	B	259/287 (90%)	243 (94%)	16 (6%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	524/574 (91%)	489 (93%)	33 (6%)	2 (0%)	47 66

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	399[A]	ASP
1	A	399[B]	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	220/231 (95%)	201 (91%)	19 (9%)	10 30
1	B	215/231 (93%)	202 (94%)	13 (6%)	19 48
All	All	435/462 (94%)	403 (93%)	32 (7%)	16 37

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	124	TYR
1	A	125	THR
1	A	152	THR
1	A	156	VAL
1	A	206	ARG
1	A	217	LYS
1	A	244	GLN
1	A	275	SER
1	A	297[A]	ARG
1	A	297[B]	ARG
1	A	356	ASN
1	A	359	PRO
1	A	369	VAL
1	A	371[A]	ARG
1	A	371[B]	ARG
1	A	393	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	399[A]	ASP
1	A	399[B]	ASP
1	A	407	LYS
1	B	152	THR
1	B	173	ARG
1	B	186	LYS
1	B	189	LYS
1	B	192	THR
1	B	204	ASN
1	B	275	SER
1	B	296	SER
1	B	297	ARG
1	B	370	LYS
1	B	393	ASP
1	B	399[A]	ASP
1	B	399[B]	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	205	ASN
1	A	289	ASN
1	A	356	ASN
1	A	379	ASN
1	A	395	ASN
1	B	204	ASN
1	B	289	ASN
1	B	361	ASN
1	B	379	ASN
1	B	395	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSO	B	354	1	3,6,7	0.84	0	0,6,8	0.00	-
1	CSO	A	354	1	3,6,7	0.64	0	0,6,8	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	B	354	1	-	0/1/5/7	-
1	CSO	A	354	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PEG	A	501	-	6,6,6	0.56	0	5,5,5	0.26	0
2	PEG	B	501	-	6,6,6	0.52	0	5,5,5	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	A	501	-	-	3/4/4/4	-
2	PEG	B	501	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	PEG	O2-C3-C4-O4
2	B	501	PEG	O1-C1-C2-O2
2	B	501	PEG	C1-C2-O2-C3
2	A	501	PEG	C1-C2-O2-C3
2	A	501	PEG	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	265/287 (92%)	-0.63	3 (1%) 80 75	5, 17, 45, 94	0
1	B	257/287 (89%)	-0.71	0 100 100	7, 16, 41, 52	0
All	All	522/574 (90%)	-0.67	3 (0%) 89 86	5, 17, 43, 94	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	124	TYR	3.5
1	A	236	GLY	2.5
1	A	123	ARG	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSO	B	354	7/8	0.98	0.12	7,7,11,12	0
1	CSO	A	354	7/8	0.98	0.11	5,6,9,16	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PEG	A	501	7/7	0.69	0.44	80,83,84,84	0
2	PEG	B	501	7/7	0.86	0.18	60,63,65,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.