



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2023 – 10:29 AM EDT

PDB ID : 1VDK
Title : Crystal structure of fumarase from thermus thermophilus HB8
Authors : Mizutani, H.; Kunishima, N.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2004-03-23
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	466	 11% 82% 16% .
1	B	466	 7% 81% 17% .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7680 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fumarate hydratase class II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	458	3515	2231	617	653	14	0	0	0
1	B	457	3517	2234	617	652	14	0	0	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	357	Total	O	0	0
			357	357		
2	B	291	Total	O	0	0
			291	291		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fumarate hydratase class II



- Molecule 1: Fumarate hydratase class II



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	125.34Å 140.12Å 135.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.80 29.95 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.00-1.80) 100.0 (29.95-1.80)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.31 (at 1.80Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.185 , 0.201 0.185 , 0.190	Depositor DCC
R_{free} test set	5490 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	23.5	Xtrriage
Anisotropy	0.116	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 55.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7680	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/3580	0.57	1/4853 (0.0%)
1	B	0.28	0/3582	0.54	0/4855
All	All	0.29	0/7162	0.55	1/9708 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	128	ALA	N-CA-C	-5.43	96.33	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	367	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3515	0	3557	55	0
1	B	3517	0	3563	48	2
2	A	357	0	0	2	0
2	B	291	0	0	7	0
All	All	7680	0	7120	102	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:7:ARG:HG2	1:B:12:GLU:HG2	1.45	0.94
1:A:428:ALA:HB3	1:A:435:LEU:HD23	1.64	0.79
1:B:436:LYS:HD3	1:B:447:GLU:HG3	1.65	0.79
1:A:440:LEU:HD21	1:A:447:GLU:N	2.03	0.73
1:A:446:THR:HG23	1:A:449:GLU:H	1.55	0.72
1:B:113:ASN:HB3	1:B:123:LEU:HD22	1.72	0.71
1:B:145:PRO:HA	1:B:148:MET:HG2	1.73	0.70
1:A:145:PRO:HA	1:A:148:MET:HG2	1.75	0.68
1:A:105:MET:O	1:A:109:GLU:HG3	1.99	0.62
1:A:196:THR:OG1	1:A:199:GLN:HG3	2.00	0.61
1:A:402:LEU:HG	1:A:408:LEU:HD11	1.81	0.61
1:A:434:THR:HG23	1:A:437:GLN:HB2	1.81	0.61
1:A:122:PRO:HG2	1:A:125:SER:HB3	1.81	0.61
1:A:420:LYS:N	1:A:420:LYS:HD2	2.15	0.61
1:B:248:LYS:O	1:B:252:GLU:HG3	2.01	0.60
1:A:440:LEU:HD21	1:A:447:GLU:H	1.70	0.56
1:A:7:ARG:HB3	1:A:12:GLU:HG2	1.87	0.56
1:B:260:VAL:HG11	1:B:264:ARG:HE	1.70	0.56
1:B:303:TYR:HA	1:B:397:ARG:NH2	2.20	0.56
1:B:320:ILE:N	1:B:320:ILE:HD12	2.21	0.56
1:B:16:PRO:HB2	1:B:19:LYS:HG2	1.88	0.55
1:A:408:LEU:HD22	1:A:456:PRO:HG3	1.89	0.55
1:A:172:THR:O	1:A:176:GLN:HG3	2.07	0.54
1:B:172:THR:O	1:B:176:GLN:HG3	2.07	0.54
1:A:229:GLY:HA3	2:A:696:HOH:O	2.08	0.54
1:B:264:ARG:O	1:B:268:LEU:HD13	2.08	0.53
1:B:105:MET:O	1:B:109:GLU:HG3	2.09	0.53
1:A:436:LYS:HE2	1:A:447:GLU:HB2	1.90	0.53
1:A:407:MET:HB3	1:A:459:LEU:HD13	1.91	0.52
1:B:387:LEU:HD23	1:B:387:LEU:C	2.30	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:387:LEU:C	1:A:387:LEU:HD23	2.29	0.52
1:B:282:ARG:HD2	2:B:583:HOH:O	2.10	0.52
1:A:114:ARG:O	1:A:118:ILE:HG13	2.10	0.52
1:A:236:LEU:O	1:A:237:ASN:HB2	2.10	0.52
1:B:446:THR:C	1:B:448:GLU:H	2.13	0.52
1:B:260:VAL:CG1	1:B:264:ARG:HE	2.22	0.51
1:A:428:ALA:HB2	1:A:438:ALA:HB2	1.93	0.51
1:A:362:LYS:HB2	1:A:363:PRO:HD3	1.93	0.50
1:A:410:THR:HA	1:A:413:ASN:HD22	1.77	0.50
1:A:447:GLU:OE2	1:A:448:GLU:HG3	2.11	0.50
1:B:92:LEU:HD13	1:B:102:GLN:HB3	1.94	0.50
1:B:236:LEU:O	1:B:237:ASN:HB2	2.10	0.50
1:A:440:LEU:HD11	1:A:447:GLU:HB3	1.93	0.50
1:A:427:LYS:HE2	1:A:441:GLU:HB2	1.94	0.49
1:B:161:TYR:HB2	1:B:162:PRO:HD3	1.94	0.49
1:B:196:THR:OG1	1:B:199:GLN:HG3	2.11	0.49
1:A:126:LYS:HA	2:A:693:HOH:O	2.13	0.48
1:B:362:LYS:HB2	1:B:363:PRO:HD3	1.94	0.48
1:B:427:LYS:HD2	1:B:442:LEU:HD21	1.95	0.48
1:A:324:LYS:HE2	1:B:191:ASP:O	2.14	0.48
1:B:306:ILE:HG22	1:B:308:GLU:HG3	1.96	0.48
1:B:420:LYS:HG3	2:B:694:HOH:O	2.14	0.48
1:A:180:GLN:HE21	1:A:180:GLN:HA	1.79	0.48
1:A:188:HIS:O	1:A:189:LEU:HB2	2.14	0.47
1:B:197:LEU:CD2	1:B:393:PRO:HB3	2.45	0.47
1:B:52:LEU:C	1:B:52:LEU:HD23	2.34	0.47
1:B:80:GLU:HG2	1:B:85:LYS:HD2	1.95	0.47
1:B:184:VAL:HG13	1:B:407:MET:CE	2.45	0.47
1:B:223:TYR:CD1	1:B:257:PRO:HG2	2.49	0.47
1:A:434:THR:CG2	1:A:437:GLN:HB2	2.46	0.46
1:A:436:LYS:CE	1:A:447:GLU:HB2	2.46	0.46
1:A:427:LYS:HD3	1:A:438:ALA:HB1	1.97	0.46
1:B:420:LYS:O	1:B:424:ILE:HG13	2.15	0.46
1:A:180:GLN:HA	1:A:180:GLN:NE2	2.30	0.46
1:A:52:LEU:C	1:A:52:LEU:HD23	2.36	0.46
1:B:105:MET:HG3	2:B:723:HOH:O	2.16	0.46
1:B:187:THR:O	1:B:188:HIS:HB2	2.16	0.46
1:B:114:ARG:O	1:B:118:ILE:HG13	2.16	0.45
1:A:25:THR:OG1	1:A:106:ASN:HA	2.16	0.45
1:A:223:TYR:CD1	1:A:257:PRO:HG2	2.51	0.45
1:B:446:THR:C	1:B:448:GLU:N	2.71	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:399:GLU:HG3	2:B:612:HOH:O	2.18	0.44
1:A:97:THR:HG21	1:A:101:THR:HB	1.99	0.44
1:A:306:ILE:HG22	1:A:308:GLU:HG3	2.00	0.44
1:A:455:VAL:HB	1:A:458:ARG:HD2	1.99	0.44
1:A:161:TYR:HB2	1:A:162:PRO:HD3	2.00	0.43
1:A:339:ARG:NH1	1:A:343:ASN:OD1	2.51	0.43
1:B:326:ASN:HB3	2:B:591:HOH:O	2.17	0.43
1:A:28:SER:HB2	1:A:102:GLN:HG2	2.01	0.43
1:B:428:ALA:HB1	1:B:433:LYS:O	2.19	0.43
1:A:446:THR:HG22	1:A:449:GLU:OE1	2.18	0.43
1:A:43:LEU:O	1:A:47:ARG:HG2	2.19	0.42
1:B:129:HIS:HA	1:B:130:PRO:HD3	1.91	0.42
1:B:145:PRO:HB3	1:B:271:HIS:CE1	2.55	0.42
1:B:66:LEU:HD11	2:B:531:HOH:O	2.19	0.42
1:A:434:THR:HG23	1:A:437:GLN:CB	2.49	0.42
1:A:125:SER:O	1:A:126:LYS:HB2	2.20	0.42
1:A:86:TRP:CZ2	1:A:114:ARG:HD3	2.54	0.42
1:B:125:SER:O	1:B:126:LYS:HB2	2.19	0.42
1:A:397:ARG:HD2	1:A:401:TYR:CZ	2.55	0.42
1:A:427:LYS:HE3	1:A:441:GLU:OE1	2.20	0.42
1:B:4:ARG:HG2	1:B:4:ARG:HH11	1.85	0.41
1:A:442:LEU:HB3	1:A:444:TYR:CD1	2.55	0.41
1:B:144:PHE:N	1:B:145:PRO:HD2	2.35	0.41
1:A:420:LYS:N	1:A:420:LYS:CD	2.84	0.41
1:A:187:THR:O	1:A:188:HIS:HB2	2.21	0.41
1:B:405:ASN:HA	1:B:406:PRO:HD3	1.86	0.41
1:A:399:GLU:OE2	1:A:403:GLN:NE2	2.53	0.41
1:B:126:LYS:HA	2:B:751:HOH:O	2.21	0.41
1:A:455:VAL:HB	1:A:458:ARG:CD	2.51	0.40
1:B:226:ALA:HB1	1:B:268:LEU:HD12	2.02	0.40
1:B:401:TYR:O	1:B:405:ASN:HB2	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:339:ARG:NH2	1:B:339:ARG:NH2[3_655]	1.75	0.45
1:B:452:ARG:NH1	1:B:452:ARG:NH1[4_555]	1.83	0.37

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	454/466 (97%)	431 (95%)	23 (5%)	0	100	100
1	B	453/466 (97%)	439 (97%)	14 (3%)	0	100	100
All	All	907/932 (97%)	870 (96%)	37 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	361/367 (98%)	356 (99%)	5 (1%)	67	59
1	B	361/367 (98%)	355 (98%)	6 (2%)	60	51
All	All	722/734 (98%)	711 (98%)	11 (2%)	65	56

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	TYR
1	A	91	PRO
1	A	106	ASN
1	A	349	PHE
1	A	447	GLU
1	B	49	TYR
1	B	91	PRO
1	B	106	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	259	ARG
1	B	315	GLU
1	B	349	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	GLN
1	A	131	ASN
1	A	180	GLN
1	A	437	GLN
1	B	31	ASN
1	B	326	ASN
1	B	329	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	458/466 (98%)	0.42	51 (11%) 5 4	13, 23, 76, 80	0
1	B	457/466 (98%)	0.22	31 (6%) 17 13	15, 28, 58, 72	0
All	All	915/932 (98%)	0.32	82 (8%) 9 7	13, 26, 68, 80	0

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	435	LEU	12.1
1	A	445	LEU	10.0
1	A	428	ALA	9.8
1	A	444	TYR	9.2
1	A	429	LEU	8.5
1	A	440	LEU	8.5
1	A	453	ILE	7.9
1	A	439	ALA	7.6
1	A	421	ALA	7.5
1	A	319	SER	7.3
1	B	316	PRO	7.3
1	A	426	LYS	7.1
1	A	430	LYS	6.7
1	B	321	MET	6.5
1	A	447	GLU	6.4
1	A	438	ALA	6.4
1	A	425	VAL	6.1
1	A	422	ALA	6.0
1	A	414	LYS	5.9
1	A	427	LYS	5.5
1	A	442	LEU	5.5
1	B	5	ILE	5.4
1	A	441	GLU	5.3
1	A	419	ASP	5.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	318	SER	5.2
1	A	423	GLU	5.1
1	A	420	LYS	5.0
1	A	437	GLN	4.9
1	A	450	PHE	4.9
1	A	413	ASN	4.8
1	A	424	ILE	4.8
1	A	443	GLY	4.7
1	B	9	THR	4.7
1	A	433	LYS	4.6
1	B	446	THR	4.5
1	A	418	TYR	4.4
1	B	448	GLU	4.4
1	A	446	THR	4.4
1	B	430	LYS	4.3
1	B	10	MET	4.2
1	A	448	GLU	4.1
1	A	434	THR	4.1
1	A	451	ASP	3.9
1	A	449	GLU	3.8
1	A	410	THR	3.7
1	A	458	ARG	3.7
1	B	14	ARG	3.6
1	B	453	ILE	3.6
1	B	440	LEU	3.5
1	B	414	LYS	3.4
1	A	14	ARG	3.4
1	A	322	PRO	3.4
1	B	413	ASN	3.1
1	B	322	PRO	3.0
1	B	447	GLU	3.0
1	B	320	ILE	2.9
1	A	407	MET	2.9
1	A	452	ARG	2.9
1	B	443	GLY	2.8
1	B	122	PRO	2.8
1	A	436	LYS	2.8
1	A	320	ILE	2.8
1	B	418	TYR	2.6
1	B	315	GLU	2.6
1	A	417	GLY	2.6
1	B	444	TYR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	12	GLU	2.5
1	A	463	HIS	2.5
1	B	449	GLU	2.4
1	A	408	LEU	2.4
1	A	412	LEU	2.3
1	A	415	ALA	2.3
1	B	445	LEU	2.3
1	A	455	VAL	2.3
1	B	15	VAL	2.2
1	B	450	PHE	2.2
1	B	452	ARG	2.2
1	B	432	LYS	2.2
1	A	454	VAL	2.2
1	B	415	ALA	2.1
1	B	68	GLU	2.1
1	B	438	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.