Full wwPDB X-ray Structure Validation Report (i)

May 16, 2020 – 02:40 am BST

PDB ID	:	1VFB
Title	:	BOUND WATER MOLECULES AND CONFORMATIONAL STABILIZATION HELP MEDIATE AN ANTIGEN-ANTIBODY ASSOCIATION
Authors	:	Bhat, T.N.; Poljak, R.J.
Deposited on	:	1993-12-03
Resolution	:	1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	107	92% 7% •
2	B	116	83% 16% •
3	C	129	86% 13% •

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3456 atoms, of which 677 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IGG1-KAPPA D1.3 FV (LIGHT CHAIN).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	107	1020	523	194	138	163	2	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	3	VAL	GLU	CONFLICT	GB 545862
A	50	TYR	LYS	CONFLICT	GB 545862
A	51	THR	ALA	CONFLICT	GB 545862
A	52	THR	GLN	CONFLICT	GB 545862
A	?	-	PRO	DELETION	GB 545862
A	96	ARG	TRP	CONFLICT	GB 545862

- Molecule 2 is a protein called IGG1-KAPPA D1.3 FV (HEAVY CHAIN).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	116	1123	564	219	157	179	4	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	112	LEU	VAL	CONFLICT	GB 896294

- Molecule 3 is a protein called HEN EGG WHITE LYSOZYME.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
3	C	129	1265	613	264	193	185	10	0	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	8	8	8	0	0
			Total	O		
4	B	14	14	14	0	0
			Total	O		
4	C	26	26	26	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: IGG1-KAPPA D1.3 FV (LIGHT CHAIN)

Chain A:



- Molecule 2: IGG1-KAPPA D1.3 FV (HEAVY CHAIN)

Chain B:

- Molecule 3: HEN EGG WHITE LYSOZYME

Chain C:



4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	129.20Å 60.80Å 56.90Å 90.00° 119.30° 90.00°	Depositor
Resolution (Å)	(Not available) – 1.80	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-1.80)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.185 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3456	wwPDB-VP
Average B, all atoms ($Å^2$)	24.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.72	0/847	1.34	6/1150 (0.5%)
2	B	0.75	0/924	1.50	17/1252 (1.4%)
3	C	0.76	0/1021	1.58	20/1379 (1.5%)
All	All	0.74	0/2792	1.48	43/3781 (1.1%)

There are no bond length outliers.

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	C	73	ARG	NE-CZ-NH2	-8.38	116.11	120.30
3	C	123	TRP	CD1-CG-CD2	8.34	112.97	106.30
2	B	99	ARG	NE-CZ-NH2	-8.22	116.19	120.30
3	C	28	TRP	CD1-CG-CD2	8.16	112.83	106.30
3	C	62	TRP	CD1-CG-CD2	8.08	112.76	106.30
2	B	36	TRP	CD1-CG-CD2	7.98	112.68	106.30
1	A	92	TRP	CD1-CG-CD2	7.87	112.60	106.30
3	C	28	TRP	CE2-CD2-CG	-7.64	101.19	107.30
2	B	47	TRP	CD1-CG-CD2	7.62	112.40	106.30
2	B	52	TRP	CD1-CG-CD2	7.61	112.39	106.30
1	A	35	TRP	CD1-CG-CD2	7.61	112.38	106.30
2	B	106	TRP	CD1-CG-CD2	7.56	112.35	106.30
1	A	92	TRP	CE2-CD2-CG	-7.49	101.31	107.30
3	C	63	TRP	CD1-CG-CD2	7.35	112.18	106.30
3	C	111	TRP	CE2-CD2-CG	-7.35	101.42	107.30
3	C	123	TRP	CE2-CD2-CG	-7.33	101.43	107.30
3	C	62	TRP	CE2-CD2-CG	-7.17	101.56	107.30
3	C	111	TRP	CD1-CG-CD2	7.13	112.01	106.30
2	B	106	TRP	CE2-CD2-CG	-7.02	101.68	107.30
2	B	47	TRP	CE2-CD2-CG	-6.87	101.80	107.30
3	C	108	TRP	CD1-CG-CD2	6.76	111.71	106.30
3	C	108	TRP	CE2-CD2-CG	-6.72	101.92	107.30
2	B	52	TRP	CE2-CD2-CG	-6.71	101.93	107.30
3	C	63	TRP	CE2-CD2-CG	-6.70	101.94	107.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	36	TRP	CE2-CD2-CG	-6.70	101.94	107.30
3	C	73	ARG	NE-CZ-NH1	6.67	123.64	120.30
1	A	35	TRP	CE2-CD2-CG	-6.63	102.00	107.30
3	C	68	ARG	NE-CZ-NH1	6.55	123.58	120.30
2	B	47	TRP	CG-CD2-CE3	6.23	139.51	133.90
3	C	108	TRP	CG-CD2-CE3	5.86	139.18	133.90
1	A	86	TYR	CB-CG-CD2	-5.83	117.50	121.00
3	C	108	TRP	CB-CG-CD1	-5.73	119.55	127.00
1	A	24	ARG	NE-CZ-NH1	5.71	123.16	120.30
2	B	36	TRP	CG-CD1-NE1	-5.51	104.59	110.10
2	B	47	TRP	CB-CG-CD1	-5.44	119.93	127.00
2	B	102	ARG	NE-CZ-NH2	-5.39	117.61	120.30
3	C	123	TRP	CB-CG-CD1	-5.37	120.02	127.00
2	B	59	TYR	CB-CG-CD1	-5.24	117.86	121.00
2	B	47	TRP	CG-CD1-NE1	-5.23	104.87	110.10
2	B	52	TRP	CG-CD2-CE3	5.20	138.58	133.90
3	C	5	ARG	NE-CZ-NH2	-5.17	117.71	120.30
3	C	112	ARG	NE-CZ-NH2	-5.15	117.72	120.30
2	B	66	ARG	NE-CZ-NH2	-5.13	117.73	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	826	194	798	2	0
2	B	904	219	872	6	0
3	C	1001	264	963	6	0
4	A	8	0	0	0	0
4	B	14	0	0	0	0
4	C	26	0	0	1	0
All	All	2779	677	2633	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:90:HIS:HD2	1:A:92:TRP:H	1.46	0.62
3:C:48:ASP:HB2	3:C:61:ARG:HH12	1.72	0.54
3:C:61:ARG:HH21	3:C:70:PRO:HG2	1.72	0.53
1:A:39:LYS:HE2	1:A:81:GLU:O	2.09	0.53
2:B:6:GLU:HA	2:B:21:THR:O	2.13	0.49
2:B:67:LEU:HD22	2:B:80:LEU:HD11	1.95	0.47
3:C:24:SER:HA	4:C:200:HOH:O	2.15	0.46
2:B:97:ARG:O	2:B:103:LEU:HA	2.18	0.43
2:B:87:THR:HA	2:B:114:VAL:HB	2.02	0.42
3:C:61:ARG:NH2	3:C:70:PRO:HG2	2.34	0.41
2:B:89:ASP:HB2	2:B:114:VAL:HG21	2.01	0.41
3:C:69:THR:HA	3:C:70:PRO:HD2	1.78	0.41
2:B:75:LYS:HE3	2:B:77:GLN:NE2	2.35	0.41
3:C:44:ASN:O	3:C:51:THR:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	105/107 (98%)	102 (97%)	3 (3%)	0	100	100
2	B	114/116 (98%)	112 (98%)	2 (2%)	0	100	100
3	C	127/129 (98%)	122 (96%)	4 (3%)	1 (1%)	19	7
All	All	346/352 (98%)	336 (97%)	9 (3%)	1 (0%)	41	27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	48	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	91/91 (100%)	89 (98%)	2 (2%)	52	39
2	B	100/100 (100%)	98 (98%)	2 (2%)	55	44
3	C	105/105 (100%)	103 (98%)	2 (2%)	57	46
All	All	296/296 (100%)	290 (98%)	6 (2%)	55	44

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	LEU
1	A	70	GLN
2	B	102	ARG
2	B	110	THR
3	C	48	ASP
3	C	75	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	90	HIS
2	B	77	GLN
3	C	37	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.