

Full wwPDB X-ray Structure Validation Report (i)

May 23, 2020 – 04:42 pm BST

PDB ID : 2VG3

Title: Rv2361 with citronellyl pyrophosphate Authors: Naismith, J.H.; Wang, W.; Dong, C.

Deposited on : 2007-11-08

Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.11

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	284	88%	11%
1	В	284	89%	9% •
1	С	284	93%	7%
1	D	284	95%	•



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 9701 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called UNDECAPRENYL PYROPHOSPHATE SYNTHETASE.


Mol	Chain	Residues	${f Atoms}$			ZeroOcc	AltConf	Trace		
1	Λ	284	Total	С	N	О	S	0	4	0
1	A	204	2307	1457	422	419	9	0	4	
1	В	284	Total	С	N	О	S	0	0	0
1	Б	204	2288	1443	422	414	9	0	U	
1	С	284	Total	С	N	О	S	0	1	0
1		204	2294	1447	422	416	9	0	1	
1	D	283	Total	С	N	О	S	0	0	0
1	ש	200	2277	1434	421	413	9	U	U	U

• Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0
2	С	1	Total Mg 1 1	0	0


• Molecule 3 is GERANYL DIPHOSPHATE (three-letter code: GPP) (formula: C₁₀H₂₀O₇P₂).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	Λ	1	Total	С	О	Р	0	0
)	A	1	19	10	7	2	0	0
3	D	1	Total	С	О	Р	0	0
)	Б	1	19	10	7	2	0	0
3	С	1	Total	С	О	Р	0	0
)		1	19	10	7	2	0	0
3	D	1	Total	С	О	Р	0	0
)	ש	1	19	10	7	2	0	U

 \bullet Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: $\mathrm{O_4P}).$






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O P 5 4 1	0	0
4	В	1	Total O P 5 4 1	0	0
4	В	1	Total O P 5 4 1	0	0

• Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	2	Total Cl 2 2	0	0


• Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
6	С	1	Total 5	O 4	S 1	0	0

 \bullet Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: $\mathrm{C_3H_8O_3}).$





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	С	1	Total C C 6 3 3)	0	0

• Molecule 8 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	139	Total O 139 139	0	0
8	В	106	Total O 106 106	0	0
8	С	99	Total O 99 99	0	0
8	С	1	Total O 1 1	0	0
8	D	82	Total O 82 82	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: UNDECAPRENYL PYROPHOSPHATE SYNTHETASE





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32	Depositor
Cell constants	86.87Å 86.87Å 184.07Å	Danasitan
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	75.16 - 1.80	Depositor
rtesoration (A)	69.64 - 1.80	EDS
% Data completeness	91.6 (75.16-1.80)	Depositor
(in resolution range)	87.1 (69.64-1.80)	EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	0.67 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.203 , 0.231	Depositor
	0.206 , 0.231	DCC
R_{free} test set	6269 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	19.2	Xtriage
Anisotropy	0.022	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	$0.35\;,28.8$	EDS
L-test for twinning ²	$< L > = 0.47, < L^2> = 0.30$	Xtriage
	0.003 for -h,-k,l	
Estimated twinning fraction	0.259 for h,-h-k,-l	Xtriage
	0.002 for -k,-h,-l	
F_o, F_c correlation	0.95	EDS
Total number of atoms	9701	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.44% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, CL, PO4, SO4, GPP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.53	0/2381	0.70	0/3238	
1	В	0.49	0/2350	0.67	3/3196~(0.1%)	
1	С	0.46	0/2359	0.63	0/3208	
1	D	0.42	0/2338	0.62	0/3179	
All	All	0.48	0/9428	0.66	3/12821~(0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	Α	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\mathbf{Ideal}(^{o})$
1	В	145	ARG	NE-CZ-NH2	-7.38	116.61	120.30
1	В	145	ARG	NE-CZ-NH1	6.81	123.71	120.30
1	В	49	ARG	NE-CZ-NH2	-6.12	117.24	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	44	TYR	Peptide



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2307	0	2271	21	0
1	В	2288	0	2245	23	0
1	С	2294	0	2251	11	0
1	D	2277	0	2237	7	0
2	A	1	0	0	0	0
2	В	1	0	0	0	0
2	С	1	0	0	0	0
2	D	1	0	0	0	0
3	A	19	0	17	0	0
3	В	19	0	17	0	0
3	С	19	0	17	0	0
3	D	19	0	17	4	0
4	A	5	0	0	0	0
4	В	10	0	0	0	0
5	В	2	0	0	1	0
6	С	5	0	0	0	0
7	С	6	0	8	0	0
8	A	139	0	0	0	0
8	В	106	0	0	0	0
8	С	100	0	0	0	0
8	D	82	0	0	0	0
All	All	9701	0	9080	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{array}{l} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{array}$	Clash overlap (Å)
1:B:255:MET:HE3	1:B:258:GLN:HB2	1.61	0.82
1:A:31:THR:HG21	1:A:35[B]:VAL:HG23	1.71	0.73
1:A:147:ARG:HG2	1:A:180:MET:SD	2.34	0.67
1:C:13:PHE:N	1:C:14:PRO:HA	2.10	0.67
1:D:139:ASN:HD21	3:D:1298:GPP:H91	1.60	0.66
1:B:255:MET:HE3	1:B:258:GLN:CB	2.27	0.65

Continued on next page...



Continued from previous page...

Continuea from previ	1 0	Interatomic	Clash
Atom-1	Atom-2	${f distance} \; ({f \AA})$	overlap $(ext{Å})$
1:B:70:HIS:CE1	1:B:255:MET:HE2	2.35	0.62
1:B:66:ARG:NH2	5:B:1301:CL:CL	2.70	0.61
1:B:70:HIS:HE1	1:B:255:MET:HE2	1.66	0.60
1:D:139:ASN:ND2	3:D:1298:GPP:H91	2.18	0.59
1:B:119:ALA:HB3	1:B:193:VAL:HG22	1.88	0.55
1:C:13:PHE:N	1:C:14:PRO:CA	2.69	0.55
1:A:118:TYR:OH	1:A:121:SER:HB2	2.07	0.55
1:A:44:TYR:HB2	1:A:45:GLY:CA	2.37	0.54
1:A:206:ARG:CZ	1:C:222:ILE:O	2.56	0.54
1:B:227:ILE:HD11	1:D:202:THR:HA	1.91	0.53
1:B:97:GLU:OE2	1:B:145:ARG:HD2	2.09	0.53
1:A:162[A]:SER:OG	1:A:164:PRO:HD2	2.10	0.52
1:A:44:TYR:HB2	1:A:45:GLY:C	2.30	0.52
1:A:31:THR:HG23	1:A:34:VAL:HA	1.93	0.51
1:D:139:ASN:HD21	3:D:1298:GPP:C9	2.24	0.50
1:B:43:PRO:O	1:B:44:TYR:CG	2.65	0.50
1:B:70:HIS:HE1	1:B:255:MET:CE	2.25	0.49
1:C:160:VAL:O	1:C:192:CYS:HA	2.12	0.49
1:A:143:VAL:O	1:A:147:ARG:HB2	2.13	0.48
1:C:159:TRP:HB2	1:C:174:LEU:HD13	1.95	0.48
1:C:160:VAL:HG21	1:C:190:ASN:HB3	1.96	0.48
1:D:138:PHE:O	1:D:142:VAL:HG23	2.14	0.48
1:B:43:PRO:O	1:B:44:TYR:CD2	2.67	0.47
1:D:119:ALA:O	3:D:1298:GPP:H93	2.14	0.47
1:B:223:THR:HG23	1:B:225:SER:H	1.78	0.47
1:B:223:THR:HG23	1:B:225:SER:N	2.30	0.47
1:A:13:PHE:HB3	1:A:15:GLN:OE1	2.15	0.46
1:A:206:ARG:NH1	1:C:222:ILE:O	2.49	0.46
1:A:169:SER:O	1:A:173:GLU:HG2	2.16	0.46
1:A:155:VAL:HG22	1:A:187:ILE:CG2	2.46	0.45
1:B:136:MET:SD	1:B:166:LEU:HD12	2.57	0.44
1:B:160:VAL:HG22	1:B:233:ARG:HG3	1.99	0.44
1:B:13:PHE:CD2	1:B:61:ARG:HD2	2.52	0.44
1:B:223:THR:CG2	1:B:225:SER:OG	2.67	0.43
1:A:208:ILE:CD1	1:A:227:ILE:HD13	2.49	0.43
1:A:44:TYR:CD2	1:A:45:GLY:HA2	2.53	0.43
1:C:25:THR:HA	1:C:52:GLN:O	2.19	0.42
1:A:159:TRP:HB2	1:A:174:LEU:HD13	2.01	0.42
1:B:43:PRO:C	1:B:44:TYR:CG	2.93	0.42
1:B:99:VAL:HG12	1:B:271:TRP:CZ2	2.55	0.42
1:C:81:TRP:CH2	1:C:95:MET:HG3	2.56	0.41

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	$egin{array}{c} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{array}$	Clash overlap (Å)
1:A:160:VAL:O	1:A:192:CYS:HA	2.21	0.41
1:A:69:ASN:HB3	1:A:240:ASP:OD2	2.20	0.41
1:B:32:TRP:HA	1:B:33:PRO:C	2.40	0.41
1:B:245:THR:HB	1:B:269:LYS:O	2.21	0.41
1:B:160:VAL:O	1:B:192:CYS:HA	2.20	0.41
1:C:243:LEU:CD2	1:C:265:ILE:HB	2.50	0.41
1:A:269:LYS:HD3	1:A:274:TYR:HA	2.03	0.41
1:A:106:GLY:HA3	1:A:276:ARG:HB2	2.02	0.41
1:A:44:TYR:CD1	1:A:44:TYR:N	2.89	0.41
1:C:67:LEU:HD11	1:C:110:LEU:HG	2.02	0.41
1:B:118:TYR:OH	1:B:121:SER:HB2	2.21	0.41
1:D:169:SER:O	1:D:173:GLU:HG2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	${f ntiles}$
1	A	$286/284 \; (101\%)$	279 (98%)	6 (2%)	1 (0%)	41	27
1	В	282/284 (99%)	274 (97%)	7 (2%)	1 (0%)	34	21
1	С	283/284 (100%)	279 (99%)	4 (1%)	0	100	100
1	D	281/284 (99%)	278 (99%)	3 (1%)	0	100	100
All	All	$1132/1136 \; (100\%)$	1110 (98%)	20 (2%)	2 (0%)	47	33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	44	TYR
1	A	46	GLY



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	$243/239 \ (102\%)$	240 (99%)	3 (1%)	71 65
1	В	239/239 (100%)	236 (99%)	3 (1%)	69 62
1	С	240/239 (100%)	236 (98%)	4 (2%)	60 51
1	D	238/239 (100%)	233 (98%)	5 (2%)	53 42
All	All	960/956 (100%)	945 (98%)	15 (2%)	62 54

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	TYR
1	A	118	TYR
1	A	158	ARG
1	В	66	ARG
1	В	118	TYR
1	В	158	ARG
1	С	22	ASP
1	С	118	TYR
1	С	145	ARG
1	С	158	ARG
1	D	31	THR
1	D	44	TYR
1	D	118	TYR
1	D	158	ARG
1	D	225	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	С	69	ASN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 15 ligands modelled in this entry, 6 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tune	Chain	Res	Link	Во	ond leng	ths	В	ond ang	les
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PO4	A	1299	-	4,4,4	1.31	1 (25%)	6,6,6	0.83	0
7	GOL	С	1300	-	5,5,5	0.35	0	5,5,5	0.17	0
4	PO4	В	1300	_	4,4,4	0.76	0	6,6,6	0.67	0
4	PO4	В	1299	-	4,4,4	0.80	0	6,6,6	1.39	1 (16%)
3	GPP	С	1298	2	16,18,18	2.72	4 (25%)	21,25,25	2.10	5 (23%)
3	GPP	D	1298	2	16,18,18	2.48	3 (18%)	21,25,25	2.76	6 (28%)
6	SO4	С	1299	-	4,4,4	0.11	0	6,6,6	0.22	0
3	GPP	A	1298	2	16,18,18	2.75	4 (25%)	21,25,25	2.43	6 (28%)
3	GPP	В	1298	2	16,18,18	2.78	3 (18%)	21,25,25	2.12	5 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	${ m Res}$	Link	Chirals	${f Torsions}$	Rings
3	GPP	D	1298	2	-	7/19/19/19	-
3	GPP	A	1298	2	-	7/19/19/19	-
3	GPP	В	1298	2	-	5/19/19/19	-
3	GPP	С	1298	2	-	6/19/19/19	-
7	GOL	С	1300	-	-	0/4/4/4	-

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(ext{\AA})$
3	В	1298	GPP	C2-C3	9.38	1.55	1.33
3	С	1298	GPP	C2-C3	9.16	1.54	1.33
3	A	1298	GPP	C2-C3	9.12	1.54	1.33
3	D	1298	GPP	C2-C3	8.29	1.52	1.33
3	С	1298	GPP	C6-C7	-3.53	1.38	1.50
3	D	1298	GPP	C6-C7	-3.28	1.39	1.50
3	В	1298	GPP	C6-C7	-3.26	1.39	1.50
3	A	1298	GPP	C6-C7	-3.26	1.39	1.50
3	В	1298	GPP	C7-C8	2.70	1.40	1.32
3	A	1298	GPP	C7-C8	2.68	1.40	1.32
3	D	1298	GPP	C7-C8	2.63	1.39	1.32
3	С	1298	GPP	C7-C8	2.50	1.39	1.32
4	A	1299	PO4	P-O2	-2.24	1.47	1.54
3	A	1298	GPP	PA-O1	-2.13	1.50	1.59
3	С	1298	GPP	PB-O3B	-2.00	1.47	1.54

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^o)$
3	D	1298	GPP	C1-C2-C3	-8.78	110.86	126.04
3	A	1298	GPP	C1-C2-C3	-7.49	113.09	126.04
3	В	1298	GPP	C1-C2-C3	-5.69	116.19	126.04
3	D	1298	GPP	C5-C3-C2	-5.67	109.64	121.12
3	С	1298	GPP	C1-C2-C3	-5.56	116.43	126.04
3	D	1298	GPP	C4-C3-C2	-5.06	110.70	123.68
3	A	1298	GPP	C5-C3-C2	-4.98	111.04	121.12
3	С	1298	GPP	C4-C3-C2	-4.70	111.63	123.68
3	В	1298	GPP	C4-C3-C2	-4.48	112.19	123.68
3	A	1298	GPP	C4-C3-C2	-4.26	112.76	123.68
3	В	1298	GPP	C5-C3-C2	-4.23	112.56	121.12
3	С	1298	GPP	C5-C3-C2	-4.04	112.94	121.12
3	С	1298	GPP	C10-C8-C9	2.77	120.72	114.60
3	D	1298	GPP	C4-C3-C5	-2.68	110.76	115.27

Continued on next page...



 $Continued\ from\ previous\ page...$

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
3	A	1298	GPP	O2B-PB-O3A	2.59	113.33	104.64
4	В	1299	PO4	O4-P-O3	2.53	116.10	107.97
3	В	1298	GPP	O2B-PB-O3A	2.31	112.38	104.64
3	D	1298	GPP	PA-O3A-PB	-2.31	124.91	132.83
3	D	1298	GPP	C10-C8-C9	2.27	119.62	114.60
3	A	1298	GPP	C10-C8-C9	2.19	119.45	114.60
3	В	1298	GPP	C10-C8-C9	2.18	119.43	114.60
3	A	1298	GPP	C4-C3-C5	-2.17	111.62	115.27
3	С	1298	GPP	C4-C3-C5	-2.12	111.70	115.27

There are no chirality outliers.

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	С	1298	GPP	C2-C1-O1-PA
3	С	1298	GPP	C1-O1-PA-O1A
3	С	1298	GPP	C1-C2-C3-C4
3	D	1298	GPP	C1-O1-PA-O1A
3	D	1298	GPP	C1-C2-C3-C4
3	D	1298	GPP	C1-C2-C3-C5
3	A	1298	GPP	C1-O1-PA-O1A
3	A	1298	GPP	C1-C2-C3-C4
3	A	1298	GPP	C1-C2-C3-C5
3	В	1298	GPP	C2-C1-O1-PA
3	В	1298	GPP	C1-C2-C3-C4
3	В	1298	GPP	C3-C5-C6-C7
3	D	1298	GPP	C3-C5-C6-C7
3	С	1298	GPP	C4-C3-C5-C6
3	D	1298	GPP	C4-C3-C5-C6
3	В	1298	GPP	C4-C3-C5-C6
3	D	1298	GPP	C2-C1-O1-PA
3	A	1298	GPP	C2-C1-O1-PA
3	A	1298	GPP	C4-C3-C5-C6
3	D	1298	GPP	C1-O1-PA-O3A
3	A	1298	GPP	C3-C5-C6-C7
3	С	1298	GPP	C3-C5-C6-C7
3	С	1298	GPP	C1-O1-PA-O3A
3	A	1298	GPP	C1-O1-PA-O3A
3	В	1298	GPP	C1-O1-PA-O1A


There are no ring outliers.

1 monomer is involved in 4 short contacts:





Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	1298	GPP	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	284/284 (100%)	-0.87	2 (0%) 87 86	8, 15, 23, 31	0
1	В	$284/284 \; (100\%)$	-0.84	2 (0%) 87 86	11, 15, 23, 33	0
1	С	$284/284 \ (100\%)$	-0.81	2 (0%) 87 86	8, 15, 23, 31	0
1	D	283/284 (99%)	-0.79	5 (1%) 68 64	5, 15, 23, 31	0
All	All	$1135/1136 \ (99\%)$	-0.83	11 (0%) 82 80	5, 15, 23, 33	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	44	TYR	4.3
1	D	45	GLY	3.8
1	D	43	PRO	3.6
1	С	44	TYR	3.4
1	A	44	TYR	3.2
1	С	43	PRO	3.0
1	В	44	TYR	2.8
1	D	46	GLY	2.3
1	D	42	ALA	2.1
1	В	43	PRO	2.1
1	A	45	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
7	GOL	С	1300	6/6	0.86	0.23	53,54,54,54	0
2	MG	D	1297	1/1	0.99	0.04	16,16,16,16	0
2	MG	В	1297	1/1	0.99	0.03	17,17,17,17	0
3	GPP	D	1298	19/19	0.99	0.07	15,20,35,35	0
3	GPP	A	1298	19/19	0.99	0.06	13,17,35,35	0
3	GPP	В	1298	19/19	0.99	0.05	14,20,37,38	0
2	MG	A	1297	1/1	1.00	0.03	16,16,16,16	0
4	PO4	В	1299	5/5	1.00	0.05	19,19,19,21	0
2	MG	С	1297	1/1	1.00	0.04	16,16,16,16	0
3	GPP	С	1298	19/19	1.00	0.06	12,16,32,32	0
5	CL	В	1302	1/1	1.00	0.02	17,17,17,17	0
6	SO4	С	1299	5/5	1.00	0.06	25,25,27,27	0
4	PO4	A	1299	5/5	1.00	0.06	20,20,23,23	0
4	PO4	В	1300	5/5	1.00	0.08	23,24,26,26	0
5	CL	В	1301	1/1	1.00	0.02	18,18,18,18	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around GPP D 1298: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive) Electron density around GPP A 1298:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











Electron density around GPP B 1298:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around GPP C 1298:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

