



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 2, 2023 – 11:22 PM EDT

PDB ID : 3VGD  
Title : Crystal structure of glycosyltrehalose trehalohydrolase (D252E)  
Authors : Okazaki, N.; Tamada, T.; Feese, M.D.; Kato, M.; Miura, Y.; Komeda, T.; Kobayashi, K.; Kondo, K.; Kuroki, R.  
Deposited on : 2011-08-09  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	558	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	1104	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4972 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Malto-oligosyltrehalose trehalohydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	553	4537	2925	744	859	9	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	252	GLU	ASP	engineered mutation	UNP Q55088

- Molecule 2 is CITRATE ANION (three-letter code: FLC) (formula:  $C_6H_5O_7$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	13	6	7	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	392	Total O 392 392	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Malto-oligosyltrehalose trehalohydrolase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.88Å 78.88Å 282.53Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	61.50 – 2.40 61.50 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (61.50-2.40) 99.9 (61.50-2.40)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.77 (at 2.40Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.158 , 0.198 0.156 , 0.193	Depositor DCC
$R_{free}$ test set	2050 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.6	Xtrriage
Anisotropy	0.108	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 46.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.028 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4972	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.71	0/4653	0.79	7/6286 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	493	ARG	NE-CZ-NH2	-7.79	116.41	120.30
1	A	278	ARG	NE-CZ-NH2	-7.37	116.61	120.30
1	A	493	ARG	NE-CZ-NH1	7.32	123.96	120.30
1	A	250	ARG	NE-CZ-NH2	-7.01	116.80	120.30
1	A	280	VAL	CB-CA-C	-6.55	98.95	111.40
1	A	365	ASP	CB-CG-OD1	5.46	123.21	118.30
1	A	278	ARG	NE-CZ-NH1	5.08	122.84	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4537	0	4393	87	0
2	A	13	0	5	1	0
3	A	30	0	40	14	0
4	A	392	0	0	13	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4972	0	4438	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1104:GOL:H2	4:A:2256:HOH:O	1.56	1.03
1:A:264:HIS:HD2	1:A:266:LEU:H	1.24	0.85
1:A:113:THR:HG22	1:A:116:GLY:H	1.44	0.83
1:A:493:ARG:HD2	4:A:2023:HOH:O	1.77	0.82
1:A:113:THR:HG21	1:A:120:GLY:HA3	1.61	0.81
1:A:311:HIS:HD2	1:A:312:HIS:ND1	1.81	0.78
1:A:143:GLN:HE22	1:A:162:ASN:H	1.31	0.77
1:A:133:ILE:HD11	1:A:413:ILE:HD12	1.73	0.71
1:A:191:ASN:OD1	1:A:192:HIS:HD2	1.75	0.70
1:A:336:VAL:HB	1:A:497:VAL:HG21	1.72	0.69
1:A:155:VAL:HG22	1:A:198:ASN:HB2	1.73	0.69
1:A:368:ASN:ND2	3:A:1104:GOL:C1	2.56	0.68
1:A:214:PRO:HD3	4:A:2390:HOH:O	1.96	0.64
1:A:357:HIS:HD2	4:A:2015:HOH:O	1.81	0.62
1:A:35:MET:HG2	1:A:45:ILE:HD11	1.81	0.62
1:A:480:LEU:HD13	1:A:532:LEU:HD23	1.81	0.61
3:A:1104:GOL:H11	4:A:2192:HOH:O	1.99	0.61
1:A:217:LEU:HD12	1:A:217:LEU:H	1.64	0.61
1:A:205:PRO:HG2	1:A:225:GLU:HG3	1.82	0.61
1:A:444:ARG:HE	1:A:450:GLN:NE2	1.99	0.60
1:A:155:VAL:CG2	1:A:198:ASN:HB2	2.31	0.60
1:A:255:HIS:CD2	1:A:285:ASP:H	2.19	0.60
1:A:357:HIS:HE1	4:A:2025:HOH:O	1.85	0.59
1:A:250:ARG:HD3	1:A:250:ARG:C	2.22	0.59
1:A:200:MET:HE3	1:A:207:PHE:CZ	2.37	0.58
1:A:100:GLU:HB2	1:A:491:CYS:SG	2.45	0.57
1:A:52:VAL:O	1:A:53:ARG:HB2	2.04	0.57
1:A:99:LYS:HZ3	3:A:1104:GOL:H12	1.70	0.57
1:A:368:ASN:HD21	3:A:1104:GOL:C1	2.16	0.57
1:A:92:ASN:H	1:A:245:ASN:HD21	1.54	0.56
1:A:427:PHE:HB3	1:A:450:GLN:HE22	1.71	0.56
1:A:72:ARG:HD3	1:A:239:TYR:CE1	2.41	0.56
1:A:264:HIS:CD2	1:A:266:LEU:H	2.15	0.56

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:152:TYR:O	1:A:192:HIS:HE1	1.88	0.55
3:A:1104:GOL:C1	4:A:2192:HOH:O	2.55	0.55
1:A:323:GLN:HE21	1:A:323:GLN:H	1.55	0.55
1:A:113:THR:HG23	1:A:115:GLU:H	1.71	0.55
1:A:273:VAL:HG21	1:A:280:VAL:HG22	1.89	0.55
1:A:20:GLN:OE1	1:A:60:LEU:HB3	2.06	0.55
1:A:273:VAL:CG2	1:A:280:VAL:HG22	2.37	0.54
1:A:20:GLN:HG2	1:A:23:VAL:HG22	1.89	0.54
1:A:368:ASN:HD21	3:A:1104:GOL:H11	1.73	0.54
1:A:92:ASN:H	1:A:245:ASN:ND2	2.06	0.54
1:A:368:ASN:ND2	3:A:1104:GOL:H12	2.24	0.53
1:A:113:THR:CG2	1:A:116:GLY:H	2.18	0.52
1:A:72:ARG:HD3	1:A:239:TYR:HE1	1.72	0.52
1:A:200:MET:CE	1:A:207:PHE:CZ	2.93	0.51
1:A:108:HIS:HE1	1:A:149:ASP:O	1.93	0.51
1:A:321:GLU:HB3	1:A:323:GLN:HE22	1.76	0.51
1:A:467:TRP:CB	3:A:1103:GOL:H12	2.41	0.50
1:A:99:LYS:NZ	3:A:1104:GOL:H12	2.26	0.50
1:A:467:TRP:HB3	3:A:1103:GOL:H12	1.94	0.50
2:A:1001:FLC:OA2	3:A:1101:GOL:H11	2.12	0.50
1:A:48:ASN:OD1	1:A:48:ASN:N	2.44	0.49
1:A:499:ASN:HB3	1:A:504:LEU:HD12	1.94	0.49
1:A:323:GLN:H	1:A:323:GLN:NE2	2.11	0.48
1:A:116:GLY:O	3:A:1105:GOL:O3	2.30	0.48
1:A:121:VAL:HG22	1:A:173:LEU:CD1	2.43	0.48
1:A:296:GLU:H	1:A:296:GLU:CD	2.17	0.48
1:A:444:ARG:HE	1:A:450:GLN:HE21	1.59	0.48
1:A:357:HIS:CD2	4:A:2015:HOH:O	2.62	0.47
1:A:264:HIS:HE1	4:A:2210:HOH:O	1.98	0.46
1:A:179:LYS:HD2	4:A:2089:HOH:O	2.16	0.46
1:A:529:THR:HB	1:A:541:HIS:HD2	1.81	0.46
1:A:311:HIS:HE1	4:A:2040:HOH:O	1.99	0.46
1:A:191:ASN:OD1	1:A:192:HIS:CD2	2.62	0.46
1:A:250:ARG:C	1:A:250:ARG:CD	2.84	0.46
1:A:529:THR:HB	1:A:541:HIS:CD2	2.51	0.45
1:A:42:TYR:OH	1:A:228:GLU:HB3	2.16	0.44
1:A:544:GLU:H	1:A:544:GLU:HG2	1.37	0.44
1:A:121:VAL:HG22	1:A:173:LEU:HD13	1.99	0.44
1:A:499:ASN:ND2	4:A:2349:HOH:O	2.50	0.44
1:A:105:TYR:HB2	1:A:133:ILE:HG13	1.98	0.44
3:A:1104:GOL:C2	4:A:2256:HOH:O	2.37	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:103:ILE:HG22	1:A:133:ILE:HD12	2.00	0.43
1:A:200:MET:CE	1:A:207:PHE:HZ	2.31	0.43
1:A:255:HIS:CD2	1:A:255:HIS:H	2.35	0.43
1:A:336:VAL:HG12	1:A:504:LEU:HD11	2.00	0.43
1:A:345:TYR:CD2	1:A:348:LYS:HG2	2.54	0.43
1:A:113:THR:HG22	1:A:116:GLY:N	2.23	0.43
1:A:416:GLY:HA2	1:A:419:TYR:CE2	2.54	0.43
1:A:226:SER:HA	1:A:229:VAL:HG13	2.01	0.42
1:A:396:SER:OG	1:A:551:LYS:NZ	2.52	0.42
1:A:213:THR:HB	1:A:214:PRO:HD2	2.01	0.42
1:A:25:LEU:HB2	1:A:45:ILE:HD11	2.01	0.42
1:A:127:TYR:CD2	1:A:128:LEU:HD13	2.55	0.41
1:A:480:LEU:HD13	1:A:532:LEU:CD2	2.47	0.41
1:A:311:HIS:CD2	1:A:312:HIS:ND1	2.74	0.41
1:A:35:MET:HG2	1:A:45:ILE:CG1	2.51	0.41
1:A:198:ASN:HD21	1:A:200:MET:HB2	1.86	0.40
1:A:246:VAL:O	1:A:278:ARG:HD2	2.22	0.40
1:A:315:HIS:CD2	1:A:326:TYR:CZ	3.10	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	549/558 (98%)	533 (97%)	16 (3%)	0	<a href="#">100</a> <a href="#">100</a>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	489/494 (99%)	438 (90%)	51 (10%)	7 10

All (51) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LYS
1	A	9	ASN
1	A	11	VAL
1	A	12	ILE
1	A	15	LEU
1	A	21	LYS
1	A	25	LEU
1	A	28	LEU
1	A	39	GLU
1	A	45	ILE
1	A	47	LEU
1	A	59	VAL
1	A	76	GLU
1	A	87	GLU
1	A	93	ASN
1	A	94	GLU
1	A	113	THR
1	A	121	VAL
1	A	125	LEU
1	A	128	LEU
1	A	133	ILE
1	A	155	VAL
1	A	168	GLU
1	A	179	LYS
1	A	193	VAL
1	A	229	VAL
1	A	246	VAL
1	A	250	ARG
1	A	278	ARG
1	A	280	VAL
1	A	296	GLU
1	A	297	LYS
1	A	323	GLN
1	A	332	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	339	TYR
1	A	348	LYS
1	A	361	VAL
1	A	372	TYR
1	A	438	GLN
1	A	480	LEU
1	A	487	LEU
1	A	499	ASN
1	A	514	LEU
1	A	518	SER
1	A	525	LYS
1	A	529	THR
1	A	531	LEU
1	A	536	ASN
1	A	544	GLU
1	A	553	PHE
1	A	555	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	ASN
1	A	74	GLN
1	A	108	HIS
1	A	143	GLN
1	A	192	HIS
1	A	245	ASN
1	A	255	HIS
1	A	264	HIS
1	A	301	ASN
1	A	311	HIS
1	A	323	GLN
1	A	331	ASN
1	A	357	HIS
1	A	368	ASN
1	A	450	GLN
1	A	541	HIS

### 5.3.3 RNA

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GOL	A	1103	-	5,5,5	0.48	0	5,5,5	1.26	1 (20%)
3	GOL	A	1102	-	5,5,5	0.32	0	5,5,5	0.36	0
3	GOL	A	1101	-	5,5,5	0.40	0	5,5,5	0.52	0
3	GOL	A	1105	-	5,5,5	0.30	0	5,5,5	0.77	0
3	GOL	A	1104	-	5,5,5	0.48	0	5,5,5	0.54	0
2	FLC	A	1001	-	12,12,12	1.24	0	17,17,17	1.30	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1103	-	-	2/4/4/4	-
3	GOL	A	1102	-	-	2/4/4/4	-
3	GOL	A	1101	-	-	1/4/4/4	-
3	GOL	A	1105	-	-	2/4/4/4	-
3	GOL	A	1104	-	-	4/4/4/4	-
2	FLC	A	1001	-	-	0/16/16/16	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	FLC	OB2-CBC-CB	3.54	119.20	113.05
2	A	1001	FLC	OB1-CBC-CB	-3.16	117.78	122.25
3	A	1103	GOL	O2-C2-C1	2.02	118.03	109.12

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1102	GOL	O1-C1-C2-C3
3	A	1104	GOL	O1-C1-C2-C3
3	A	1103	GOL	O1-C1-C2-C3
3	A	1104	GOL	C1-C2-C3-O3
3	A	1105	GOL	O1-C1-C2-C3
3	A	1104	GOL	O1-C1-C2-O2
3	A	1105	GOL	O1-C1-C2-O2
3	A	1103	GOL	O1-C1-C2-O2
3	A	1101	GOL	O1-C1-C2-C3
3	A	1102	GOL	O1-C1-C2-O2
3	A	1104	GOL	O2-C2-C3-O3

There are no ring outliers.

5 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1103	GOL	2	0
3	A	1101	GOL	1	0
3	A	1105	GOL	1	0
3	A	1104	GOL	10	0
2	A	1001	FLC	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	553/558 (99%)	-0.22	28 (5%) <span style="border: 1px solid red; padding: 2px;">28</span> <span style="border: 1px solid red; padding: 2px;">26</span>	20, 31, 67, 90	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	9	ASN	4.4
1	A	6	ILE	4.2
1	A	46	THR	4.2
1	A	8	GLY	3.7
1	A	33	TYR	3.6
1	A	3	ALA	3.6
1	A	435	LYS	3.5
1	A	5	LYS	3.2
1	A	45	ILE	3.1
1	A	31	GLY	3.0
1	A	87	GLU	3.0
1	A	11	VAL	2.8
1	A	50	VAL	2.8
1	A	7	ASP	2.7
1	A	36	GLU	2.7
1	A	32	LEU	2.7
1	A	12	ILE	2.6
1	A	63	ALA	2.6
1	A	39	GLU	2.6
1	A	21	LYS	2.5
1	A	23	VAL	2.3
1	A	51	LYS	2.3
1	A	10	GLU	2.2
1	A	47	LEU	2.2
1	A	61	ASP	2.2
1	A	22	SER	2.2
1	A	438	GLN	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	24	LYS	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	A	1103	6/6	0.83	0.20	44,48,49,51	0
3	GOL	A	1101	6/6	0.86	0.20	46,52,53,54	0
3	GOL	A	1102	6/6	0.91	0.24	55,60,60,61	0
3	GOL	A	1105	6/6	0.91	0.22	59,62,63,64	0
3	GOL	A	1104	6/6	0.93	0.15	40,45,48,51	0
2	FLC	A	1001	13/13	0.96	0.11	35,37,39,40	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.