

Full wwPDB X-ray Structure Validation Report (i)

Oct 5, 2023 – 07:51 AM EDT

PDB ID : 6VLE

Title : Crystal structure of human alpha 1,6-fucosyltransferase, FUT8 in its Apo-form Authors : Jarva, M.A.; Dramicanin, M.; Lingford, J.P.; Mao, R.; John, A.; Goddard-

Borger, E.D.

Deposited on : 2020-01-23

Resolution : 2.28 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity : FAILED

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : FAILED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.28 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 7873 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Alpha-(1,6)-fucosyltransferase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Δ	460	Total	С	N	О	S	0	0	0
1	Λ		3752	2390	661	687	14	U		
1	P	458	Total	С	N	О	S	0	0	0
1	D	498	3740	2383	659	684	14			U

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference	
A	576	HIS	-	expression tag	UNP Q9BYC5	
A	577	HIS	-	expression tag	UNP Q9BYC5	
A	578	HIS	-	expression tag	UNP Q9BYC5	
A	579	HIS	-	expression tag	UNP Q9BYC5	
A	580	HIS	-	expression tag	UNP Q9BYC5	
A	581	HIS	-	expression tag	UNP Q9BYC5	
A	582	HIS	-	expression tag	UNP Q9BYC5	
A	583	HIS	-	expression tag	UNP Q9BYC5	
A	584	HIS	-	expression tag	UNP Q9BYC5	
A	585	HIS	-	expression tag	UNP Q9BYC5	
В	576	HIS	-	expression tag	UNP Q9BYC5	
В	577	HIS	-	expression tag	UNP Q9BYC5	
В	578	HIS	-	expression tag	UNP Q9BYC5	
В	579	HIS	-	expression tag	UNP Q9BYC5	
В	580	HIS	-	expression tag	UNP Q9BYC5	
В	581	HIS	_	expression tag	UNP Q9BYC5	
В	582	HIS	-	expression tag	UNP Q9BYC5	
В	583	HIS	-	expression tag	UNP Q9BYC5	
В	584	HIS	-	expression tag	UNP Q9BYC5	
В	585	HIS	-	expression tag	UNP Q9BYC5	

• Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).





Mol	Chain	Residues	Atoms	;	ZeroOcc	AltConf
2	A	1	Total O 5 4	S 1	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	188	Total O 188 188	0	0
3	В	188	Total O 188 188	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source	
Space group	P 1 21 1	Depositor	
Cell constants	95.02Å 62.20Å 109.91Å	Depositor	
a, b, c, α , β , γ	90.00° 90.88° 90.00°	Depositor	
Resolution (Å)	47.25 - 2.28	Depositor	
% Data completeness	99.7 (47.25-2.28)	Depositor	
(in resolution range)	, ,		
R_{merge}	0.14	Depositor	
R_{sym}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	1.49 (at 2.29Å)	Xtriage	
Refinement program	PHENIX 1.16_3549	Depositor	
R, R_{free}	0.202 , 0.234	Depositor	
Wilson B-factor $(Å^2)$	41.1	Xtriage	
Anisotropy	0.732	Xtriage	
L-test for twinning ²	$< L > = 0.50, < L^2> = 0.33$	Xtriage	
Estimated twinning fraction	0.017 for h,-k,-l	Xtriage	
Total number of atoms	7873	wwPDB-VP	
Average B, all atoms (\mathring{A}^2)	50.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.01% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol		Type	Chain	Res	Link	Bond lengths			Bond angles		
	MIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
	2	SO4	A	601	-	4,4,4	0.15	0	6,6,6	0.06	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

