



# wwPDB Geometry-Only Validation Summary Report ⓘ

Feb 5, 2024 – 08:54 AM EST

PDB ID : 1VTM  
Title : STRUCTURE OF THE U2 STRAIN OF TOBACCO MOSAIC VIRUS  
REFINED AT 3.5 ANGSTROMS RESOLUTION USING X-RAY FIBER  
DIFFRACTION  
Authors : Stubbs, G.; Pattanayek, R.  
Deposited on : 1992-03-30  
Resolution : 3.50 Å(reported)

This is a wwPDB Geometry-Only Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*FIBER DIFFRACTION*

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)
RNA backbone	3102	1002 (4.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	R	3	100%
2	P	158	42% 40% 15% .

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1307 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (5'-R(P\*GP\*AP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	R	3	67	30	15	19	3	0	0	0

- Molecule 2 is a protein called Coat protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	P	158	1232	774	211	244	3	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
P	22	GLU	GLN	conflict	UNP P03579
P	55	SER	VAL	conflict	UNP P03579
P	57	VAL	SER	conflict	UNP P03579
P	97	ASN	ASP	conflict	UNP P03579

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	R	1	Total	O	0	0
			1	1		
3	P	7	Total	O	0	0
			7	7		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: RNA (5'-R(P\*GP\*AP\*A)-3')


Chain R:  100%



- Molecule 2: Coat protein

Chain P:  42% 40% 15%







## 4 Model quality i

### 4.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	R	2.66	2/75 (2.7%)	3.68	17/115 (14.8%)
2	P	0.87	0/1259	1.92	31/1726 (1.8%)
All	All	1.06	2/1334 (0.1%)	2.07	48/1841 (2.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	P	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	R	3	A	N9-C4	9.54	1.43	1.37
1	R	1	G	P-O5'	7.45	1.67	1.59

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	R	2	A	O4'-C1'-N9	14.47	119.77	108.20
1	R	1	G	P-O3'-C3'	10.98	132.88	119.70
2	P	106	GLU	CA-C-N	-10.36	94.42	117.20
1	R	2	A	C4'-C3'-C2'	10.27	112.86	102.60
1	R	2	A	C5'-C4'-C3'	-9.82	100.28	116.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	P	99	GLN	Peptide

## 4.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	R	67	0	34	4	0
2	P	1232	0	1194	67	0
3	P	7	0	0	0	0
3	R	1	0	0	0	0
All	All	1307	0	1228	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 67 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:P:43:THR:HA	2:P:46:GLN:HB2	1.51	0.90
2:P:119:VAL:HG13	2:P:122:ARG:NH1	1.93	0.84
2:P:24:ILE:HD11	2:P:132:LEU:HD22	1.60	0.83
2:P:96:VAL:HG23	2:P:98:ASN:HB2	1.61	0.81
2:P:119:VAL:HG13	2:P:122:ARG:HH11	1.46	0.77

There are no symmetry-related clashes.

## 4.3 Torsion angles [i](#)

### 4.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	P	156/158 (99%)	113 (72%)	31 (20%)	12 (8%)	<b>1</b> <b>10</b>

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	P	55	SER
2	P	96	VAL
2	P	99	GLN
2	P	136	THR
2	P	20	PRO

#### 4.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	P	136/136 (100%)	116 (85%)	20 (15%)	<b>3</b> <b>18</b>

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	P	128	LEU
2	P	141	GLN
2	P	154	THR
2	P	150	LEU
2	P	57	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
2	P	33	ASN
2	P	47	GLN
2	P	91	ASN
2	P	103	ASN
2	P	130	ASN

#### 4.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	R	2/3 (66%)	2 (100%)	1 (50%)

All (2) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	R	2	A
1	R	3	A

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	R	2	A

#### 4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

#### 4.6 Ligand geometry [i](#)

There are no ligands in this entry.

#### 4.7 Other polymers [i](#)

There are no such residues in this entry.

#### 4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.