



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 25, 2023 – 12:20 AM EDT

PDB ID : 5VTW
Title : Crystal structure of the A/Hong Kong/1/1968 (H3N2) influenza virus hemagglutinin G225M/L226T/S228A mutant in complex with 6'-SLN
Authors : Wu, N.C.; Wilson, I.A.
Deposited on : 2017-05-18
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.












Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	 3% 94% 5%
1	C	323	 2% 94%
1	E	323	 2% 94%
2	B	174	 14% 97%
2	D	174	 11% 95%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	174	 8% 97% ..
3	G	3	 100%
3	H	3	 100%
3	J	3	 67% 33%
4	I	2	 100%
4	K	2	 100%
5	L	2	 50% 50%
5	N	2	 50% 50%
6	M	4	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	NAG	F	201	-	-	-	X

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 12410 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin HA1 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	321	Total 2481	C 1553	N 438	O 476	S 14	0	0	0
1	C	317	Total 2445	C 1532	N 429	O 470	S 14	0	0	0
1	E	317	Total 2445	C 1532	N 429	O 470	S 14	0	0	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	ALA	-	expression tag	UNP Q91MA7
A	8	ASP	-	expression tag	UNP Q91MA7
A	9	PRO	-	expression tag	UNP Q91MA7
A	10	GLY	-	expression tag	UNP Q91MA7
A	225	MET	GLY	engineered mutation	UNP Q91MA7
A	226	THR	LEU	engineered mutation	UNP Q91MA7
A	228	ALA	SER	engineered mutation	UNP Q91MA7
C	7	ALA	-	expression tag	UNP Q91MA7
C	8	ASP	-	expression tag	UNP Q91MA7
C	9	PRO	-	expression tag	UNP Q91MA7
C	10	GLY	-	expression tag	UNP Q91MA7
C	225	MET	GLY	engineered mutation	UNP Q91MA7
C	226	THR	LEU	engineered mutation	UNP Q91MA7
C	228	ALA	SER	engineered mutation	UNP Q91MA7
E	7	ALA	-	expression tag	UNP Q91MA7
E	8	ASP	-	expression tag	UNP Q91MA7
E	9	PRO	-	expression tag	UNP Q91MA7
E	10	GLY	-	expression tag	UNP Q91MA7
E	225	MET	GLY	engineered mutation	UNP Q91MA7
E	226	THR	LEU	engineered mutation	UNP Q91MA7
E	228	ALA	SER	engineered mutation	UNP Q91MA7


- Molecule 2 is a protein called Hemagglutinin HA2 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	172	1391	863	243	279	6	0	0	0
2	D	171	1382	858	241	277	6	0	0	0
2	F	171	1382	858	241	277	6	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	123	GLY	ARG	conflict	UNP Q91MA7
D	123	GLY	ARG	conflict	UNP Q91MA7
F	123	GLY	ARG	conflict	UNP Q91MA7

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	G	3	39	22	2	15	0	0	0
3	H	3	39	22	2	15	0	0	0
3	J	3	39	22	2	15	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	I	2	28	16	2	10	0	0	0
4	K	2	28	16	2	10	0	0	0

- Molecule 5 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	L	2	32	17	1	14	0	0	0
5	N	2	32	17	1	14	0	0	0

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranos e-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluco pyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
6	M	4	50	28	2	20	0	0	0

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			14	8	1	5		
7	C	1	Total	C	N	O	0	0
			14	8	1	5		
7	C	1	Total	C	N	O	0	0
			14	8	1	5		
7	E	1	Total	C	N	O	0	0
			14	8	1	5		
7	E	1	Total	C	N	O	0	0
			14	8	1	5		
7	F	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 8 is N-acetyl-alpha-neuraminic acid (three-letter code: SIA) (formula: C₁₁H₁₉NO₉).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	O	0	0
			21	11	1	9		

- Molecule 9 is TRIS(HYDROXYETHYL)AMINOMETHANE (three-letter code: TAM) (formula: C₇H₁₇NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
9	E	1	11	7	1	3	0	0


- Molecule 10 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	102	Total	O	0	0
			102	102		
10	B	49	Total	O	0	0
			49	49		
10	C	85	Total	O	0	0
			85	85		
10	D	46	Total	O	0	0
			46	46		
10	E	146	Total	O	0	0
			146	146		
10	F	53	Total	O	0	0
			53	53		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hemagglutinin HA1 chain

Chain A: 



- Molecule 1: Hemagglutinin HA1 chain

Chain C: 




- Molecule 1: Hemagglutinin HA1 chain

Chain E: 





- Molecule 2: Hemagglutinin HA2 chain

Chain B: 




- Molecule 2: Hemagglutinin HA2 chain


Chain D: 




- Molecule 2: Hemagglutinin HA2 chain




- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose



- Molecule 5: N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose

Chain N:  50% 50%

GAL1
STA2

- Molecule 6: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M:  100%

MAG1
MAG2
BGA3
MAN4

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	208.65Å 131.53Å 72.73Å 90.00° 98.05° 90.00°	Depositor
Resolution (Å)	110.95 – 2.35 45.06 – 2.35	Depositor EDS
% Data completeness (in resolution range)	98.5 (110.95-2.35) 98.6 (45.06-2.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 2.34Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.193 , 0.213 0.196 , 0.215	Depositor DCC
R_{free} test set	3928 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	36.3	Xtrriage
Anisotropy	0.356	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 37.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12410	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, GAL, TAM, MAN, NAG, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/2538	0.70	1/3457 (0.0%)
1	C	0.48	0/2502	0.71	0/3410
1	E	0.51	0/2502	0.70	1/3410 (0.0%)
2	B	0.48	0/1415	0.67	0/1902
2	D	0.48	0/1406	0.69	2/1890 (0.1%)
2	F	0.46	0/1406	0.67	0/1890
All	All	0.49	0/11769	0.70	4/15959 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	201	ARG	NE-CZ-NH2	-5.28	117.66	120.30
2	D	38	LEU	CB-CG-CD2	-5.26	102.05	111.00
2	D	38	LEU	CB-CG-CD1	5.24	119.90	111.00
1	E	31	ASP	CB-CG-OD1	5.08	122.87	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2481	0	2433	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2445	0	2392	7	0
1	E	2445	0	2393	7	0
2	B	1391	0	1307	2	0
2	D	1382	0	1299	0	0
2	F	1382	0	1298	0	0
3	G	39	0	34	0	0
3	H	39	0	34	0	0
3	J	39	0	34	1	0
4	I	28	0	25	0	0
4	K	28	0	25	0	0
5	L	32	0	28	0	0
5	N	32	0	28	0	0
6	M	50	0	43	0	0
7	A	14	0	13	0	0
7	C	28	0	26	0	0
7	E	28	0	26	0	0
7	F	14	0	13	0	0
8	A	21	0	18	0	0
9	E	11	0	17	0	0
10	A	102	0	0	0	0
10	B	49	0	0	0	0
10	C	85	0	0	0	0
10	D	46	0	0	0	0
10	E	146	0	0	1	0
10	F	53	0	0	0	0
All	All	12410	0	11486	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:212:THR:HG21	1:E:216:ASN:HB3	1.44	0.99
2:B:72:GLU:OE2	1:E:238:LYS:HE2	1.76	0.84
1:C:212:THR:CG2	1:E:216:ASN:HB3	2.22	0.64
1:C:212:THR:HG21	1:E:216:ASN:CB	2.26	0.63
1:A:222:TRP:CZ2	1:A:225:MET:HA	2.41	0.54
1:A:53:ASN:HD22	1:A:54:ASN:N	2.05	0.54
1:E:203:THR:HG23	1:E:212:THR:HG22	1.90	0.53
1:C:53:ASN:HD21	1:C:276:THR:HG22	1.72	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53:ASN:HD22	1:A:53:ASN:C	2.17	0.48
1:A:207:ARG:HG2	1:C:221:PRO:HB2	1.99	0.45
1:E:33:GLN:NE2	10:E:604:HOH:O	2.49	0.45
1:E:201:ARG:HG2	1:E:248:ASN:OD1	2.17	0.44
1:C:167:THR:OG1	3:J:1:NAG:H62	2.20	0.41
2:B:50:GLY:HA3	1:C:30:THR:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	319/323 (99%)	309 (97%)	9 (3%)	1 (0%)	41	47
1	C	315/323 (98%)	305 (97%)	9 (3%)	1 (0%)	41	47
1	E	315/323 (98%)	306 (97%)	8 (2%)	1 (0%)	41	47
2	B	170/174 (98%)	163 (96%)	7 (4%)	0	100	100
2	D	169/174 (97%)	162 (96%)	7 (4%)	0	100	100
2	F	169/174 (97%)	162 (96%)	7 (4%)	0	100	100
All	All	1457/1491 (98%)	1407 (97%)	47 (3%)	3 (0%)	47	56

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	62	ILE
1	C	62	ILE
1	E	62	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	282/283 (100%)	272 (96%)	10 (4%)	36	44
1	C	278/283 (98%)	270 (97%)	8 (3%)	42	52
1	E	278/283 (98%)	273 (98%)	5 (2%)	59	70
2	B	146/148 (99%)	144 (99%)	2 (1%)	67	78
2	D	145/148 (98%)	141 (97%)	4 (3%)	43	53
2	F	145/148 (98%)	143 (99%)	2 (1%)	67	78
All	All	1274/1293 (98%)	1243 (98%)	31 (2%)	49	59

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	HIS
1	A	53	ASN
1	A	141	ARG
1	A	189	GLN
1	A	208	ARG
1	A	210	GLN
1	A	307	LYS
1	A	325	GLU
1	A	326	LYS
1	A	329	ARG
2	B	38	LEU
2	B	172	GLN
1	C	18	HIS
1	C	22	ASN
1	C	31	ASP
1	C	141	ARG
1	C	167	THR
1	C	208	ARG
1	C	210	GLN
1	C	216	ASN
2	D	39	LYS
2	D	51	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	59	THR
2	D	60	ASN
1	E	18	HIS
1	E	22	ASN
1	E	32	ASP
1	E	189	GLN
1	E	216	ASN
2	F	38	LEU
2	F	60	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	ASN
2	D	53	ASN
1	E	296	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

21 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	G	1	3,1	14,14,15	0.32	0	17,19,21	1.16	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	G	2	3	14,14,15	0.58	0	17,19,21	1.33	3 (17%)
3	BMA	G	3	3	11,11,12	0.44	0	15,15,17	1.12	2 (13%)
3	NAG	H	1	3,1	14,14,15	0.36	0	17,19,21	1.34	2 (11%)
3	NAG	H	2	3	14,14,15	0.42	0	17,19,21	0.74	1 (5%)
3	BMA	H	3	3	11,11,12	0.52	0	15,15,17	1.19	2 (13%)
4	NAG	I	1	4,1	14,14,15	0.54	0	17,19,21	1.22	3 (17%)
4	NAG	I	2	4	14,14,15	0.42	0	17,19,21	1.69	2 (11%)
3	NAG	J	1	3,1	14,14,15	0.43	0	17,19,21	1.27	1 (5%)
3	NAG	J	2	3	14,14,15	0.41	0	17,19,21	1.14	2 (11%)
3	BMA	J	3	3	11,11,12	0.92	0	15,15,17	2.41	4 (26%)
4	NAG	K	1	4,1	14,14,15	0.46	0	17,19,21	1.02	1 (5%)
4	NAG	K	2	4	14,14,15	0.42	0	17,19,21	1.49	2 (11%)
5	GAL	L	1	5	12,12,12	0.47	0	17,17,17	0.70	0
5	SIA	L	2	5	20,20,21	0.73	0	24,28,31	1.06	2 (8%)
6	NAG	M	1	6,1	14,14,15	0.27	0	17,19,21	1.13	1 (5%)
6	NAG	M	2	6	14,14,15	0.42	0	17,19,21	1.04	1 (5%)
6	BMA	M	3	6	11,11,12	0.49	0	15,15,17	1.06	2 (13%)
6	MAN	M	4	6	11,11,12	0.52	0	15,15,17	1.31	3 (20%)
5	GAL	N	1	5	12,12,12	0.51	0	17,17,17	0.75	0
5	SIA	N	2	5	20,20,21	0.69	0	24,28,31	1.25	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	G	1	3,1	-	1/6/23/26	0/1/1/1
3	NAG	G	2	3	-	2/6/23/26	0/1/1/1
3	BMA	G	3	3	-	0/2/19/22	0/1/1/1
3	NAG	H	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	H	2	3	-	0/6/23/26	0/1/1/1
3	BMA	H	3	3	-	2/2/19/22	0/1/1/1
4	NAG	I	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	I	2	4	-	1/6/23/26	0/1/1/1
3	NAG	J	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	J	2	3	-	2/6/23/26	0/1/1/1
3	BMA	J	3	3	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	K	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	K	2	4	-	2/6/23/26	0/1/1/1
5	GAL	L	1	5	-	0/2/22/22	0/1/1/1
5	SIA	L	2	5	-	0/18/34/38	0/1/1/1
6	NAG	M	1	6,1	-	0/6/23/26	0/1/1/1
6	NAG	M	2	6	-	0/6/23/26	0/1/1/1
6	BMA	M	3	6	-	2/2/19/22	0/1/1/1
6	MAN	M	4	6	-	1/2/19/22	0/1/1/1
5	GAL	N	1	5	-	0/2/22/22	0/1/1/1
5	SIA	N	2	5	-	0/18/34/38	0/1/1/1

There are no bond length outliers.

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	I	2	NAG	C1-O5-C5	5.86	120.13	112.19
3	J	3	BMA	C1-C2-C3	5.79	116.78	109.67
4	K	2	NAG	C1-O5-C5	4.62	118.45	112.19
3	J	3	BMA	C1-O5-C5	4.37	118.11	112.19
3	J	1	NAG	C1-O5-C5	4.33	118.06	112.19
3	J	3	BMA	O5-C1-C2	4.19	117.24	110.77
3	H	1	NAG	C1-O5-C5	4.02	117.63	112.19
3	G	1	NAG	C1-C2-N2	-3.65	104.26	110.49
3	G	2	NAG	C4-C3-C2	3.40	116.00	111.02
6	M	2	NAG	C1-O5-C5	3.09	116.37	112.19
6	M	4	MAN	C1-O5-C5	2.92	116.14	112.19
6	M	1	NAG	C1-O5-C5	2.84	116.04	112.19
3	H	3	BMA	C1-C2-C3	2.80	113.11	109.67
5	N	2	SIA	O8-C8-C7	2.62	115.48	109.10
5	L	2	SIA	O6-C2-C1	2.59	112.78	107.70
3	H	3	BMA	C1-O5-C5	2.56	115.65	112.19
3	G	3	BMA	C1-C2-C3	2.55	112.80	109.67
6	M	3	BMA	C1-O5-C5	2.54	115.63	112.19
3	J	2	NAG	C1-C2-N2	2.43	114.65	110.49
3	G	3	BMA	O5-C5-C6	2.35	110.89	107.20
3	H	2	NAG	C1-O5-C5	2.33	115.36	112.19
5	L	2	SIA	C6-O6-C2	2.33	116.32	111.34
5	N	2	SIA	O6-C2-C1	2.32	112.25	107.70
6	M	3	BMA	C3-C4-C5	2.32	114.37	110.24
5	N	2	SIA	O1B-C1-C2	2.29	119.58	113.03
3	J	3	BMA	C2-C3-C4	2.29	114.86	110.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	2	NAG	C1-O5-C5	2.25	115.23	112.19
5	N	2	SIA	C6-O6-C2	2.21	116.07	111.34
4	I	2	NAG	C4-C3-C2	-2.18	107.83	111.02
6	M	4	MAN	C3-C4-C5	2.17	114.10	110.24
6	M	4	MAN	C2-C3-C4	2.15	114.61	110.89
4	I	1	NAG	C6-C5-C4	2.14	118.01	113.00
4	K	1	NAG	C1-C2-N2	2.12	114.11	110.49
4	I	1	NAG	O4-C4-C5	2.11	114.54	109.30
4	I	1	NAG	C3-C4-C5	-2.09	106.51	110.24
4	K	2	NAG	C4-C3-C2	-2.06	108.00	111.02
3	H	1	NAG	O5-C1-C2	-2.04	108.06	111.29
3	G	1	NAG	C1-O5-C5	2.03	114.95	112.19
3	G	2	NAG	C3-C4-C5	2.01	113.82	110.24
3	J	2	NAG	O5-C5-C6	2.00	110.34	107.20

There are no chirality outliers.

All (13) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	J	2	NAG	O5-C5-C6-O6
3	G	2	NAG	C4-C5-C6-O6
3	J	2	NAG	C4-C5-C6-O6
4	K	2	NAG	C4-C5-C6-O6
3	G	2	NAG	O5-C5-C6-O6
3	H	3	BMA	C4-C5-C6-O6
3	H	3	BMA	O5-C5-C6-O6
6	M	3	BMA	C4-C5-C6-O6
4	I	2	NAG	O5-C5-C6-O6
4	K	2	NAG	O5-C5-C6-O6
3	G	1	NAG	C4-C5-C6-O6
6	M	4	MAN	C4-C5-C6-O6
6	M	3	BMA	O5-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	J	1	NAG	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.














5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	NAG	F	201	2	14,14,15	0.57	0	17,19,21	2.45	4 (23%)
7	NAG	E	501	1	14,14,15	0.50	0	17,19,21	1.35	1 (5%)
7	NAG	E	506	1	14,14,15	0.51	0	17,19,21	1.15	0
8	SIA	A	410	-	21,21,21	1.31	2 (9%)	25,31,31	0.95	2 (8%)
7	NAG	C	502	1	14,14,15	0.64	0	17,19,21	1.67	1 (5%)
7	NAG	C	501	1	14,14,15	0.61	0	17,19,21	2.47	5 (29%)
9	TAM	E	507	-	7,10,10	0.70	0	9,12,12	0.90	0
7	NAG	A	404	1	14,14,15	0.60	0	17,19,21	1.36	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	F	201	2	-	0/6/23/26	0/1/1/1
7	NAG	E	501	1	-	0/6/23/26	0/1/1/1
7	NAG	E	506	1	-	0/6/23/26	0/1/1/1
8	SIA	A	410	-	-	1/20/38/38	0/1/1/1
7	NAG	C	502	1	-	2/6/23/26	0/1/1/1
7	NAG	C	501	1	-	2/6/23/26	0/1/1/1
9	TAM	E	507	-	-	2/12/12/12	-
7	NAG	A	404	1	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	410	SIA	O2-C2	4.15	1.45	1.39
8	A	410	SIA	C3-C2	2.42	1.54	1.51

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	501	NAG	C1-O5-C5	8.20	123.30	112.19
7	F	201	NAG	C1-O5-C5	7.92	122.93	112.19
7	C	502	NAG	C1-O5-C5	5.21	119.25	112.19
7	E	501	NAG	C1-O5-C5	4.94	118.88	112.19
7	F	201	NAG	C3-C4-C5	3.49	116.47	110.24
7	C	501	NAG	C3-C4-C5	2.73	115.12	110.24
7	C	501	NAG	O5-C5-C4	2.72	117.44	110.83
7	F	201	NAG	O5-C5-C4	2.66	117.29	110.83
7	F	201	NAG	C4-C3-C2	2.60	114.83	111.02
7	A	404	NAG	C1-O5-C5	2.43	115.49	112.19
7	C	501	NAG	O5-C1-C2	2.38	115.04	111.29
8	A	410	SIA	O1A-C1-C2	-2.32	120.08	123.59
7	A	404	NAG	O5-C5-C6	2.18	110.62	107.20
7	C	501	NAG	C2-N2-C7	2.12	125.92	122.90
7	A	404	NAG	C1-C2-N2	2.12	114.11	110.49
8	A	410	SIA	O2-C2-C1	-2.09	106.50	110.76

There are no chirality outliers.


All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
9	E	507	TAM	C-C1-C4-O4
7	C	501	NAG	O5-C5-C6-O6
7	A	404	NAG	O5-C5-C6-O6
7	C	502	NAG	O5-C5-C6-O6
7	C	502	NAG	C4-C5-C6-O6
7	A	404	NAG	C4-C5-C6-O6
7	C	501	NAG	C4-C5-C6-O6
9	E	507	TAM	C-C2-C5-O5
8	A	410	SIA	C4-C5-N5-C10

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	321/323 (99%)	0.21	10 (3%) 49 61	25, 44, 77, 98	0
1	C	317/323 (98%)	0.17	8 (2%) 57 67	28, 43, 64, 94	0
1	E	317/323 (98%)	0.15	6 (1%) 66 76	21, 35, 58, 102	0
2	B	172/174 (98%)	0.76	25 (14%) 2 4	23, 56, 85, 115	0
2	D	171/174 (98%)	0.57	19 (11%) 5 8	26, 55, 84, 100	0
2	F	171/174 (98%)	0.60	14 (8%) 11 17	24, 56, 95, 108	0
All	All	1469/1491 (98%)	0.34	82 (5%) 24 35	21, 44, 83, 115	0

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	9	PRO	6.1
2	B	29	SER	5.0
2	D	171	PHE	4.8
2	B	143	LYS	4.8
2	B	30	GLU	4.7
2	B	147	ALA	4.6
2	D	162	TYR	4.4
2	B	167	LEU	4.3
1	E	222	TRP	4.2
2	F	147	ALA	4.2
2	B	148	CYS	4.2
2	B	145	ASP	3.7
1	C	9	PRO	3.7
2	B	146	ASN	3.6
2	F	23	GLY	3.6
1	E	22	ASN	3.5
2	D	168	ASN	3.5
1	C	12	THR	3.5
1	A	164	LEU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	154	ASN	3.4
1	C	22	ASN	3.3
2	F	168	ASN	3.3
2	B	164	ASP	3.0
2	D	167	LEU	3.0
2	F	166	ALA	3.0
1	C	225	MET	2.9
2	B	160	ASP	2.9
2	B	31	GLY	2.9
2	D	141	TYR	2.9
1	A	160	THR	2.8
2	D	59	THR	2.8
2	B	168	ASN	2.8
2	D	157	TYR	2.8
2	F	164	ASP	2.8
2	D	161	VAL	2.8
2	F	143	LYS	2.8
2	F	141	TYR	2.8
1	C	222	TRP	2.8
2	D	164	ASP	2.7
2	F	171	PHE	2.7
2	D	140	ILE	2.7
2	B	59	THR	2.7
2	F	2	LEU	2.7
2	B	158	ASP	2.6
2	F	29	SER	2.6
2	D	143	LYS	2.5
1	A	162	PRO	2.5
1	A	10	GLY	2.5
2	F	161	VAL	2.5
1	A	161	TYR	2.5
1	A	128	THR	2.5
1	C	221	PRO	2.4
2	B	28	ASN	2.4
2	B	144	CYS	2.4
1	A	124	GLY	2.4
2	B	18	ILE	2.4
1	E	14	CYS	2.4
1	A	208	ARG	2.4
2	D	147	ALA	2.3
2	B	172	GLN	2.3
2	D	142	HIS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	225	MET	2.3
2	B	32	THR	2.3
1	E	208	ARG	2.3
2	B	162	TYR	2.2
2	D	155	GLY	2.2
2	F	38	LEU	2.2
2	B	142	HIS	2.2
2	D	166	ALA	2.2
2	F	167	LEU	2.2
2	B	149	ILE	2.2
1	C	23	GLY	2.2
2	D	163	ARG	2.1
2	B	156	THR	2.1
2	D	169	ASN	2.1
2	D	150	GLU	2.1
2	D	158	ASP	2.1
2	F	16	GLY	2.1
2	B	171	PHE	2.0
1	A	129	GLY	2.0
1	C	208	ARG	2.0
1	E	159	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BMA	J	3	11/12	0.51	0.31	78,81,85,88	0
3	BMA	H	3	11/12	0.64	0.30	92,97,100,101	0
4	NAG	I	2	14/15	0.73	0.32	68,81,87,91	0
5	GAL	L	1	12/12	0.73	0.34	89,99,106,108	0
6	MAN	M	4	11/12	0.77	0.33	81,84,86,89	0
3	BMA	G	3	11/12	0.81	0.30	93,96,99,100	0
5	SIA	L	2	20/21	0.82	0.20	61,68,76,76	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	BMA	M	3	11/12	0.84	0.18	67,72,75,79	0
4	NAG	K	2	14/15	0.85	0.23	69,75,78,78	0
5	GAL	N	1	12/12	0.87	0.53	70,84,92,92	0
3	NAG	G	2	14/15	0.88	0.42	82,89,92,96	0
3	NAG	H	2	14/15	0.90	0.32	68,75,87,90	0
3	NAG	J	2	14/15	0.90	0.14	48,53,64,72	0
3	NAG	H	1	14/15	0.91	0.23	61,65,78,79	0
3	NAG	G	1	14/15	0.92	0.21	55,65,69,75	0
4	NAG	I	1	14/15	0.93	0.15	42,46,54,65	0
6	NAG	M	2	14/15	0.93	0.15	43,47,53,57	0
5	SIA	N	2	20/21	0.94	0.19	42,51,55,56	0
4	NAG	K	1	14/15	0.94	0.14	41,48,52,60	0
6	NAG	M	1	14/15	0.95	0.21	44,48,52,57	0
3	NAG	J	1	14/15	0.95	0.10	38,43,46,48	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain J:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain K:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain L:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain N:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	NAG	C	502	14/15	0.61	0.38	71,80,89,91	0
8	SIA	A	410	21/21	0.70	0.26	77,93,99,103	0
9	TAM	E	507	11/11	0.73	0.25	72,79,82,83	0
7	NAG	F	201	14/15	0.76	0.46	98,104,117,121	0
7	NAG	E	501	14/15	0.82	0.21	56,62,66,69	0
7	NAG	C	501	14/15	0.85	0.22	59,69,75,77	0
7	NAG	A	404	14/15	0.90	0.21	56,62,65,66	0
7	NAG	E	506	14/15	0.93	0.19	34,36,45,45	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.