



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 25, 2023 – 09:59 AM EDT

PDB ID : 5VTX
Title : Crystal structure of the A/Hong Kong/1/1968 (H3N2) influenza virus hemagglutinin G225M/L226T/S228A mutant apo form
Authors : Wu, N.C.; Wilson, I.A.
Deposited on : 2017-05-18
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 94%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 5%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: orange; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">94% 5% ..</p>
1	C	323	<div style="width: 93%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 5%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div>
1	E	323	<div style="width: 2%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 93%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 5%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div>
2	B	174	<div style="width: 2%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 95%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: orange; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div>
2	D	174	<div style="width: 2%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 95%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: orange; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	174	 2% 94%
3	G	3	 33% 67%
3	I	3	 33% 67%
4	H	2	 50% 50%
4	J	2	 50% 50%
4	L	2	 50% 50%
5	K	4	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	H	2	-	-	-	X
4	NAG	L	2	-	-	-	X
5	MAN	K	4	-	-	-	X
6	NAG	D	201	-	-	-	X
6	NAG	F	201	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 12092 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin HA1 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	321	Total 2481	C 1553	N 438	O 476	S 14	0	0	0
1	C	317	Total 2445	C 1532	N 429	O 470	S 14	0	0	0
1	E	317	Total 2445	C 1532	N 429	O 470	S 14	0	0	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	ALA	-	expression tag	UNP Q91MA7
A	8	ASP	-	expression tag	UNP Q91MA7
A	9	PRO	-	expression tag	UNP Q91MA7
A	10	GLY	-	expression tag	UNP Q91MA7
A	225	MET	GLY	engineered mutation	UNP Q91MA7
A	226	THR	LEU	engineered mutation	UNP Q91MA7
A	228	ALA	SER	engineered mutation	UNP Q91MA7
C	7	ALA	-	expression tag	UNP Q91MA7
C	8	ASP	-	expression tag	UNP Q91MA7
C	9	PRO	-	expression tag	UNP Q91MA7
C	10	GLY	-	expression tag	UNP Q91MA7
C	225	MET	GLY	engineered mutation	UNP Q91MA7
C	226	THR	LEU	engineered mutation	UNP Q91MA7
C	228	ALA	SER	engineered mutation	UNP Q91MA7
E	7	ALA	-	expression tag	UNP Q91MA7
E	8	ASP	-	expression tag	UNP Q91MA7
E	9	PRO	-	expression tag	UNP Q91MA7
E	10	GLY	-	expression tag	UNP Q91MA7
E	225	MET	GLY	engineered mutation	UNP Q91MA7
E	226	THR	LEU	engineered mutation	UNP Q91MA7
E	228	ALA	SER	engineered mutation	UNP Q91MA7


- Molecule 2 is a protein called Hemagglutinin HA2 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	172	1391	863	243	279	6	0	0	0
2	D	171	1382	858	241	277	6	0	0	0
2	F	171	1382	858	241	277	6	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	123	GLY	ARG	conflict	UNP Q91MA7
D	123	GLY	ARG	conflict	UNP Q91MA7
F	123	GLY	ARG	conflict	UNP Q91MA7

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	G	3	39	22	2	15	0	0	0
3	I	3	39	22	2	15	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	H	2	28	16	2	10	0	0	0
4	J	2	28	16	2	10	0	0	0
4	L	2	28	16	2	10	0	0	0

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranos e-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	K	4	50	28	2	20	0	0	0

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
6	A	1	14	8	1	5	0	0
6	A	1	14	8	1	5	0	0
6	C	1	14	8	1	5	0	0
6	C	1	14	8	1	5	0	0
6	D	1	14	8	1	5	0	0
6	E	1	14	8	1	5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
6	F	1	14	8	1	5	0	0



- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	48	Total 48	O 48	0	0
7	B	23	Total 23	O 23	0	0
7	C	53	Total 53	O 53	0	0
7	D	23	Total 23	O 23	0	0
7	E	82	Total 82	O 82	0	0
7	F	27	Total 27	O 27	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hemagglutinin HA1 chain




- Molecule 1: Hemagglutinin HA1 chain




- Molecule 1: Hemagglutinin HA1 chain





- Molecule 2: Hemagglutinin HA2 chain



- Molecule 2: Hemagglutinin HA2 chain



- Molecule 2: Hemagglutinin HA2 chain



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1
MAG2
BMA3
MAY4

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	208.59Å 131.06Å 72.25Å 90.00° 97.98° 90.00°	Depositor
Resolution (Å)	50.00 – 2.65 39.40 – 2.65	Depositor EDS
% Data completeness (in resolution range)	95.0 (50.00-2.65) 95.1 (39.40-2.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.92 (at 2.65Å)	Xtrriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.196 , 0.217 0.198 , 0.220	Depositor DCC
R_{free} test set	2676 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	40.1	Xtrriage
Anisotropy	0.297	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 34.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12092	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, BMA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/2538	0.71	0/3457
1	C	0.50	0/2502	0.70	1/3410 (0.0%)
1	E	0.54	0/2502	0.71	0/3410
2	B	0.49	0/1415	0.68	0/1902
2	D	0.49	0/1406	0.68	0/1890
2	F	0.50	0/1406	0.66	1/1890 (0.1%)
All	All	0.51	0/11769	0.69	2/15959 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	38	LEU	CA-CB-CG	5.49	127.92	115.30
1	C	208	ARG	NE-CZ-NH1	5.03	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2481	0	2433	5	1
1	C	2445	0	2392	8	0
1	E	2445	0	2393	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1391	0	1307	3	0
2	D	1382	0	1298	2	0
2	F	1382	0	1298	4	0
3	G	39	0	34	0	0
3	I	39	0	34	2	0
4	H	28	0	25	0	0
4	J	28	0	25	0	0
4	L	28	0	25	0	0
5	K	50	0	43	0	0
6	A	28	0	26	0	0
6	C	28	0	26	0	0
6	D	14	0	13	0	0
6	E	14	0	13	0	0
6	F	14	0	13	0	0
7	A	48	0	0	0	0
7	B	23	0	0	0	0
7	C	53	0	0	0	0
7	D	23	0	0	0	0
7	E	82	0	0	0	0
7	F	27	0	0	0	0
All	All	12092	0	11398	23	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:212:THR:HG21	1:E:216:ASN:HB3	1.47	0.93
1:C:212:THR:HG21	1:E:216:ASN:CB	2.20	0.71
2:F:53:ASN:HA	2:F:56:ILE:HG22	1.80	0.64
1:C:212:THR:CG2	1:E:216:ASN:HB3	2.27	0.63
1:A:53:ASN:HD22	1:A:54:ASN:N	2.07	0.52
2:B:58:LYS:N	2:B:58:LYS:HD3	2.25	0.51
1:C:53:ASN:HD21	1:C:276:THR:HG22	1.77	0.49
2:F:146:ASN:O	2:F:150:GLU:CG	2.63	0.47
1:C:167:THR:HG21	3:I:2:NAG:H82	1.97	0.47
2:F:146:ASN:O	2:F:150:GLU:HG3	2.15	0.46
1:A:77:ASP:OD2	1:A:141:ARG:NH1	2.43	0.45
1:A:53:ASN:HD22	1:A:53:ASN:C	2.19	0.44
1:C:77:ASP:OD2	1:C:141:ARG:NH1	2.43	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:325:GLU:O	1:A:327:GLN:NE2	2.51	0.44
1:C:167:THR:OG1	3:I:1:NAG:H62	2.17	0.43
2:D:154:ASN:O	2:D:154:ASN:OD1	2.36	0.43
1:E:77:ASP:OD2	1:E:141:ARG:NH1	2.45	0.42
1:E:203:THR:HG23	1:E:212:THR:HG22	2.02	0.42
1:E:66:LEU:HD22	1:E:267:ILE:HD12	2.03	0.41
1:C:66:LEU:HD22	1:C:267:ILE:HD12	2.03	0.41
2:B:86:ASP:OD2	2:F:62:LYS:NZ	2.47	0.40
1:A:220:ARG:HB3	1:A:221:PRO:CD	2.52	0.40
2:B:62:LYS:NZ	2:D:86:ASP:OD2	2.44	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:31:ASP:OD1	1:A:329:ARG:NH1[2_556]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	319/323 (99%)	311 (98%)	7 (2%)	1 (0%)	41	56
1	C	315/323 (98%)	308 (98%)	6 (2%)	1 (0%)	41	56
1	E	315/323 (98%)	307 (98%)	7 (2%)	1 (0%)	41	56
2	B	170/174 (98%)	162 (95%)	8 (5%)	0	100	100
2	D	169/174 (97%)	160 (95%)	9 (5%)	0	100	100
2	F	169/174 (97%)	161 (95%)	8 (5%)	0	100	100
All	All	1457/1491 (98%)	1409 (97%)	45 (3%)	3 (0%)	47	64

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	62	ILE
1	C	62	ILE
1	E	62	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	282/283 (100%)	272 (96%)	10 (4%)	36	52
1	C	278/283 (98%)	271 (98%)	7 (2%)	47	66
1	E	278/283 (98%)	269 (97%)	9 (3%)	39	56
2	B	146/148 (99%)	142 (97%)	4 (3%)	44	63
2	D	145/148 (98%)	140 (97%)	5 (3%)	37	53
2	F	145/148 (98%)	141 (97%)	4 (3%)	43	61
All	All	1274/1293 (98%)	1235 (97%)	39 (3%)	40	57

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	HIS
1	A	53	ASN
1	A	189	GLN
1	A	210	GLN
1	A	219	SER
1	A	238	LYS
1	A	276	THR
1	A	326	LYS
1	A	327	GLN
1	A	329	ARG
2	B	12	ASN
2	B	40	SER
2	B	58	LYS
2	B	172	GLN
1	C	18	HIS
1	C	31	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	167	THR
1	C	189	GLN
1	C	210	GLN
1	C	216	ASN
1	C	219	SER
2	D	12	ASN
2	D	51	LYS
2	D	59	THR
2	D	60	ASN
2	D	154	ASN
1	E	18	HIS
1	E	32	ASP
1	E	57	ARG
1	E	189	GLN
1	E	201	ARG
1	E	216	ASN
1	E	219	SER
1	E	238	LYS
1	E	325	GLU
2	F	12	ASN
2	F	38	LEU
2	F	56	ILE
2	F	150	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	GLN
1	A	53	ASN
1	A	327	GLN
2	F	27	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates i

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	G	1	3,1	14,14,15	0.41	0	17,19,21	1.20	2 (11%)
3	NAG	G	2	3	14,14,15	0.34	0	17,19,21	0.79	0
3	BMA	G	3	3	11,11,12	0.62	0	15,15,17	1.70	4 (26%)
4	NAG	H	1	4,1	14,14,15	0.54	0	17,19,21	0.67	0
4	NAG	H	2	4	14,14,15	0.38	0	17,19,21	1.39	1 (5%)
3	NAG	I	1	3,1	14,14,15	0.46	0	17,19,21	1.46	1 (5%)
3	NAG	I	2	3	14,14,15	0.44	0	17,19,21	0.99	2 (11%)
3	BMA	I	3	3	11,11,12	0.97	0	15,15,17	2.19	3 (20%)
4	NAG	J	1	4,1	14,14,15	0.33	0	17,19,21	0.86	0
4	NAG	J	2	4	14,14,15	0.45	0	17,19,21	1.53	3 (17%)
5	NAG	K	1	5,1	14,14,15	0.34	0	17,19,21	1.53	3 (17%)
5	NAG	K	2	5	14,14,15	0.27	0	17,19,21	0.92	2 (11%)
5	BMA	K	3	5	11,11,12	0.41	0	15,15,17	1.18	1 (6%)
5	MAN	K	4	5	11,11,12	0.49	0	15,15,17	1.20	1 (6%)
4	NAG	L	1	4,1	14,14,15	0.43	0	17,19,21	0.91	0
4	NAG	L	2	4	14,14,15	0.61	0	17,19,21	1.29	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	G	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	G	2	3	-	2/6/23/26	0/1/1/1
3	BMA	G	3	3	-	1/2/19/22	0/1/1/1
4	NAG	H	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	H	2	4	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	I	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	I	2	3	-	2/6/23/26	0/1/1/1
3	BMA	I	3	3	-	0/2/19/22	0/1/1/1
4	NAG	J	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	J	2	4	-	0/6/23/26	0/1/1/1
5	NAG	K	1	5,1	-	0/6/23/26	0/1/1/1
5	NAG	K	2	5	-	0/6/23/26	0/1/1/1
5	BMA	K	3	5	-	0/2/19/22	0/1/1/1
5	MAN	K	4	5	-	2/2/19/22	0/1/1/1
4	NAG	L	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	L	2	4	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	3	BMA	C1-C2-C3	5.94	116.96	109.67
3	I	1	NAG	C1-O5-C5	5.31	119.38	112.19
4	H	2	NAG	C1-O5-C5	5.01	118.99	112.19
4	J	2	NAG	C1-O5-C5	4.44	118.20	112.19
5	K	4	MAN	C1-O5-C5	3.81	117.36	112.19
5	K	1	NAG	C1-O5-C5	3.79	117.32	112.19
3	I	3	BMA	O5-C1-C2	3.72	116.51	110.77
3	G	3	BMA	C1-C2-C3	3.65	114.15	109.67
5	K	1	NAG	O5-C1-C2	-3.33	106.02	111.29
3	I	3	BMA	C1-O5-C5	3.25	116.60	112.19
3	G	3	BMA	O5-C1-C2	2.96	115.34	110.77
3	G	3	BMA	C1-O5-C5	2.85	116.05	112.19
3	G	1	NAG	O5-C1-C2	-2.79	106.88	111.29
3	G	3	BMA	O5-C5-C6	2.70	111.44	107.20
3	G	1	NAG	C1-O5-C5	2.69	115.83	112.19
5	K	2	NAG	C1-C2-N2	2.33	114.47	110.49
4	L	2	NAG	C1-O5-C5	2.29	115.29	112.19
4	J	2	NAG	O5-C1-C2	-2.28	107.69	111.29
5	K	2	NAG	C1-O5-C5	2.27	115.27	112.19
5	K	3	BMA	C3-C4-C5	2.23	114.22	110.24
5	K	1	NAG	C3-C4-C5	2.21	114.18	110.24
3	I	2	NAG	C1-O5-C5	2.10	115.03	112.19
3	I	2	NAG	O5-C5-C6	2.06	110.44	107.20
4	J	2	NAG	C4-C3-C2	-2.04	108.03	111.02

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	I	2	NAG	O5-C5-C6-O6
5	K	4	MAN	C4-C5-C6-O6
5	K	4	MAN	O5-C5-C6-O6
4	H	1	NAG	O5-C5-C6-O6
3	I	2	NAG	C4-C5-C6-O6
4	H	1	NAG	C4-C5-C6-O6
3	G	2	NAG	O5-C5-C6-O6
3	G	2	NAG	C4-C5-C6-O6
3	G	3	BMA	O5-C5-C6-O6


There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	1	NAG	1	0
3	I	2	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	C	502	1	14,14,15	0.53	0	17,19,21	1.18	2 (11%)
6	NAG	A	401	1	14,14,15	0.50	0	17,19,21	1.33	1 (5%)
6	NAG	F	201	2	14,14,15	0.67	0	17,19,21	2.18	5 (29%)
6	NAG	C	501	1	14,14,15	0.72	0	17,19,21	2.12	4 (23%)
6	NAG	A	402	1	14,14,15	0.68	0	17,19,21	1.43	2 (11%)
6	NAG	D	201	2	14,14,15	0.41	0	17,19,21	0.73	0
6	NAG	E	501	1	14,14,15	0.52	0	17,19,21	1.18	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	C	502	1	-	2/6/23/26	0/1/1/1
6	NAG	A	401	1	-	0/6/23/26	0/1/1/1
6	NAG	F	201	2	-	0/6/23/26	0/1/1/1
6	NAG	C	501	1	-	2/6/23/26	0/1/1/1
6	NAG	A	402	1	-	2/6/23/26	0/1/1/1
6	NAG	D	201	2	-	2/6/23/26	0/1/1/1
6	NAG	E	501	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	F	201	NAG	C1-O5-C5	6.69	121.25	112.19
6	C	501	NAG	C1-O5-C5	5.81	120.06	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	402	NAG	C1-O5-C5	3.77	117.30	112.19
6	A	401	NAG	C1-O5-C5	3.75	117.28	112.19
6	E	501	NAG	C1-O5-C5	3.30	116.66	112.19
6	C	501	NAG	O5-C5-C6	3.27	112.33	107.20
6	C	501	NAG	C3-C4-C5	3.16	115.87	110.24
6	F	201	NAG	C3-C4-C5	3.12	115.80	110.24
6	F	201	NAG	O5-C5-C4	2.40	116.66	110.83
6	A	402	NAG	C4-C3-C2	-2.38	107.53	111.02
6	F	201	NAG	C2-N2-C7	2.26	126.12	122.90
6	C	502	NAG	C1-C2-N2	2.09	114.05	110.49
6	C	502	NAG	O5-C1-C2	-2.07	108.02	111.29
6	F	201	NAG	C4-C3-C2	2.06	114.03	111.02
6	C	501	NAG	O5-C1-C2	2.05	114.53	111.29

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	C	502	NAG	O5-C5-C6-O6
6	A	402	NAG	O5-C5-C6-O6
6	C	501	NAG	C4-C5-C6-O6
6	A	402	NAG	C4-C5-C6-O6
6	C	502	NAG	C4-C5-C6-O6
6	C	501	NAG	O5-C5-C6-O6
6	D	201	NAG	C4-C5-C6-O6
6	D	201	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	321/323 (99%)	-0.17	3 (0%) 84 83	25, 46, 78, 112	0
1	C	317/323 (98%)	-0.28	0 100 100	29, 44, 63, 104	0
1	E	317/323 (98%)	-0.30	4 (1%) 77 75	21, 34, 56, 96	0
2	B	172/174 (98%)	-0.03	3 (1%) 70 67	23, 58, 86, 101	0
2	D	171/174 (98%)	0.17	2 (1%) 79 77	27, 57, 86, 104	0
2	F	171/174 (98%)	0.13	4 (2%) 60 56	23, 58, 92, 105	0
All	All	1469/1491 (98%)	-0.13	16 (1%) 80 79	21, 44, 83, 112	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	9	PRO	3.7
2	F	164	ASP	3.4
2	B	31	GLY	3.2
2	D	33	GLY	3.1
1	E	189	GLN	2.9
2	F	29	SER	2.8
1	E	22	ASN	2.7
1	A	143	PRO	2.5
2	F	33	GLY	2.2
2	F	30	GLU	2.1
2	B	147	ALA	2.1
1	E	159	SER	2.1
1	A	208	ARG	2.1
2	B	30	GLU	2.1
2	D	30	GLU	2.0
1	E	9	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	H	2	14/15	0.72	0.43	61,82,87,87	0
4	NAG	L	2	14/15	0.73	0.41	79,85,92,95	0
3	BMA	I	3	11/12	0.77	0.29	72,78,81,82	0
5	MAN	K	4	11/12	0.77	0.41	89,93,96,96	0
5	BMA	K	3	11/12	0.84	0.20	77,80,84,88	0
4	NAG	J	2	14/15	0.86	0.27	66,71,79,79	0
3	BMA	G	3	11/12	0.87	0.24	69,71,72,72	0
5	NAG	K	2	14/15	0.90	0.17	47,59,62,67	0
3	NAG	G	1	14/15	0.91	0.21	56,61,70,71	0
4	NAG	H	1	14/15	0.92	0.23	40,45,53,64	0
3	NAG	I	2	14/15	0.94	0.18	44,53,63,68	0
4	NAG	L	1	14/15	0.94	0.17	37,45,52,65	0
3	NAG	G	2	14/15	0.94	0.29	64,66,72,73	0
5	NAG	K	1	14/15	0.95	0.13	40,47,51,53	0
4	NAG	J	1	14/15	0.96	0.15	35,41,49,56	0
3	NAG	I	1	14/15	0.96	0.11	34,39,44,45	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain I:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain J:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain L:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain K:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	NAG	D	201	14/15	0.62	0.43	91,96,99,100	0
6	NAG	C	502	14/15	0.69	0.33	67,77,82,82	0
6	NAG	F	201	14/15	0.71	0.42	99,105,114,116	0
6	NAG	C	501	14/15	0.83	0.27	56,68,73,74	0
6	NAG	A	401	14/15	0.84	0.23	48,54,58,58	0
6	NAG	E	501	14/15	0.89	0.27	53,60,62,62	0
6	NAG	A	402	14/15	0.92	0.23	47,55,56,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.