



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 06:03 pm BST

PDB ID : 2W1C  
Title : Structure determination of Aurora Kinase in complex with inhibitor  
Authors : Howard, S.; Berdini, V.; Boulstridge, J.A.; Carr, M.G.; Cross, D.M.; Curry, J.; Devine, L.A.; Early, T.R.; Fazal, L.; Gill, A.L.; Heathcote, M.; Maman, S.; Matthews, J.E.; McMenamin, R.L.; Navarro, E.F.; O'Brien, M.A.; O'Reilly, M.; Rees, D.C.; Reule, M.; Tisi, D.; Williams, G.; Vinkovic, M.; Wyatt, P.G.  
Deposited on : 2008-10-17  
Resolution : 3.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1619 (3.28-3.20)
Clashscore	141614	1755 (3.28-3.20)
Ramachandran outliers	138981	1728 (3.28-3.20)
Sidechain outliers	138945	1727 (3.28-3.20)
RSRZ outliers	127900	1567 (3.28-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	TPO	A	287	-	-	-	X
1	TPO	A	288	-	-	-	X

## 2 Entry composition [i](#)


There are 2 unique types of molecules in this entry. The entry contains 2211 atoms, of which 22 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SERINE/THREONINE-PROTEIN KINASE 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	P				S
1	A	264	2158	1381	381	389	2	5	0	0	1

- Molecule 2 is 4-{2-[4-{{(4-FLUOROPHENYL)CARBONYLAMINO}-1H-PYRAZOL-3-YL)-1H-BENZIMIDAZOL-6-YL]METHYL}MORPHOLIN-4-IUM (three-letter code: LOC) (formula: C<sub>22</sub>H<sub>22</sub>FN<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	F	H	N			O
2	A	1	53	22	1	22	6	2	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: SERINE/THREONINE-PROTEIN KINASE 6



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.86Å 82.86Å 168.19Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	54.59 – 3.24 54.59 – 3.24	Depositor EDS
% Data completeness (in resolution range)	94.7 (54.59-3.24) 94.7 (54.59-3.24)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.36 (at 3.26Å)	Xtrriage
Refinement program	REFMAC 5.4.0069A	Depositor
R, $R_{free}$	0.244 , 0.288 0.231 , 0.284	Depositor DCC
$R_{free}$ test set	283 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	99.1	Xtrriage
Anisotropy	0.041	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 51.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	2211	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	83.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.40% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, LOC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.49	0/2189	0.69	0/2958

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2158	0	2145	99	0
2	A	31	22	22	7	0
All	All	2189	22	2167	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:221:GLU:HG2	1:A:232:ARG:NH2	1.86	0.90
1:A:293:LEU:HD13	1:A:301:ILE:HD11	1.66	0.77

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288:TPO:O1P	1:A:288:TPO:HG21	1.85	0.75
1:A:162:LYS:HB3	1:A:208:LEU:HB2	1.69	0.74
2:A:1391:L0C:H52	2:A:1391:L0C:C21	2.16	0.74
1:A:358:ASP:OD2	1:A:362:ARG:NE	2.20	0.73
2:A:1391:L0C:H52	2:A:1391:L0C:H42	1.71	0.71
1:A:133:PHE:HE2	1:A:159:LEU:HD21	1.58	0.68
1:A:373:MET:HE2	1:A:373:MET:HA	1.74	0.67
1:A:344:VAL:O	1:A:344:VAL:HG12	1.94	0.67
2:A:1391:L0C:C30	2:A:1391:L0C:H42	2.25	0.66
1:A:309:LYS:NZ	1:A:371:ARG:O	2.21	0.65
1:A:310:VAL:HG23	1:A:311:ASP:N	2.12	0.65
1:A:389:LYS:HG3	1:A:390:HIS:N	2.13	0.64
1:A:287:TPO:O3P	1:A:287:TPO:HG21	1.98	0.62
1:A:285:ARG:HH22	1:A:303:GLY:HA2	1.65	0.61
1:A:312:LEU:HD22	1:A:372:PRO:O	2.01	0.61
1:A:317:VAL:HG13	1:A:328:PRO:HD2	1.81	0.61
1:A:367:ASN:HB3	1:A:370:GLN:NE2	2.16	0.60
1:A:240:LEU:HD12	1:A:272:ILE:HD11	1.84	0.60
1:A:144:PHE:HB2	1:A:164:LEU:HD22	1.82	0.60
2:A:1391:L0C:C21	2:A:1391:L0C:C30	2.80	0.60
1:A:240:LEU:CD1	1:A:272:ILE:HD11	2.32	0.59
1:A:194:LEU:HD22	1:A:263:LEU:HD12	1.86	0.58
1:A:338:TYR:O	1:A:342:SER:HB2	2.03	0.58
1:A:219:TYR:CD1	1:A:260:GLU:HG2	2.40	0.57
1:A:244:LEU:HD21	1:A:257:ILE:HD11	1.84	0.57
1:A:236:TYR:CD1	1:A:270:LEU:HD13	2.40	0.56
1:A:231:GLN:O	1:A:235:THR:OG1	2.24	0.56
1:A:281:ALA:HB1	1:A:282:PRO:CD	2.35	0.56
1:A:293:LEU:HD22	1:A:301:ILE:HD11	1.88	0.55
1:A:152:GLU:OE2	1:A:155:SER:HB2	2.07	0.54
1:A:184:ILE:O	1:A:188:LEU:HD13	2.08	0.54
1:A:285:ARG:O	1:A:285:ARG:HD2	2.08	0.54
1:A:377:VAL:O	1:A:383:ILE:HG13	2.09	0.53
1:A:152:GLU:OE2	1:A:155:SER:CB	2.57	0.52
1:A:184:ILE:HG21	1:A:252:VAL:HG11	1.92	0.52
1:A:147:VAL:HG21	2:A:1391:L0C:O9	2.10	0.51
1:A:324:VAL:HG13	1:A:349:PRO:HG2	1.93	0.51
1:A:190:HIS:HB3	1:A:193:ILE:HG13	1.93	0.51
1:A:137:ARG:NH2	2:A:1391:L0C:H44	2.26	0.50
1:A:236:TYR:O	1:A:240:LEU:HB2	2.12	0.49
1:A:195:ARG:HB2	1:A:211:GLU:HB2	1.95	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:215:LEU:HB2	1:A:264:LEU:HB2	1.94	0.49
1:A:353:THR:O	1:A:357:ARG:HG3	2.13	0.49
1:A:219:TYR:HE2	1:A:223:GLN:OE1	1.95	0.49
1:A:184:ILE:HG22	1:A:188:LEU:HD22	1.95	0.49
1:A:297:PRO:HB3	1:A:313:TRP:CG	2.47	0.49
1:A:310:VAL:CG2	1:A:311:ASP:N	2.76	0.48
1:A:298:PRO:HD3	1:A:313:TRP:CZ2	2.47	0.48
1:A:373:MET:HA	1:A:373:MET:CE	2.42	0.48
1:A:281:ALA:HB1	1:A:282:PRO:HD2	1.94	0.48
1:A:330:GLU:O	1:A:340:ARG:NH1	2.46	0.48
1:A:163:VAL:HG13	1:A:207:TYR:CE2	2.49	0.47
1:A:257:ILE:HG22	1:A:318:LEU:HD21	1.97	0.47
1:A:389:LYS:CG	1:A:390:HIS:N	2.71	0.47
1:A:380:HIS:CG	1:A:381:PRO:HD2	2.50	0.47
1:A:331:ALA:HB3	1:A:337:THR:OG1	2.14	0.47
1:A:166:LYS:HD2	1:A:203:ALA:O	2.15	0.47
1:A:293:LEU:CD1	1:A:301:ILE:HD11	2.41	0.47
1:A:351:PHE:C	1:A:351:PHE:CD2	2.88	0.46
1:A:293:LEU:HD22	1:A:301:ILE:CD1	2.46	0.46
1:A:235:THR:HG23	1:A:389:LYS:HB3	1.97	0.46
1:A:367:ASN:CB	1:A:370:GLN:NE2	2.79	0.46
1:A:200:PHE:CE1	1:A:207:TYR:CD1	3.04	0.46
1:A:183:GLU:HG3	1:A:184:ILE:N	2.31	0.45
1:A:221:GLU:HG2	1:A:232:ARG:HH21	1.75	0.45
1:A:299:GLU:OE2	1:A:371:ARG:NH1	2.45	0.45
1:A:190:HIS:CG	1:A:191:PRO:HD2	2.51	0.45
1:A:266:SER:C	1:A:268:GLY:H	2.18	0.45
1:A:137:ARG:HA	1:A:138:PRO:HD3	1.78	0.44
1:A:196:LEU:HD13	1:A:210:LEU:CD2	2.48	0.44
1:A:285:ARG:HG2	1:A:287:TPO:N	2.33	0.44
1:A:374:LEU:O	1:A:378:LEU:HD22	2.17	0.44
1:A:166:LYS:O	1:A:167:ALA:C	2.56	0.44
1:A:266:SER:C	1:A:268:GLY:N	2.71	0.44
1:A:286:ARG:HB2	1:A:286:ARG:HH11	1.84	0.43
1:A:190:HIS:ND1	1:A:191:PRO:HD2	2.34	0.43
1:A:313:TRP:CE3	1:A:364:LEU:HD23	2.54	0.43
1:A:288:TPO:O1P	1:A:288:TPO:CG2	2.59	0.43
1:A:251:ARG:HD2	1:A:281:ALA:HA	2.01	0.43
1:A:269:GLU:N	1:A:269:GLU:OE1	2.41	0.42
1:A:146:ASN:HB2	1:A:148:TYR:CE1	2.54	0.42
1:A:141:LYS:O	2:A:1391:L0C:H38	2.18	0.42

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:147:VAL:HG22	1:A:162:LYS:HG3	2.01	0.42
1:A:374:LEU:HA	1:A:374:LEU:HD23	1.93	0.42
1:A:289:LEU:O	1:A:291:GLY:N	2.53	0.42
1:A:285:ARG:HD3	1:A:287:TPO:HA	2.02	0.42
1:A:318:LEU:HD13	1:A:318:LEU:HA	1.53	0.41
1:A:316:GLY:HA2	1:A:363:LEU:HD13	2.01	0.41
1:A:186:SER:C	1:A:188:LEU:H	2.22	0.41
1:A:258:LYS:HB2	1:A:259:PRO:HD2	2.02	0.41
1:A:188:LEU:HA	1:A:188:LEU:HD12	1.85	0.41
1:A:254:HIS:NE2	1:A:274:ASP:O	2.53	0.41
1:A:184:ILE:HD13	1:A:252:VAL:HG13	2.03	0.41
1:A:299:GLU:HG2	1:A:300:MET:H	1.85	0.41
1:A:194:LEU:HD22	1:A:263:LEU:CD1	2.50	0.41
1:A:220:ARG:O	1:A:224:LYS:HB2	2.21	0.41
1:A:267:ALA:HB3	1:A:269:GLU:OE1	2.20	0.41
1:A:133:PHE:CE2	1:A:159:LEU:HD21	2.46	0.41
1:A:140:GLY:HA2	1:A:141:LYS:HZ3	1.86	0.41
1:A:215:LEU:HD23	1:A:215:LEU:HA	1.82	0.41
1:A:184:ILE:CG2	1:A:252:VAL:HG11	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	260/275 (94%)	231 (89%)	23 (9%)	6 (2%)	<b>6</b>   <b>31</b>

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	389	LYS
1	A	173	GLY

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	274	ASP
1	A	290	CYS
1	A	167	ALA
1	A	281	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/242 (94%)	211 (93%)	16 (7%)	15	46

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	152	GLU
1	A	161	LEU
1	A	174	VAL
1	A	176	HIS
1	A	183	GLU
1	A	220	ARG
1	A	231	GLN
1	A	285	ARG
1	A	286	ARG
1	A	294	ASP
1	A	305	MET
1	A	312	LEU
1	A	318	LEU
1	A	335	GLN
1	A	342	SER
1	A	373	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	127	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	146	ASN
1	A	168	GLN
1	A	177	GLN
1	A	187	HIS
1	A	223	GLN
1	A	242	ASN
1	A	335	GLN
1	A	370	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	TPO	A	287	1	8,10,11	1.64	2 (25%)	10,14,16	1.03	0
1	TPO	A	288	1	8,10,11	1.73	2 (25%)	10,14,16	1.11	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	287	1	-	3/9/11/13	-
1	TPO	A	288	1	-	5/9/11/13	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	288	TPO	P-OG1	2.97	1.64	1.59
1	A	287	TPO	P-OG1	2.73	1.64	1.59
1	A	287	TPO	P-O1P	2.62	1.59	1.50
1	A	288	TPO	P-O1P	2.60	1.58	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	288	TPO	CG2-CB-CA	-2.85	107.54	113.16

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	287	TPO	N-CA-CB-CG2
1	A	287	TPO	N-CA-CB-OG1
1	A	287	TPO	C-CA-CB-CG2
1	A	288	TPO	N-CA-CB-OG1
1	A	288	TPO	C-CA-CB-CG2
1	A	288	TPO	CB-OG1-P-O2P
1	A	288	TPO	CB-OG1-P-O3P
1	A	288	TPO	O-C-CA-CB

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	287	TPO	3	0
1	A	288	TPO	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	L0C	A	1391	-	33,35,35	1.16	3 (9%)	36,49,49	0.83	2 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	L0C	A	1391	-	-	1/10/24/24	0/5/5/5

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1391	L0C	C12-C11	-3.39	1.34	1.38
2	A	1391	L0C	C11-N10	-2.83	1.36	1.41
2	A	1391	L0C	C31-C32	-2.06	1.38	1.41

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1391	L0C	C17-N33-C32	2.73	109.19	103.78
2	A	1391	L0C	C16-C17-N18	2.06	126.36	122.27

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1391	L0C	C22-C23-N24-C30


There are no ring outliers.

1 monomer is involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1391	L0C	7	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	262/275 (95%)	0.05	6 (2%) 60 50	51, 74, 130, 167	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	169	LEU	3.3
1	A	289	LEU	3.3
1	A	286	ARG	3.0
1	A	290	CYS	2.6
1	A	277	TRP	2.5
1	A	282	PRO	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	TPO	A	288	11/12	-0.15	0.76	158,171,185,185	0
1	TPO	A	287	11/12	0.49	0.50	169,175,187,187	0

### 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	L0C	A	1391	31/31	0.83	0.47	83,100,109,109	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.