



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 5, 2022 – 04:47 PM EDT

PDB ID : 6W1W
Title : Crystal Structure of Motility Associated Killing Factor B from *Vibrio cholerae*
Authors : Kim, Y.; Welk, L.; Jedrzejczak, R.; Endres, M.; Joachimiak, A.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2020-03-04
Resolution : 2.58 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.29
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.29


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3676 (2.60-2.56)
Clashscore	141614	4049 (2.60-2.56)
Ramachandran outliers	138981	3979 (2.60-2.56)
Sidechain outliers	138945	3979 (2.60-2.56)
RSRZ outliers	127900	3614 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	355	 3% 79% 17% ..
1	B	355	 2% 83% 13% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5411 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called motility-associated killing factor MakB.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	348	2699	1707	440	541	9	2	0	2	0
1	B	348	2692	1703	438	540	9	2	0	1	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q9KL65
A	-1	ASN	-	expression tag	UNP Q9KL65
A	0	ALA	-	expression tag	UNP Q9KL65
A	?	-	GLY	deletion	UNP Q9KL65
A	?	-	ALA	deletion	UNP Q9KL65
B	-2	SER	-	expression tag	UNP Q9KL65
B	-1	ASN	-	expression tag	UNP Q9KL65
B	0	ALA	-	expression tag	UNP Q9KL65
B	?	-	GLY	deletion	UNP Q9KL65
B	?	-	ALA	deletion	UNP Q9KL65

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total O 2 2	0	0
3	B	6	Total O 6 6	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: motility-associated killing factor MakB



- Molecule 1: motility-associated killing factor MakB



4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	137.72Å 137.72Å 233.83Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	66.05 – 2.58 66.05 – 2.58	Depositor EDS
% Data completeness (in resolution range)	99.5 (66.05-2.58) 100.0 (66.05-2.58)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 2.58Å)	Xtrriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.276 , 0.286 0.275 , 0.284	Depositor DCC
R_{free} test set	4058 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	63.7	Xtrriage
Anisotropy	0.412	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 45.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.478 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5411	wwPDB-VP
Average B, all atoms (Å ²)	120.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/2736	0.40	0/3719
1	B	0.26	0/2728	0.41	0/3706
All	All	0.26	0/5464	0.40	0/7425

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2699	0	2709	33	0
1	B	2692	0	2708	25	0
2	A	8	0	12	0	0
2	B	4	0	6	0	0
3	A	2	0	0	0	0
3	B	6	0	0	0	0
All	All	5411	0	5435	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:133:LYS:NZ	1:B:269:SER:OG	2.29	0.64
1:A:4:PRO:HB3	1:A:285:LEU:HD11	1.84	0.60
1:A:143:ILE:HD12	1:A:259:LYS:HA	1.85	0.58
1:A:36:GLY:O	1:A:49:ARG:NH2	2.38	0.57
1:B:23:ILE:HG13	1:B:272:PHE:CZ	2.40	0.57
1:A:296[B]:LEU:HD21	1:A:312:LEU:HD13	1.87	0.57
1:B:31:ASN:HA	1:B:56:ARG:HD3	1.87	0.55
1:B:123:GLN:HA	1:B:126:THR:HG22	1.88	0.55
1:A:111:LEU:O	1:A:115:ILE:HG22	2.07	0.53
1:A:307:ALA:O	1:A:313:GLN:NE2	2.42	0.53
1:B:51:ARG:HE	1:B:138:LEU:HB3	1.75	0.52
1:A:144:GLN:OE1	1:A:148:ASN:ND2	2.43	0.50
1:A:323:LYS:NZ	1:A:327:GLU:OE2	2.41	0.50
1:B:85:ASP:OD1	1:B:319:SER:OG	2.28	0.50
1:B:16:VAL:O	1:B:20:PHE:HB2	2.12	0.50
1:A:111:LEU:HD13	1:A:290:LEU:HA	1.92	0.49
1:B:125:PHE:HD2	1:B:275:ILE:HG12	1.77	0.49
1:B:200:VAL:HG11	1:B:250:GLU:HG3	1.95	0.49
1:A:266:SER:O	1:A:270:GLU:HG2	2.13	0.49
1:B:51:ARG:NH2	1:B:142:SER:OG	2.47	0.48
1:A:74:ILE:HD12	1:A:333:SER:HB3	1.96	0.48
1:A:47:GLU:O	1:A:51:ARG:HG3	2.14	0.47
1:A:17:TYR:CZ	1:A:21:LEU:HD11	2.50	0.47
1:A:72:GLN:NE2	1:A:117:LEU:HB3	2.30	0.46
1:A:105:GLN:HA	1:A:108:GLN:HB3	1.97	0.46
1:B:96:SER:HB2	1:B:99:GLN:HG3	1.98	0.46
1:B:211:LEU:HD23	1:B:212:SER:N	2.30	0.45
1:A:66:LYS:HE2	1:A:66:LYS:HB3	1.78	0.45
1:B:87:CYS:HB3	1:B:103:CYS:SG	2.57	0.44
1:B:74:ILE:HD12	1:B:333:SER:HB3	1.99	0.44
1:A:54:ILE:HD12	1:B:54:ILE:HD12	2.00	0.43
1:A:144:GLN:O	1:A:148:ASN:ND2	2.45	0.43
1:B:296[B]:LEU:HD21	1:B:312:LEU:HD13	1.99	0.43
1:A:51:ARG:HD3	1:A:138:LEU:HB3	2.00	0.43
1:B:296[A]:LEU:HD11	1:B:311:ASP:HB3	2.01	0.43
1:A:279:LEU:HD23	1:A:279:LEU:HA	1.85	0.43
1:A:211:LEU:HD23	1:A:212:SER:N	2.34	0.42
1:B:125:PHE:CD2	1:B:275:ILE:HG12	2.54	0.42
1:A:316:LYS:HE3	1:A:316:LYS:HB2	1.89	0.42
1:A:74:ILE:HD13	1:A:329:VAL:HG13	2.02	0.42
1:A:133:LYS:HA	1:A:133:LYS:HD3	1.75	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:LEU:HD13	1:A:13:LEU:HA	1.88	0.41
1:A:150:LEU:HD11	1:A:251:GLU:HB3	2.01	0.41
1:B:123:GLN:HE21	1:B:123:GLN:HB2	1.61	0.41
1:A:85:ASP:OD1	1:A:319:SER:HB2	2.21	0.41
1:A:90:ASN:O	1:A:90:ASN:ND2	2.53	0.41
1:B:134:ASN:HD22	1:B:134:ASN:HA	1.64	0.41
1:A:96:SER:HB2	1:A:99:GLN:HG3	2.01	0.41
1:A:4:PRO:CB	1:A:285:LEU:HD11	2.50	0.41
1:A:178:VAL:HA	1:A:189:TYR:OH	2.20	0.41
1:A:19:ASP:HB3	1:A:275:ILE:HG12	2.03	0.41
1:B:71:ALA:HA	1:B:74:ILE:HB	2.02	0.41
1:B:313:GLN:H	1:B:313:GLN:HG2	1.70	0.41
1:A:202:MET:C	1:A:206:THR:H	2.25	0.40
1:B:29:THR:HG23	1:B:213:PHE:HD2	1.85	0.40
1:B:115:ILE:HD12	1:B:287:GLN:HG2	2.02	0.40
1:B:64:GLN:HE21	1:B:64:GLN:HB2	1.71	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	346/355 (98%)	332 (96%)	13 (4%)	1 (0%)	41	62
1	B	343/355 (97%)	325 (95%)	17 (5%)	1 (0%)	41	62
All	All	689/710 (97%)	657 (95%)	30 (4%)	2 (0%)	41	62

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	39	ALA
1	B	221	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	303/305 (99%)	282 (93%)	21 (7%)	15	30
1	B	303/305 (99%)	285 (94%)	18 (6%)	19	37
All	All	606/610 (99%)	567 (94%)	39 (6%)	17	34

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MSE
1	A	7	ASP
1	A	13	LEU
1	A	20	PHE
1	A	62	PHE
1	A	90	ASN
1	A	95	THR
1	A	103	CYS
1	A	202	MET
1	A	212	SER
1	A	220	VAL
1	A	230	SER
1	A	243	GLN
1	A	274	LYS
1	A	275	ILE
1	A	283	SER
1	A	311	ASP
1	A	316	LYS
1	A	317	ILE
1	A	342	VAL
1	A	347	LEU
1	B	1	MSE
1	B	20	PHE
1	B	62	PHE
1	B	64	GLN
1	B	66	LYS
1	B	70	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	133	LYS
1	B	134	ASN
1	B	179	THR
1	B	197	SER
1	B	212	SER
1	B	217	LEU
1	B	228	ILE
1	B	246	THR
1	B	308	LEU
1	B	311	ASP
1	B	317	ILE
1	B	347	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	HIS
1	A	64	GLN
1	A	72	GLN
1	A	89	GLN
1	A	90	ASN
1	A	108	GLN
1	A	122	GLN
1	A	123	GLN
1	A	348	ASN
1	B	25	ASN
1	B	64	GLN
1	B	99	GLN
1	B	108	GLN
1	B	113	GLN
1	B	122	GLN
1	B	123	GLN
1	B	131	GLN
1	B	134	ASN
1	B	144	GLN
1	B	182	GLN
1	B	191	GLN
1	B	287	GLN
1	B	313	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	401	-	3,3,3	0.49	0	2,2,2	0.28	0
2	EDO	B	401	-	3,3,3	0.52	0	2,2,2	0.24	0
2	EDO	A	402	-	3,3,3	0.50	0	2,2,2	0.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	401	-	-	0/1/1/1	-
2	EDO	B	401	-	-	0/1/1/1	-
2	EDO	A	402	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	346/355 (97%)	0.31	9 (2%) 56 52	85, 118, 153, 183	0
1	B	346/355 (97%)	0.31	8 (2%) 60 57	85, 118, 148, 177	0
All	All	692/710 (97%)	0.31	17 (2%) 57 53	85, 118, 152, 183	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	62	PHE	5.5
1	B	206	THR	4.1
1	A	2	LEU	3.8
1	B	70	PHE	3.7
1	A	70	PHE	3.2
1	B	62	PHE	2.7
1	A	74	ILE	2.7
1	B	2	LEU	2.6
1	A	13	LEU	2.5
1	B	208	VAL	2.4
1	B	213	PHE	2.3
1	A	262	ILE	2.2
1	A	125	PHE	2.2
1	B	125	PHE	2.2
1	B	20	PHE	2.1
1	A	296[A]	LEU	2.1
1	A	20	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	EDO	A	401	4/4	0.90	0.17	84,104,112,116	0
2	EDO	B	401	4/4	0.91	0.21	77,104,120,121	0
2	EDO	A	402	4/4	0.92	0.40	110,115,123,124	0

6.5 Other polymers [i](#)

There are no such residues in this entry.