



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 07:35 am BST

PDB ID : 1W2C  
Title : Human Inositol (1,4,5) trisphosphate 3-kinase complexed with Mn<sup>2+</sup>/AMPP  
NP/Ins(1,4,5)P<sub>3</sub>  
Authors : Gonzalez, B.; Schell, M.J.; Irvine, R.F.; Williams, R.L.  
Deposited on : 2004-07-01  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	265	 10% 86% 12% •
1	B	265	 18% 83% 11% • 6%

## 2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 4615 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INOSITOL-TRISPHOSPHATE 3-KINASE A.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	265	Total 2152	1345	395	400	12	0	3	0
1	B	249	Total 2012	1264	363	373	12	0	2	0

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula:  $C_{10}H_{17}N_6O_{12}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total 31	10	6	12	3	0	0
2	B	1	Total 31	10	6	12	3	0	0

- Molecule 3 is D-MYO-INOSITOL-1,4,5-TRIPHOSPHATE (three-letter code: I3P) (formula:  $C_6H_{15}O_{15}P_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	O			P
3	A	1	24	6	15	3	0	0
3	B	1	24	6	15	3	0	0

- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mn		
4	B	1	1	1	0	0
4	A	1	1	1	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	166	Total O 166 166	0	0
6	B	158	Total O 158 158	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: INOSITOL-TRISPHOSPHATE 3-KINASE A



- Molecule 1: INOSITOL-TRISPHOSPHATE 3-KINASE A



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	72.06Å 97.52Å 191.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.80 – 1.95 48.76 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.6 (48.80-1.95) 88.0 (48.76-1.90)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.21 (at 1.90Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.199 , 0.243 0.209 , 0.249	Depositor DCC
$R_{free}$ test set	4529 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.9	Xtrriage
Anisotropy	0.253	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 47.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4615	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.89% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: I3P, ANP, SO4, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.74	0/2204	0.92	7/2964 (0.2%)
1	B	0.74	0/2055	0.95	7/2760 (0.3%)
All	All	0.74	0/4259	0.93	14/5724 (0.2%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	339	ASP	CB-CG-OD2	7.79	125.31	118.30
1	A	262	ASP	CB-CG-OD1	7.02	124.62	118.30
1	B	376	ARG	NE-CZ-NH1	6.71	123.65	120.30
1	A	225	ASP	CB-CG-OD2	6.42	124.08	118.30
1	A	376	ARG	NE-CZ-NH1	6.39	123.49	120.30
1	B	392	ARG	NE-CZ-NH1	5.94	123.27	120.30
1	B	339	ASP	CB-CG-OD2	5.91	123.62	118.30
1	B	253	ASP	CB-CG-OD2	5.90	123.61	118.30
1	B	241	ASP	CB-CG-OD2	5.68	123.42	118.30
1	B	319	ARG	NE-CZ-NH1	5.45	123.03	120.30
1	A	319	ARG	NE-CZ-NH1	5.27	122.94	120.30
1	B	430	ASP	CB-CG-OD2	5.26	123.03	118.30
1	A	371	ARG	NE-CZ-NH2	-5.19	117.70	120.30
1	A	382	ASP	CB-CG-OD1	5.14	122.93	118.30

There are no chirality outliers.

There are no planarity outliers.



## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2152	0	2135	23	0
1	B	2012	0	2002	12	0
2	A	31	0	12	0	0
2	B	31	0	12	0	0
3	A	24	0	9	2	0
3	B	24	0	9	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	10	0	0	0	0
5	B	5	0	0	0	0
6	A	166	0	0	4	0
6	B	158	0	0	3	0
All	All	4615	0	4179	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:409[B]:ARG:HD2	6:B:2112:HOH:O	1.92	0.68
1:A:299:ALA:HB3	1:A:300:PRO:HD3	1.78	0.66
1:A:419:LYS:HE2	6:A:2139:HOH:O	1.98	0.63
1:A:423:LEU:HD21	1:A:429:LEU:HG	1.79	0.63
1:A:203:THR:HB	1:A:206:LEU:HD12	1.82	0.62
1:A:366:ASP:OD2	1:A:408:HIS:HD2	1.83	0.62
1:A:341:SER:HB2	1:B:345:ASP:HA	1.83	0.58
1:A:425:ASP:CG	1:A:425:ASP:O	2.42	0.57
1:B:358:VAL:O	1:B:361[A]:GLU:HG3	2.05	0.56
1:A:419:LYS:HE3	3:A:1463:I3P:O43	2.06	0.56
1:A:339:ASP:OD1	1:A:341:SER:HB3	2.08	0.54
1:B:274:GLU:HB3	1:B:313:PRO:HG3	1.91	0.51
1:A:213:GLU:OE1	1:A:213:GLU:HA	2.11	0.50
1:A:364:GLN:NE2	6:A:2102:HOH:O	2.43	0.50
1:A:419:LYS:CE	3:A:1463:I3P:O43	2.62	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:296:ASP:HB3	1:A:299:ALA:HB2	1.96	0.47
1:A:213:GLU:N	1:A:214:PRO:CD	2.77	0.47
1:A:338:ALA:C	1:A:340:GLY:N	2.69	0.46
1:B:332:ILE:HD13	1:B:358:VAL:HG11	1.97	0.46
1:A:353:GLU:OE2	6:A:2093:HOH:O	2.21	0.46
1:A:339:ASP:N	1:B:294:ALA:HB1	2.31	0.46
1:B:361[B]:GLU:OE1	6:B:2089:HOH:O	2.20	0.45
1:A:338:ALA:HB3	1:B:294:ALA:CB	2.48	0.44
1:B:289:TYR:HB2	1:B:309:ALA:HA	1.99	0.44
1:B:461:ARG:NH1	6:B:2153:HOH:O	2.50	0.43
1:A:339:ASP:N	1:B:294:ALA:CB	2.81	0.43
1:A:390:PHE:CE2	1:A:431:HIS:CD2	3.07	0.42
1:A:350[B]:ARG:NH1	6:A:2085:HOH:O	2.44	0.42
1:B:213:GLU:N	1:B:214:PRO:CD	2.82	0.42
1:A:293:LEU:HD12	1:A:297:PRO:HA	2.02	0.41
1:A:312:LYS:O	1:A:316:MET:HG2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	266/265 (100%)	261 (98%)	4 (2%)	1 (0%)	34	22
1	B	245/265 (92%)	237 (97%)	7 (3%)	1 (0%)	34	22
All	All	511/530 (96%)	498 (98%)	11 (2%)	2 (0%)	34	22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	339	ASP
1	B	425	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/229 (101%)	225 (97%)	7 (3%)	41	30
1	B	218/229 (95%)	207 (95%)	11 (5%)	24	11
All	All	450/458 (98%)	432 (96%)	18 (4%)	31	19

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	197	SER
1	A	212	SER
1	A	286	LYS
1	A	339	ASP
1	A	391	ARG
1	A	423	LEU
1	A	425	ASP
1	B	203	THR
1	B	241	ASP
1	B	243	GLU
1	B	244	SER
1	B	262	ASP
1	B	291	LYS
1	B	314	ARG
1	B	316	MET
1	B	352	ARG
1	B	392	ARG
1	B	433	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	247	GLN
1	A	408	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	I3P	A	1463	-	24,24,24	1.50	5 (20%)	36,39,39	1.27	5 (13%)
3	I3P	B	1463	-	24,24,24	1.62	5 (20%)	36,39,39	1.65	9 (25%)
5	SO4	B	1465	-	4,4,4	0.17	0	6,6,6	0.32	0
5	SO4	A	1466	-	4,4,4	0.20	0	6,6,6	0.45	0
2	ANP	A	1462	4	29,33,33	2.69	14 (48%)	31,52,52	2.26	11 (35%)
2	ANP	B	1462	4	29,33,33	2.80	15 (51%)	31,52,52	2.29	12 (38%)
5	SO4	A	1465	-	4,4,4	0.20	0	6,6,6	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	I3P	A	1463	-	-	2/15/39/39	0/1/1/1
3	I3P	B	1463	-	-	3/15/39/39	0/1/1/1

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ANP	A	1462	4	-	4/14/38/38	0/3/3/3
2	ANP	B	1462	4	-	3/14/38/38	0/3/3/3

All (39) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1462	ANP	C5'-C4'	-5.69	1.33	1.51
2	A	1462	ANP	C5'-C4'	-5.49	1.34	1.51
2	A	1462	ANP	C4-N3	4.92	1.42	1.35
2	B	1462	ANP	O2'-C2'	4.83	1.54	1.43
2	B	1462	ANP	PG-O3G	-4.49	1.44	1.56
2	A	1462	ANP	PG-O3G	-4.48	1.44	1.56
2	A	1462	ANP	O3'-C3'	4.29	1.53	1.43
2	B	1462	ANP	C4-N3	4.26	1.41	1.35
2	B	1462	ANP	O3'-C3'	4.23	1.52	1.43
2	A	1462	ANP	O2'-C2'	4.14	1.52	1.43
2	B	1462	ANP	C2'-C1'	-3.90	1.47	1.53
2	B	1462	ANP	C2-N1	3.87	1.41	1.33
2	A	1462	ANP	C8-N7	-3.76	1.28	1.34
2	A	1462	ANP	C2-N1	3.71	1.40	1.33
2	B	1462	ANP	C8-N7	-3.60	1.28	1.34
2	B	1462	ANP	PB-O2B	-3.54	1.47	1.56
3	A	1463	I3P	P1-O1	3.52	1.66	1.59
3	B	1463	I3P	P1-O1	3.47	1.65	1.59
2	A	1462	ANP	PB-O2B	-3.36	1.47	1.56
2	A	1462	ANP	C2'-C1'	-3.08	1.49	1.53
3	A	1463	I3P	P1-O12	-3.00	1.43	1.54
2	B	1462	ANP	C3'-C4'	2.90	1.60	1.53
2	A	1462	ANP	C2'-C3'	2.73	1.60	1.53
3	A	1463	I3P	P1-O11	-2.71	1.41	1.50
3	B	1463	I3P	P4-O41	2.71	1.59	1.50
2	A	1462	ANP	C3'-C4'	2.58	1.59	1.53
3	B	1463	I3P	P4-O42	-2.53	1.45	1.54
3	B	1463	I3P	P1-O11	-2.41	1.42	1.50
2	B	1462	ANP	C2'-C3'	2.38	1.59	1.53
3	B	1463	I3P	P1-O12	-2.38	1.45	1.54
3	A	1463	I3P	P4-O41	2.36	1.58	1.50
2	A	1462	ANP	O5'-C5'	2.34	1.53	1.44
2	B	1462	ANP	PB-N3B	-2.33	1.57	1.63
2	A	1462	ANP	PB-N3B	-2.27	1.57	1.63
2	B	1462	ANP	O4'-C4'	2.21	1.49	1.45
2	A	1462	ANP	PB-O1B	2.19	1.49	1.46

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1462	ANP	O5'-C5'	2.16	1.53	1.44
3	A	1463	I3P	P4-O42	-2.09	1.46	1.54
2	B	1462	ANP	PB-O1B	2.00	1.49	1.46

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1462	ANP	C4-C5-N7	5.05	114.66	109.40
2	A	1462	ANP	C4-C5-N7	4.93	114.54	109.40
2	A	1462	ANP	C5-C6-N6	4.88	127.77	120.35
3	B	1463	I3P	C6-C1-C2	4.67	117.58	110.85
2	B	1462	ANP	C5-C6-N6	4.66	127.43	120.35
2	B	1462	ANP	O3A-PB-N3B	4.36	118.69	106.59
2	A	1462	ANP	O3A-PB-N3B	4.03	117.77	106.59
3	A	1463	I3P	C6-C1-C2	3.66	116.12	110.85
2	B	1462	ANP	O3G-PG-O1G	-3.60	104.39	113.45
2	A	1462	ANP	O2B-PB-O1B	3.56	117.38	109.92
2	A	1462	ANP	O3'-C3'-C4'	3.54	121.27	111.05
2	B	1462	ANP	O3'-C3'-C4'	3.49	121.13	111.05
2	A	1462	ANP	O3G-PG-O2G	3.27	116.34	107.64
3	B	1463	I3P	O1-C1-C6	-3.21	101.20	108.66
2	A	1462	ANP	O3G-PG-O1G	-3.02	105.86	113.45
2	B	1462	ANP	O2B-PB-O1B	2.93	116.07	109.92
2	B	1462	ANP	O3G-PG-O2G	2.84	115.21	107.64
3	B	1463	I3P	O43-P4-O42	2.79	118.31	107.64
2	B	1462	ANP	O1B-PB-N3B	-2.70	107.80	111.77
2	A	1462	ANP	O4'-C4'-C5'	-2.64	100.68	109.37
2	A	1462	ANP	C1'-N9-C4	2.59	131.19	126.64
2	A	1462	ANP	C2'-C3'-C4'	-2.56	97.67	102.64
3	B	1463	I3P	O1-C1-C2	2.48	114.43	108.66
2	A	1462	ANP	O3'-C3'-C2'	2.43	119.68	111.82
3	B	1463	I3P	O13-P1-O1	-2.40	95.22	105.99
2	B	1462	ANP	O2B-PB-O3A	-2.39	96.65	104.64
3	A	1463	I3P	O1-C1-C6	-2.38	103.14	108.66
2	B	1462	ANP	C2'-C3'-C4'	-2.36	98.05	102.64
3	A	1463	I3P	O3-C3-C4	2.29	116.02	109.94
3	B	1463	I3P	O53-P5-O52	-2.21	99.20	107.64
3	B	1463	I3P	O6-C6-C5	2.18	115.72	109.94
3	B	1463	I3P	O2-C2-C1	2.16	115.67	109.94
2	B	1462	ANP	C1'-N9-C4	2.16	130.44	126.64
3	A	1463	I3P	O6-C6-C1	2.11	115.54	109.94
3	A	1463	I3P	O1-C1-C2	2.05	113.44	108.66

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1462	ANP	O2'-C2'-C3'	-2.05	105.19	111.82
3	B	1463	I3P	O3-C3-C4	2.03	115.31	109.94

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1463	I3P	C1-O1-P1-O12
3	B	1463	I3P	C2-C1-O1-P1
2	B	1462	ANP	PG-N3B-PB-O1B
2	B	1462	ANP	C5'-O5'-PA-O1A
2	A	1462	ANP	PA-O3A-PB-O1B
2	A	1462	ANP	C5'-O5'-PA-O1A
3	B	1463	I3P	C1-O1-P1-O12
2	A	1462	ANP	PB-O3A-PA-O2A
3	A	1463	I3P	C2-C1-O1-P1
3	B	1463	I3P	C6-C1-O1-P1
2	B	1462	ANP	PG-N3B-PB-O3A
2	A	1462	ANP	PG-N3B-PB-O3A

There are no ring outliers.


1 monomer is involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1463	I3P	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	265/265 (100%)	0.66	26 (9%) <b>7</b>   <b>12</b>	6, 17, 42, 50	0
1	B	249/265 (93%)	1.13	49 (19%) <b>1</b>   <b>1</b>	5, 18, 53, 66	0
All	All	514/530 (96%)	0.89	75 (14%) <b>2</b>   <b>3</b>	5, 17, 45, 66	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	338	ALA	15.4
1	B	276	THR	8.1
1	B	284	LEU	7.4
1	B	286	LYS	6.6
1	B	311	THR	6.1
1	B	289	TYR	5.6
1	B	313	PRO	5.2
1	B	202	GLY	5.1
1	B	243	GLU	4.8
1	B	338	ALA	4.8
1	B	285	ARG	4.8
1	A	294	ALA	4.7
1	B	294	ALA	4.6
1	B	287	ASP	4.5
1	A	295	VAL	4.4
1	B	293	LEU	4.4
1	B	239	GLU	4.4
1	B	310	VAL	4.3
1	A	203	THR	4.3
1	B	290	LYS	4.1
1	B	283	LYS	4.0
1	B	309	ALA	4.0
1	A	339	ASP	3.9
1	B	242	GLY	3.9

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	273	GLU	3.8
1	B	197	SER	3.7
1	B	277	LYS	3.6
1	B	198	PHE	3.6
1	B	241	ASP	3.6
1	A	204	SER	3.6
1	B	288	MET	3.5
1	A	284	LEU	3.5
1	A	425	ASP	3.5
1	B	295	VAL	3.4
1	A	241	ASP	3.4
1	A	281	ARG	3.4
1	B	244	SER	3.3
1	B	315	TYR	3.3
1	B	240	ARG	3.1
1	A	299	ALA	3.1
1	B	312	LYS	3.0
1	B	425	ASP	3.0
1	A	280	GLU	3.0
1	B	377	LEU	2.9
1	B	210	ARG	2.9
1	A	198	PHE	2.9
1	B	297	PRO	2.7
1	B	200	ALA	2.7
1	B	245	TYR	2.6
1	B	452	ILE	2.6
1	A	210	ARG	2.5
1	A	277	LYS	2.5
1	B	446	LEU	2.5
1	A	197	SER	2.5
1	B	314	ARG	2.5
1	A	243	GLU	2.4
1	B	203	THR	2.4
1	B	201	ALA	2.4
1	A	199	LYS	2.3
1	B	238	VAL	2.3
1	A	292	MET	2.2
1	A	434	PRO	2.2
1	A	290	LYS	2.2
1	B	445	LEU	2.2
1	A	426	GLY	2.2
1	B	339	ASP	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	298	GLU	2.1
1	B	199	LYS	2.1
1	B	204	SER	2.1
1	B	291	LYS	2.1
1	A	240	ARG	2.1
1	B	328	LEU	2.1
1	B	412	VAL	2.0
1	A	202	GLY	2.0
1	A	201	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	MN	B	1464	1/1	0.76	0.28	57,57,57,57	1
2	ANP	B	1462	31/31	0.84	0.24	37,48,78,78	0
2	ANP	A	1462	31/31	0.84	0.18	32,42,75,76	0
4	MN	A	1464	1/1	0.85	0.23	64,64,64,64	1
3	I3P	B	1463	24/24	0.86	0.33	62,67,74,75	0
5	SO4	B	1465	5/5	0.89	0.24	37,41,42,43	5
3	I3P	A	1463	24/24	0.93	0.12	44,50,53,56	0
5	SO4	A	1466	5/5	0.93	0.15	36,36,38,38	5
5	SO4	A	1465	5/5	0.97	0.19	53,54,55,55	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around ANP B 1462:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around ANP A 1462:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around I3P B 1463:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.