



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 10, 2023 – 07:37 pm GMT

PDB ID : 2W3S
Title : Crystal Structure of Xanthine Dehydrogenase (desulfo form) from Rhodobacter capsulatus in Complex with Xanthine
Authors : Dietzel, U.; Kuper, J.; Leimkuhler, S.; Kisker, C.
Deposited on : 2008-11-14
Resolution : 2.60 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	462	 6% 62% 31% . .
1	C	462	 2% 80% 16% . .
1	E	462	 5% 66% 28% . .
1	G	462	 8% 60% 33% . .
2	B	777	 71% 23% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	777	 73% 23% ..
2	F	777	 70% 26% ..
2	H	777	 % 71% 24% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MTE	B	1778	X	-	-	-
5	MTE	D	1778	X	-	-	-
5	MTE	F	1778	X	-	-	-
5	MTE	H	1778	X	-	-	-
8	MOM	B	1781	-	-	X	-
8	MOM	D	1781	-	-	X	-
8	MOM	F	1781	-	-	X	-
8	MOM	H	1781	-	-	X	-

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 36858 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called XANTHINE DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	450	3376	2115	608	628	25	0	0	0
1	C	450	3376	2115	608	628	25	0	0	0
1	E	450	3376	2115	608	628	25	0	0	0
1	G	450	3376	2115	608	628	25	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	26	TRP	LEU	conflict	UNP O54050
C	26	TRP	LEU	conflict	UNP O54050
E	26	TRP	LEU	conflict	UNP O54050
G	26	TRP	LEU	conflict	UNP O54050


- Molecule 2 is a protein called XANTHINE DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	760	5717	3581	1056	1054	26	0	0	0
2	D	760	5717	3581	1056	1054	26	0	0	0
2	F	760	5717	3581	1056	1054	26	0	0	0
2	H	760	5717	3581	1056	1054	26	0	0	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	772	ARG	GLY	conflict	UNP O54051
D	772	ARG	GLY	conflict	UNP O54051
F	772	ARG	GLY	conflict	UNP O54051
H	772	ARG	GLY	conflict	UNP O54051

- Molecule 3 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe₂S₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	Fe	S	0	0
			4	2	2		
3	A	1	Total	Fe	S	0	0
			4	2	2		
3	C	1	Total	Fe	S	0	0
			4	2	2		
3	C	1	Total	Fe	S	0	0
			4	2	2		
3	E	1	Total	Fe	S	0	0
			4	2	2		
3	E	1	Total	Fe	S	0	0
			4	2	2		
3	G	1	Total	Fe	S	0	0
			4	2	2		
3	G	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
4	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
4	E	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
4	G	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 5 is PHOSPHONIC ACIDMONO-(2-AMINO-5,6-DIMERCAPTO-4-OXO-3,7,8A, 9,10,10A-HEXAHYDRO-4H-8-OXA-1,3,9,10-TETRAAZA-ANTHRACEN-7-YLMETHYL) ESTER (three-letter code: MTE) (formula: $C_{10}H_{14}N_5O_6PS_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
5	B	1	Total	C	N	O	P	S	0	0
			24	10	5	6	1	2		
5	D	1	Total	C	N	O	P	S	0	0
			24	10	5	6	1	2		
5	F	1	Total	C	N	O	P	S	0	0
			24	10	5	6	1	2		
5	H	1	Total	C	N	O	P	S	0	0
			24	10	5	6	1	2		

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Ca	0	0
			1	1		
6	D	1	Total	Ca	0	0
			1	1		
6	F	1	Total	Ca	0	0
			1	1		
6	H	1	Total	Ca	0	0
			1	1		

- Molecule 7 is XANTHINE (three-letter code: XAN) (formula: C₅H₄N₄O₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	B	1	Total	C	N	O	0	1
			22	10	8	4		
7	D	1	Total	C	N	O	0	1
			22	10	8	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	F	1	Total	C	N	O	0	1
			22	10	8	4		
7	H	1	Total	C	N	O	0	1
			22	10	8	4		

- Molecule 8 is HYDROXY(DIOXO)MOLYBDENUM (three-letter code: MOM) (formula: HMoO_3).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	Mo	O	0	0
			4	1	3		
8	D	1	Total	Mo	O	0	0
			4	1	3		
8	F	1	Total	Mo	O	0	0
			4	1	3		
8	H	1	Total	Mo	O	0	0
			4	1	3		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	3	Total	O	0	0
			3	3		
9	B	9	Total	O	0	0
			9	9		
9	C	4	Total	O	0	0
			4	4		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	D	7	Total O 7 7	0	0
9	E	2	Total O 2 2	0	0
9	F	6	Total O 6 6	0	0
9	H	7	Total O 7 7	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: XANTHINE DEHYDROGENASE




• Molecule 1: XANTHINE DEHYDROGENASE




• Molecule 1: XANTHINE DEHYDROGENASE






• Molecule 1: XANTHINE DEHYDROGENASE




• Molecule 2: XANTHINE DEHYDROGENASE







• Molecule 2: XANTHINE DEHYDROGENASE




• Molecule 2: XANTHINE DEHYDROGENASE





● Molecule 2: XANTHINE DEHYDROGENASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	92.83Å 140.56Å 158.17Å 109.60° 105.89° 101.18°	Depositor
Resolution (Å)	50.12 – 2.60 50.01 – 2.60	Depositor EDS
% Data completeness (in resolution range)	97.7 (50.12-2.60) 97.7 (50.01-2.60)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 2.61Å)	Xtrriage
Refinement program	REFMAC 5.5.0055	Depositor
R, R_{free}	0.232 , 0.284 0.234 , 0.282	Depositor DCC
R_{free} test set	10343 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	41.5	Xtrriage
Anisotropy	0.200	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 55.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	36858	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, FES, CA, MTE, MOM, XAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/3439	0.67	0/4659
1	C	0.50	0/3439	0.64	0/4659
1	E	0.51	0/3439	0.66	0/4659
1	G	0.54	1/3439 (0.0%)	0.69	0/4659
2	B	0.56	0/5845	0.70	0/7942
2	D	0.57	1/5845 (0.0%)	0.70	0/7942
2	F	0.58	0/5845	0.71	0/7942
2	H	0.57	0/5845	0.69	0/7942
All	All	0.55	2/37136 (0.0%)	0.69	0/50404

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	103	CYS	CB-SG	-5.70	1.72	1.81
2	D	750	CYS	CB-SG	-5.15	1.73	1.81

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3376	0	3367	127	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3376	0	3367	52	0
1	E	3376	0	3367	94	0
1	G	3376	0	3367	123	0
2	B	5717	0	5631	154	0
2	D	5717	0	5631	125	0
2	F	5717	0	5631	143	0
2	H	5717	0	5630	141	0
3	A	8	0	0	1	0
3	C	8	0	0	0	0
3	E	8	0	0	0	0
3	G	8	0	0	1	0
4	A	53	0	31	6	0
4	C	53	0	31	3	0
4	E	53	0	31	3	0
4	G	53	0	31	4	0
5	B	24	0	8	0	0
5	D	24	0	9	1	0
5	F	24	0	8	2	0
5	H	24	0	8	3	0
6	B	1	0	0	0	0
6	D	1	0	0	0	0
6	F	1	0	0	0	0
6	H	1	0	0	0	0
7	B	22	0	8	0	0
7	D	22	0	8	3	0
7	F	22	0	8	0	0
7	H	22	0	8	0	0
8	B	4	0	0	2	0
8	D	4	0	0	5	0
8	F	4	0	0	2	0
8	H	4	0	0	2	0
9	A	3	0	0	0	0
9	B	9	0	0	0	0
9	C	4	0	0	0	0
9	D	7	0	0	0	0
9	E	2	0	0	0	0
9	F	6	0	0	0	0
9	H	7	0	0	0	0
All	All	36858	0	36180	896	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 896 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:247:ARG:HG2	2:F:247:ARG:HH11	1.11	1.07
1:G:240:GLY:HA2	1:G:343:LYS:HG2	1.34	1.04
2:F:221:MET:HE1	2:F:224:MET:HG3	1.40	1.03
1:A:240:GLY:HA2	1:A:343:LYS:HG2	1.41	1.03
1:G:361:ILE:HD11	1:G:429:ARG:NH2	1.75	1.01

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	446/462 (96%)	402 (90%)	38 (8%)	6 (1%)	12 24
1	C	446/462 (96%)	422 (95%)	20 (4%)	4 (1%)	17 35
1	E	446/462 (96%)	389 (87%)	46 (10%)	11 (2%)	5 9
1	G	446/462 (96%)	383 (86%)	56 (13%)	7 (2%)	9 19
2	B	756/777 (97%)	706 (93%)	39 (5%)	11 (2%)	10 21
2	D	756/777 (97%)	713 (94%)	36 (5%)	7 (1%)	17 35
2	F	756/777 (97%)	707 (94%)	41 (5%)	8 (1%)	14 30
2	H	756/777 (97%)	689 (91%)	58 (8%)	9 (1%)	13 27
All	All	4808/4956 (97%)	4411 (92%)	334 (7%)	63 (1%)	12 24

5 of 63 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	165	THR
1	A	345	ALA
2	B	187	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	458	SER
2	D	458	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	339/347 (98%)	311 (92%)	28 (8%)	11	22
1	C	339/347 (98%)	311 (92%)	28 (8%)	11	22
1	E	339/347 (98%)	306 (90%)	33 (10%)	8	15
1	G	339/347 (98%)	304 (90%)	35 (10%)	7	13
2	B	571/584 (98%)	524 (92%)	47 (8%)	11	22
2	D	571/584 (98%)	530 (93%)	41 (7%)	14	29
2	F	571/584 (98%)	523 (92%)	48 (8%)	11	21
2	H	571/584 (98%)	529 (93%)	42 (7%)	13	28
All	All	3640/3724 (98%)	3338 (92%)	302 (8%)	11	22

5 of 302 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	79	GLU
2	H	424	SER
1	G	231	LEU
2	H	10	ASP
2	H	741	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 47 such sidechains are listed below:

Mol	Chain	Res	Type
2	F	204	HIS
2	F	744	HIS
2	F	236	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	F	463	HIS
1	G	196	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 32 ligands modelled in this entry, 4 are monoatomic - leaving 28 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FES	A	1463	1	0,4,4	-	-	-		
3	FES	E	1463	1	0,4,4	-	-	-		
7	XAN	D	1780[A]	-	8,12,12	1.87	2 (25%)	4,17,17	5.43	3 (75%)
7	XAN	F	1780[B]	-	8,12,12	2.06	2 (25%)	4,17,17	5.28	4 (100%)
4	FAD	C	1465	-	53,58,58	1.58	8 (15%)	68,89,89	1.50	17 (25%)
4	FAD	E	1465	-	53,58,58	1.52	5 (9%)	68,89,89	1.50	15 (22%)
8	MOM	B	1781	5	0,3,3	-	-	-		
7	XAN	B	1780[A]	-	8,12,12	1.96	2 (25%)	4,17,17	5.13	4 (100%)
5	MTE	B	1778	8	21,26,26	3.89	5 (23%)	21,40,40	2.73	9 (42%)
3	FES	C	1464	1	0,4,4	-	-	-		

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	XAN	H	1780[A]	-	8,12,12	1.94	2 (25%)	4,17,17	5.29	3 (75%)
3	FES	A	1464	1	0,4,4	-	-	-		
3	FES	C	1463	1	0,4,4	-	-	-		
5	MTE	H	1778	8	21,26,26	3.68	6 (28%)	21,40,40	2.34	7 (33%)
7	XAN	F	1780[A]	-	8,12,12	2.03	2 (25%)	4,17,17	5.20	4 (100%)
7	XAN	D	1780[B]	-	8,12,12	2.05	2 (25%)	4,17,17	5.22	4 (100%)
4	FAD	G	1465	-	53,58,58	1.50	4 (7%)	68,89,89	1.39	11 (16%)
3	FES	G	1464	1	0,4,4	-	-	-		
3	FES	E	1464	1	0,4,4	-	-	-		
7	XAN	B	1780[B]	-	8,12,12	2.06	2 (25%)	4,17,17	5.27	4 (100%)
5	MTE	D	1778	8	21,26,26	3.96	6 (28%)	21,40,40	2.76	8 (38%)
7	XAN	H	1780[B]	-	8,12,12	2.09	2 (25%)	4,17,17	5.19	4 (100%)
5	MTE	F	1778	8	21,26,26	4.07	7 (33%)	21,40,40	2.55	9 (42%)
8	MOM	F	1781	5	0,3,3	-	-	-		
4	FAD	A	1465	-	53,58,58	1.52	5 (9%)	68,89,89	1.44	11 (16%)
3	FES	G	1463	1	0,4,4	-	-	-		
8	MOM	D	1781	5	0,3,3	-	-	-		
8	MOM	H	1781	5	0,3,3	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FES	A	1463	1	-	-	0/1/1/1
3	FES	E	1463	1	-	-	0/1/1/1
7	XAN	D	1780[A]	-	-	-	0/2/2/2
7	XAN	F	1780[B]	-	-	-	0/2/2/2
4	FAD	C	1465	-	-	9/30/50/50	0/6/6/6
4	FAD	E	1465	-	-	15/30/50/50	0/6/6/6
7	XAN	B	1780[A]	-	-	-	0/2/2/2
5	MTE	B	1778	8	1/1/6/8	1/6/34/34	0/3/3/3
3	FES	C	1464	1	-	-	0/1/1/1
7	XAN	H	1780[A]	-	-	-	0/2/2/2
3	FES	A	1464	1	-	-	0/1/1/1
3	FES	C	1463	1	-	-	0/1/1/1
5	MTE	H	1778	8	1/1/6/8	2/6/34/34	0/3/3/3
7	XAN	F	1780[A]	-	-	-	0/2/2/2
7	XAN	D	1780[B]	-	-	-	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FAD	G	1465	-	-	9/30/50/50	0/6/6/6
3	FES	E	1464	1	-	-	0/1/1/1
3	FES	G	1464	1	-	-	0/1/1/1
7	XAN	B	1780[B]	-	-	-	0/2/2/2
5	MTE	D	1778	8	1/1/6/8	2/6/34/34	0/3/3/3
7	XAN	H	1780[B]	-	-	-	0/2/2/2
5	MTE	F	1778	8	1/1/6/8	1/6/34/34	0/3/3/3
4	FAD	A	1465	-	-	11/30/50/50	0/6/6/6
3	FES	G	1463	1	-	-	0/1/1/1

The worst 5 of 62 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	F	1778	MTE	C4'-C3'	-14.73	1.32	1.52
5	B	1778	MTE	C4'-C3'	-14.06	1.33	1.52
5	D	1778	MTE	C4'-C3'	-13.89	1.33	1.52
5	H	1778	MTE	C4'-C3'	-12.29	1.35	1.52
5	D	1778	MTE	C6-N5	-8.16	1.33	1.45

The worst 5 of 117 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	D	1780[A]	XAN	C2-N1-C6	8.91	122.67	115.14
7	H	1780[A]	XAN	C2-N1-C6	8.62	122.42	115.14
7	B	1780[B]	XAN	C2-N1-C6	8.44	122.26	115.14
7	F	1780[B]	XAN	C2-N1-C6	8.42	122.25	115.14
7	F	1780[A]	XAN	C2-N1-C6	8.35	122.19	115.14

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	B	1778	MTE	C3'
5	D	1778	MTE	C3'
5	F	1778	MTE	C3'
5	H	1778	MTE	C3'

5 of 50 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1465	FAD	C2'-C1'-N10-C10
4	A	1465	FAD	N10-C1'-C2'-O2'

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	1465	FAD	N10-C1'-C2'-C3'
4	A	1465	FAD	C2'-C3'-C4'-O4'
4	A	1465	FAD	C2'-C3'-C4'-C5'


There are no ring outliers.



15 monomers are involved in 35 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1463	FES	1	0
7	D	1780[A]	XAN	2	0
4	C	1465	FAD	3	0
4	E	1465	FAD	3	0
8	B	1781	MOM	2	0
5	H	1778	MTE	3	0
7	D	1780[B]	XAN	1	0
4	G	1465	FAD	4	0
5	D	1778	MTE	1	0
5	F	1778	MTE	2	0
8	F	1781	MOM	2	0
4	A	1465	FAD	6	0
3	G	1463	FES	1	0
8	D	1781	MOM	5	0
8	H	1781	MOM	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	450/462 (97%)	0.22	30 (6%) 17 13	25, 49, 66, 70	0
1	C	450/462 (97%)	-0.04	7 (1%) 72 68	22, 47, 66, 70	0
1	E	450/462 (97%)	0.09	23 (5%) 28 22	25, 48, 66, 71	0
1	G	450/462 (97%)	0.25	39 (8%) 10 7	26, 49, 66, 70	0
2	B	760/777 (97%)	-0.27	3 (0%) 92 91	21, 35, 50, 59	0
2	D	760/777 (97%)	-0.28	1 (0%) 95 95	20, 34, 49, 59	0
2	F	760/777 (97%)	-0.36	1 (0%) 95 95	21, 34, 49, 59	0
2	H	760/777 (97%)	-0.35	8 (1%) 80 78	22, 35, 50, 59	0
All	All	4840/4956 (97%)	-0.15	112 (2%) 60 54	20, 39, 62, 71	0

The worst 5 of 112 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	374	GLY	4.4
1	A	411	ILE	4.2
1	A	376	LYS	4.0
1	E	166	LEU	4.0
1	A	377	ILE	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	XAN	F	1780[A]	11/11	0.74	0.27	62,62,62,62	11
7	XAN	F	1780[B]	11/11	0.74	0.27	61,62,62,62	11
7	XAN	H	1780[A]	11/11	0.78	0.30	62,62,62,62	11
7	XAN	H	1780[B]	11/11	0.78	0.30	61,62,62,62	11
7	XAN	B	1780[A]	11/11	0.79	0.26	62,62,62,62	11
7	XAN	B	1780[B]	11/11	0.79	0.26	61,62,62,62	11
7	XAN	D	1780[A]	11/11	0.84	0.20	62,62,62,62	11
7	XAN	D	1780[B]	11/11	0.84	0.20	61,62,62,62	11
4	FAD	G	1465	53/53	0.91	0.17	41,50,63,64	0
5	MTE	H	1778	24/24	0.92	0.28	33,38,41,43	24
4	FAD	E	1465	53/53	0.93	0.15	35,45,53,53	0
4	FAD	A	1465	53/53	0.93	0.15	38,48,56,56	0
5	MTE	B	1778	24/24	0.93	0.34	27,36,42,43	24
5	MTE	F	1778	24/24	0.94	0.27	22,29,34,36	24
5	MTE	D	1778	24/24	0.94	0.27	9,25,32,33	24
4	FAD	C	1465	53/53	0.96	0.14	16,27,35,37	0
6	CA	F	1779	1/1	0.97	0.26	56,56,56,56	0
6	CA	H	1779	1/1	0.97	0.22	58,58,58,58	0
3	FES	G	1464	4/4	0.97	0.19	39,40,42,42	0
3	FES	E	1463	4/4	0.97	0.15	26,27,30,31	0
3	FES	E	1464	4/4	0.97	0.17	38,38,40,42	0
3	FES	G	1463	4/4	0.98	0.14	29,32,32,33	0
6	CA	D	1779	1/1	0.98	0.18	49,49,49,49	0
3	FES	A	1463	4/4	0.98	0.19	25,28,29,30	0
3	FES	C	1463	4/4	0.98	0.16	19,19,21,22	0
8	MOM	B	1781	4/4	0.98	0.12	44,45,45,47	3
8	MOM	H	1781	4/4	0.98	0.12	44,44,44,45	3
3	FES	C	1464	4/4	0.99	0.17	33,35,35,37	0
6	CA	B	1779	1/1	0.99	0.22	48,48,48,48	0
8	MOM	D	1781	4/4	0.99	0.10	34,34,34,37	3
8	MOM	F	1781	4/4	0.99	0.12	35,35,36,37	3
3	FES	A	1464	4/4	0.99	0.18	35,35,36,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around MTE H 1778:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD E 1465:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD A 1465:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MTE B 1778:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MTE F 1778:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MTE D 1778:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.