



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 2, 2024 – 11:08 PM EST

PDB ID : 5W8V
Title : HUMAN HGPRT in complex with [(2-[(guanin-9-yl)methyl]propane-1,3-diyl)bis(oxy)]bis(methylene)diphosphonic acid
Authors : Guddat, L.W.; Keough, D.T.
Deposited on : 2017-06-22
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	215	
1	B	215	
1	C	215	
1	D	215	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7052 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Hypoxanthine-guanine phosphoribosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	206	1586	1021	265	293	7	5	1	0
1	B	204	1596	1026	263	299	8	4	3	0
1	C	205	1610	1033	271	299	7	0	1	0
1	D	204	1623	1044	271	299	9	5	3	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	3	ALA	-	expression tag	UNP P00492
B	3	ALA	-	expression tag	UNP P00492
C	3	ALA	-	expression tag	UNP P00492
D	3	ALA	-	expression tag	UNP P00492

- Molecule 2 is $\{[(2R)-2-[(2\text{-amino-6-oxo-1,6-dihydro-9H-purin-9-yl)methyl]-3-(2\text{-phosphonoethoxy})\text{propoxy}]methyl\}$ phosphonic acid (three-letter code: 9YP) (formula: $C_{12}H_{21}N_5O_9P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			28	12	5	9	2		
2	B	1	Total	C	N	O	P	0	0
			28	12	5	9	2		
2	C	1	Total	C	N	O	P	0	0
			28	12	5	9	2		
2	D	1	Total	C	N	O	P	0	0
			28	12	5	9	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		
3	C	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	126	Total	O	0	0
			126	126		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	114	Total 114	O 114	0	0
4	C	135	Total 135	O 135	0	0
4	D	146	Total 146	O 146	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hypoxanthine-guanine phosphoribosyltransferase




- Molecule 1: Hypoxanthine-guanine phosphoribosyltransferase



- Molecule 1: Hypoxanthine-guanine phosphoribosyltransferase



- Molecule 1: Hypoxanthine-guanine phosphoribosyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.67Å 128.40Å 65.33Å 90.00° 102.98° 90.00°	Depositor
Resolution (Å)	47.03 – 2.35 47.30 – 2.03	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.03-2.35) 99.3 (47.30-2.03)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 2.03Å)	Xtrriage
Refinement program	PHENIX 1.8_1069	Depositor
R, R_{free}	0.196 , 0.249 0.203 , 0.248	Depositor DCC
R_{free} test set	2000 reflections (3.42%)	wwPDB-VP
Wilson B-factor (Å ²)	15.3	Xtrriage
Anisotropy	0.206	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 53.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.83	EDS
Total number of atoms	7052	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 9YP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/1620	0.41	0/2197
1	B	0.25	0/1632	0.42	0/2211
1	C	0.23	0/1644	0.41	0/2223
1	D	0.23	0/1663	0.42	0/2247
All	All	0.24	0/6559	0.42	0/8878

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1586	0	1551	18	0
1	B	1596	0	1559	17	0
1	C	1610	0	1598	18	0
1	D	1623	0	1636	21	0
2	A	28	0	0	1	0
2	B	28	0	0	1	0
2	C	28	0	0	2	0
2	D	28	0	0	1	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	126	0	0	7	2
4	B	114	0	0	4	0
4	C	135	0	0	5	0
4	D	146	0	0	13	2
All	All	7052	0	6344	70	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25:ASN:OD1	4:A:401:HOH:O	1.91	0.88
1:A:85:ASN:ND2	4:A:403:HOH:O	2.12	0.83
1:C:76:ASP:OD2	4:C:401:HOH:O	2.03	0.77
1:D:125:THR:O	4:D:402:HOH:O	2.04	0.76
1:B:127:LYS:NZ	4:B:402:HOH:O	2.18	0.72
1:D:97:ASP:OD2	4:D:403:HOH:O	2.08	0.71
2:B:301:9YP:OAD	4:B:401:HOH:O	2.08	0.71
1:A:117:GLY:O	4:A:402:HOH:O	2.11	0.67
1:C:32:GLU:OE2	4:C:402:HOH:O	2.12	0.66
1:D:210:THR:O	4:D:404:HOH:O	2.14	0.66
1:A:29:GLU:OE2	4:A:404:HOH:O	2.15	0.65
1:D:47:ARG:NH1	1:D:51:ASP:OD1	2.30	0.64
1:A:92:ILE:O	4:A:405:HOH:O	2.15	0.64
2:C:301:9YP:OAF	4:C:403:HOH:O	2.16	0.63
1:D:174:LYS:NZ	4:D:411:HOH:O	2.31	0.62
1:B:86:ARG:O	1:C:50:ARG:NH1	2.33	0.61
1:D:79:ASP:HA	1:D:82:LYS:HE2	1.82	0.60
1:D:113:ILE:N	4:D:412:HOH:O	2.34	0.59
1:B:100:ARG:HB2	1:B:116:ILE:HG13	1.83	0.59
1:C:61:ILE:HB	1:C:94:MET:HG2	1.83	0.59
1:B:173:TYR:N	4:B:406:HOH:O	2.34	0.59
1:D:216:LYS:NZ	1:D:217:ALA:O	2.36	0.59
1:D:175:PRO:HG2	1:D:178:VAL:HG22	1.85	0.59
1:A:135:ILE:HD11	1:A:165:LYS:HE3	1.86	0.58
1:B:87:ASN:HA	1:C:50:ARG:HH11	1.70	0.57
1:B:136:ILE:HG13	1:B:175:PRO:HG3	1.85	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:301:9YP:OAE	4:D:405:HOH:O	2.18	0.54
1:C:36:ILE:HB	1:C:204:VAL:HB	1.92	0.52
1:C:165:LYS:NZ	1:C:185:LYS:O	2.42	0.51
1:D:174:LYS:O	4:D:406:HOH:O	2.19	0.50
1:D:135:ILE:HD11	1:D:165:LYS:HE3	1.92	0.50
1:B:32:GLU:HG2	1:B:208:SER:HA	1.93	0.50
1:B:151:GLN:NE2	4:B:413:HOH:O	2.42	0.50
1:A:169:ARG:NE	4:A:407:HOH:O	2.43	0.50
1:B:36:ILE:HB	1:B:204:VAL:HB	1.95	0.49
1:A:36:ILE:HB	1:A:204:VAL:HB	1.95	0.49
1:A:137:ASP:HB3	2:A:301:9YP:OAD	2.14	0.48
1:B:50:ARG:NH1	1:C:86:ARG:O	2.46	0.48
1:C:7:VAL:HG22	1:D:20:LEU:HD13	1.95	0.47
1:C:38:HIS:HD2	4:D:428:HOH:O	1.97	0.47
1:A:175:PRO:HG2	1:A:178:VAL:HG22	1.96	0.47
1:B:128:ASN:OD1	1:B:155:LYS:HB3	2.16	0.46
1:C:143:GLN:NE2	4:C:423:HOH:O	2.48	0.46
1:D:4:SER:HB3	4:D:505:HOH:O	2.15	0.46
1:A:65:CYS:HB2	1:A:74:PHE:CD1	2.50	0.46
1:C:125:THR:HG21	4:C:532:HOH:O	2.15	0.46
1:A:60:HIS:HD1	1:A:127:LYS:HD2	1.81	0.45
1:C:137:ASP:HB3	2:C:301:9YP:OAD	2.17	0.44
1:D:143:GLN:NE2	4:D:426:HOH:O	2.51	0.44
1:B:100:ARG:NH2	1:D:97:ASP:OD1	2.37	0.43
1:A:147:SER:O	1:A:151:GLN:HG3	2.19	0.43
1:B:126:GLY:H	1:B:154:PRO:HA	1.83	0.43
1:C:127:LYS:HA	1:C:127:LYS:HD3	1.85	0.42
1:D:91:SER:N	4:D:419:HOH:O	2.42	0.42
1:D:51:ASP:O	1:D:54:LYS:HG2	2.20	0.42
1:D:85:ASN:ND2	4:D:418:HOH:O	2.41	0.42
1:B:54:LYS:HB2	1:B:54:LYS:HE3	1.83	0.42
1:C:125:THR:HG22	1:C:152:TYR:O	2.20	0.42
1:D:47:ARG:NH2	4:D:430:HOH:O	2.53	0.41
1:D:11:ASP:OD1	1:D:183:PRO:HB3	2.20	0.41
1:B:65:CYS:HB2	1:B:74:PHE:CD1	2.55	0.41
1:A:18:LEU:HD23	1:A:34:VAL:HG23	2.03	0.41
1:A:150:ARG:NH2	4:A:424:HOH:O	2.53	0.41
1:D:140:LYS:O	1:D:144:THR:HG22	2.20	0.41
1:A:199:ARG:HD2	1:C:96:VAL:HG22	2.03	0.41
1:C:122:SER:O	1:C:125:THR:HG23	2.21	0.41
1:A:20:LEU:HD13	1:B:7:VAL:HG22	2.03	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:127:LYS:O	1:B:154:PRO:HB2	2.21	0.40
1:A:167:THR:HA	1:A:168:PRO:HD3	1.95	0.40
1:C:160:ALA:HA	1:C:177:PHE:HB2	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:510:HOH:O	4:D:431:HOH:O[2_657]	2.05	0.15
4:A:521:HOH:O	4:D:412:HOH:O[2_657]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	203/215 (94%)	197 (97%)	6 (3%)	0	100	100
1	B	201/215 (94%)	190 (94%)	9 (4%)	2 (1%)	15	14
1	C	202/215 (94%)	199 (98%)	3 (2%)	0	100	100
1	D	203/215 (94%)	199 (98%)	4 (2%)	0	100	100
All	All	809/860 (94%)	785 (97%)	22 (3%)	2 (0%)	47	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	168	PRO
1	B	153	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/189 (88%)	165 (99%)	1 (1%)	86	92
1	B	170/189 (90%)	164 (96%)	6 (4%)	36	45
1	C	173/189 (92%)	167 (96%)	6 (4%)	36	45
1	D	178/189 (94%)	175 (98%)	3 (2%)	60	72
All	All	687/756 (91%)	671 (98%)	16 (2%)	52	61

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	TYR
1	B	71	TYR
1	B	82	LYS
1	B	127	LYS
1	B	157[A]	VAL
1	B	157[B]	VAL
1	B	202	ASN
1	C	13	GLU
1	C	40	LEU
1	C	47	ARG
1	C	71	TYR
1	C	121	LEU
1	C	157	VAL
1	D	71	TYR
1	D	140	LYS
1	D	202	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	9YP	D	301	-	25,29,29	2.33	8 (32%)	28,42,42	1.64	6 (21%)
2	9YP	C	301	-	25,29,29	2.43	10 (40%)	28,42,42	1.60	5 (17%)
2	9YP	B	301	-	25,29,29	2.35	7 (28%)	28,42,42	1.58	6 (21%)
2	9YP	A	301	-	25,29,29	2.47	10 (40%)	28,42,42	1.59	6 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	9YP	D	301	-	-	8/18/19/19	0/2/2/2
2	9YP	C	301	-	-	5/18/19/19	0/2/2/2
2	9YP	B	301	-	-	9/18/19/19	0/2/2/2
2	9YP	A	301	-	-	9/18/19/19	0/2/2/2

All (35) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	301	9YP	OAB-CAV	7.30	1.38	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	301	9YP	OAB-CAV	7.28	1.38	1.23
2	A	301	9YP	OAB-CAV	7.21	1.38	1.23
2	B	301	9YP	OAB-CAV	7.19	1.37	1.23
2	B	301	9YP	PBA-CAL	4.15	1.83	1.78
2	A	301	9YP	CAP-NAZ	-4.01	1.43	1.48
2	D	301	9YP	CAP-NAZ	-3.93	1.43	1.48
2	A	301	9YP	PBA-CAL	3.88	1.82	1.78
2	B	301	9YP	CAP-NAZ	-3.88	1.43	1.48
2	D	301	9YP	CAU-NAA	3.87	1.43	1.34
2	C	301	9YP	CAP-NAZ	-3.86	1.44	1.48
2	A	301	9YP	CAU-NAA	3.84	1.43	1.34
2	B	301	9YP	CAU-NAA	3.80	1.43	1.34
2	C	301	9YP	PBA-CAL	3.80	1.82	1.78
2	D	301	9YP	PBA-CAL	3.73	1.82	1.78
2	C	301	9YP	CAU-NAA	3.70	1.43	1.34
2	A	301	9YP	PBA-OAF	2.93	1.56	1.50
2	B	301	9YP	PBA-OAF	2.89	1.56	1.50
2	C	301	9YP	PBA-OAF	2.87	1.56	1.50
2	C	301	9YP	PBB-OAH	2.81	1.56	1.50
2	D	301	9YP	PBB-OAH	2.77	1.56	1.50
2	A	301	9YP	PBB-OAH	2.74	1.55	1.50
2	A	301	9YP	OAT-CAO	-2.67	1.38	1.42
2	B	301	9YP	OAT-CAO	-2.55	1.38	1.42
2	C	301	9YP	OAT-CAO	-2.49	1.38	1.42
2	D	301	9YP	OAT-CAO	-2.39	1.38	1.42
2	A	301	9YP	PBB-OAD	-2.16	1.50	1.54
2	C	301	9YP	PBB-OAD	-2.13	1.50	1.54
2	A	301	9YP	PBA-OAC	-2.10	1.50	1.54
2	D	301	9YP	PBB-OAD	-2.08	1.50	1.54
2	A	301	9YP	CAX-CAV	-2.06	1.43	1.47
2	D	301	9YP	CAX-CAV	-2.05	1.43	1.47
2	C	301	9YP	CAX-CAV	-2.02	1.43	1.47
2	C	301	9YP	PBA-OAC	-2.02	1.50	1.54
2	B	301	9YP	PBA-OAC	-2.02	1.50	1.54

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	9YP	OAT-CAN-CAY	3.46	119.12	109.63
2	B	301	9YP	CAI-NAQ-CAX	3.35	109.37	102.99
2	A	301	9YP	CAI-NAQ-CAX	3.28	109.24	102.99
2	D	301	9YP	OAT-CAN-CAY	3.24	118.50	109.63

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	301	9YP	CAI-NAQ-CAX	3.22	109.12	102.99
2	C	301	9YP	CAI-NAQ-CAX	3.22	109.12	102.99
2	C	301	9YP	OAS-CAM-CAY	3.21	118.44	109.63
2	D	301	9YP	CAO-OAT-CAN	3.00	119.98	112.84
2	C	301	9YP	CAO-OAT-CAN	2.98	119.95	112.84
2	B	301	9YP	OAS-CAM-CAY	2.92	117.63	109.63
2	A	301	9YP	OAS-CAM-CAY	2.92	117.62	109.63
2	B	301	9YP	OAT-CAN-CAY	2.90	117.56	109.63
2	C	301	9YP	OAT-CAN-CAY	2.82	117.36	109.63
2	B	301	9YP	CAO-OAT-CAN	2.73	119.34	112.84
2	A	301	9YP	OAS-CAK-CAL	2.46	116.26	109.19
2	B	301	9YP	OAS-CAK-CAL	2.43	116.18	109.19
2	D	301	9YP	OAS-CAK-CAL	2.41	116.11	109.19
2	D	301	9YP	CAX-CAV-NAR	2.32	118.05	113.95
2	A	301	9YP	CAX-CAV-NAR	2.31	118.03	113.95
2	B	301	9YP	CAX-CAV-NAR	2.30	118.01	113.95
2	C	301	9YP	CAX-CAV-NAR	2.24	117.91	113.95
2	D	301	9YP	OAS-CAM-CAY	2.19	115.64	109.63
2	A	301	9YP	CAO-OAT-CAN	2.04	117.70	112.84

There are no chirality outliers.

All (31) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	9YP	CAK-CAL-PBA-OAE
2	A	301	9YP	CAK-CAL-PBA-OAF
2	A	301	9YP	CAK-CAL-PBA-OAC
2	A	301	9YP	OAS-CAM-CAY-CAN
2	A	301	9YP	OAS-CAM-CAY-CAP
2	A	301	9YP	OAT-CAN-CAY-CAM
2	A	301	9YP	OAT-CAN-CAY-CAP
2	B	301	9YP	OAS-CAM-CAY-CAN
2	B	301	9YP	OAS-CAM-CAY-CAP
2	C	301	9YP	OAS-CAK-CAL-PBA
2	C	301	9YP	CAL-CAK-OAS-CAM
2	D	301	9YP	OAS-CAK-CAL-PBA
2	B	301	9YP	NAZ-CAP-CAY-CAM
2	B	301	9YP	NAZ-CAP-CAY-CAN
2	D	301	9YP	NAZ-CAP-CAY-CAN
2	C	301	9YP	CAY-CAM-OAS-CAK
2	D	301	9YP	CAY-CAM-OAS-CAK
2	D	301	9YP	CAY-CAN-OAT-CAO

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	C	301	9YP	CAY-CAN-OAT-CAO
2	A	301	9YP	CAL-CAK-OAS-CAM
2	B	301	9YP	OAT-CAN-CAY-CAP
2	D	301	9YP	OAS-CAM-CAY-CAP
2	B	301	9YP	CAL-CAK-OAS-CAM
2	B	301	9YP	PBB-CAO-OAT-CAN
2	A	301	9YP	OAS-CAK-CAL-PBA
2	D	301	9YP	CAL-CAK-OAS-CAM
2	D	301	9YP	CAK-CAL-PBA-OAC
2	B	301	9YP	OAT-CAN-CAY-CAM
2	B	301	9YP	OAS-CAK-CAL-PBA
2	C	301	9YP	PBB-CAO-OAT-CAN
2	D	301	9YP	PBB-CAO-OAT-CAN


There are no ring outliers.


4 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	301	9YP	1	0
2	C	301	9YP	2	0
2	B	301	9YP	1	0
2	A	301	9YP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.




6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 9YP A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 9YP B 301:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 9YP C 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [\(i\)](#)

Unable to reproduce the depositor's R factor - this section is therefore empty.