



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 17, 2023 – 09:20 AM EST

PDB ID : 4WIJ
Title : HUMAN SPLICING FACTOR, CONSTRUCT 1
Authors : lee, M.; bond, c.s.
Deposited on : 2014-09-26
Resolution : 3.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1659 (3.60-3.40)
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	326	76% (green), 14% (yellow), 9% (grey)
1	B	326	74% (green), 17% (yellow), 8% (grey)

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 4978 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Splicing factor, proline- and glutamine-rich.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	298	2473	1532	453	473	15	0	1	0
1	B	301	2505	1548	463	479	15	0	1	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	273	GLY	-	expression tag	UNP P23246
A	274	ALA	-	expression tag	UNP P23246
A	275	MET	-	expression tag	UNP P23246
B	273	GLY	-	expression tag	UNP P23246
B	274	ALA	-	expression tag	UNP P23246
B	275	MET	-	expression tag	UNP P23246

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Splicing factor, proline- and glutamine-rich

Chain A: 



- Molecule 1: Splicing factor, proline- and glutamine-rich

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	66.58Å 66.58Å 398.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.08 – 3.49 47.08 – 3.49	Depositor EDS
% Data completeness (in resolution range)	99.4 (47.08-3.49) 99.7 (47.08-3.49)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.87 (at 3.48Å)	Xtrriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.272 , 0.335 0.315 , 0.385	Depositor DCC
R_{free} test set	605 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	103.8	Xtrriage
Anisotropy	0.667	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 67.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.39$, $\langle L^2 \rangle = 0.21$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	4978	wwPDB-VP
Average B, all atoms (Å ²)	127.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	1/2516 (0.0%)	0.62	2/3366 (0.1%)
1	B	0.35	0/2548	0.56	1/3407 (0.0%)
All	All	0.36	1/5064 (0.0%)	0.59	3/6773 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	468	PRO	N-CD	5.93	1.56	1.47

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	474	ARG	CB-CA-C	5.70	121.81	110.40
1	B	364	VAL	N-CA-C	-5.55	96.00	111.00
1	A	454	ASP	CB-CA-C	-5.53	99.34	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2473	0	2440	53	0
1	B	2505	0	2469	55	0
All	All	4978	0	4909	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:470:TYR:CE1	1:A:474:ARG:HD3	1.60	1.33
1:B:300:PHE:N	1:B:367:ALA:HB2	1.73	1.03
1:A:320:TYR:O	1:A:344:LEU:HD22	1.55	1.02
1:B:332:LYS:HB2	1:B:334:PHE:HE1	1.30	0.94
1:A:470:TYR:CE1	1:A:474:ARG:CD	2.50	0.94
1:B:552:LEU:HD23	1:B:555:GLN:OE1	1.66	0.94
1:A:470:TYR:HE1	1:A:474:ARG:CD	1.81	0.93
1:B:299:LEU:HD12	1:B:339:LEU:HD11	1.55	0.88
1:A:470:TYR:HE1	1:A:474:ARG:HD3	1.10	0.88
1:A:470:TYR:CD1	1:A:474:ARG:HD3	2.08	0.88
1:B:303:ASN:HD21	1:B:363:ARG:NH1	1.73	0.85
1:A:467:ASN:OD1	1:A:468:PRO:HD3	1.77	0.84
1:B:365:ARG:HD2	1:B:454:ASP:OD1	1.78	0.83
1:B:300:PHE:H	1:B:367:ALA:HB2	1.42	0.81
1:B:332:LYS:HB2	1:B:334:PHE:CE1	2.15	0.81
1:B:293:TYR:O	1:B:338:LYS:HD3	1.81	0.79
1:B:591:ARG:O	1:B:592:GLN:HB2	1.82	0.78
1:A:478:PRO:HB3	1:B:438:LEU:HD23	1.68	0.74
1:B:298:ARG:NH2	1:B:325:GLU:OE2	2.24	0.70
1:A:467:ASN:OD1	1:A:468:PRO:CD	2.43	0.67
1:B:376:ARG:HG3	1:B:446:ILE:HB	1.75	0.67
1:A:377:ASN:ND2	1:B:494:TRP:CZ3	2.64	0.66
1:B:365:ARG:HD2	1:B:454:ASP:CG	2.15	0.65
1:B:339:LEU:HD13	1:B:345:ALA:HA	1.78	0.65
1:A:467:ASN:CG	1:A:468:PRO:CD	2.64	0.65
1:A:317:PHE:CE1	1:A:348:ALA:HB1	2.34	0.62
1:A:317:PHE:HE1	1:A:348:ALA:HB1	1.63	0.62
1:A:455:ASP:HB2	1:B:399:ARG:HH22	1.63	0.61
1:B:368:THR:HB	1:B:452:GLN:HB3	1.82	0.61
1:A:461:GLU:O	1:A:465:GLN:HG3	2.01	0.61
1:A:523:MET:O	1:A:527:TYR:HB3	2.01	0.60
1:A:468:PRO:HD2	1:A:469:MET:H	1.67	0.60
1:A:467:ASN:HD22	1:B:381:TYR:HE1	1.48	0.60
1:B:436:PHE:CE2	1:B:438:LEU:HD21	2.37	0.59
1:A:377:ASN:HB2	1:A:446:ILE:HD12	1.83	0.59
1:A:470:TYR:HE1	1:A:474:ARG:NH1	2.01	0.59
1:A:461:GLU:HG2	1:A:465:GLN:OE1	2.01	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:303:ASN:HD22	1:B:363:ARG:HG3	1.68	0.59
1:B:332:LYS:CB	1:B:334:PHE:CE1	2.86	0.59
1:B:303:ASN:ND2	1:B:363:ARG:NH1	2.52	0.56
1:A:470:TYR:CE1	1:A:474:ARG:NH1	2.75	0.54
1:A:465:GLN:NE2	1:A:470:TYR:OH	2.41	0.54
1:B:299:LEU:CD1	1:B:339:LEU:HD11	2.32	0.54
1:A:465:GLN:HB3	1:A:470:TYR:CE2	2.43	0.53
1:A:436:PHE:HD1	1:B:480:PHE:CD2	2.26	0.53
1:A:455:ASP:O	1:B:399:ARG:NH1	2.43	0.52
1:A:319:LYS:CG	1:A:319:LYS:O	2.57	0.52
1:A:344:LEU:HA	1:A:347:ILE:HD12	1.92	0.52
1:A:478:PRO:HB3	1:B:438:LEU:CD2	2.40	0.51
1:B:438:LEU:CD1	1:B:445:VAL:HG21	2.41	0.51
1:A:470:TYR:HE1	1:A:474:ARG:CZ	2.24	0.51
1:B:295:GLN:O	1:B:298:ARG:HB2	2.12	0.49
1:A:442:PRO:O	1:B:490:TYR:HE2	1.95	0.49
1:A:295:GLN:HE22	1:A:371:ALA:CB	2.25	0.49
1:A:467:ASN:ND2	1:B:381:TYR:CE1	2.77	0.49
1:A:293:TYR:HA	1:A:339:LEU:O	2.13	0.49
1:B:303:ASN:ND2	1:B:363:ARG:HG3	2.27	0.49
1:A:373:LEU:HD23	1:A:449:PRO:HA	1.94	0.48
1:A:467:ASN:CG	1:A:468:PRO:HD2	2.33	0.48
1:B:295:GLN:HG2	1:B:369:HIS:ND1	2.29	0.48
1:B:292:THR:HG22	1:B:340:GLU:O	2.13	0.48
1:A:436:PHE:HD1	1:B:480:PHE:HD2	1.62	0.48
1:B:422:PRO:HA	1:B:425:ARG:HD2	1.96	0.48
1:A:442:PRO:O	1:B:490:TYR:CE2	2.67	0.47
1:B:339:LEU:CD1	1:B:345:ALA:HA	2.43	0.47
1:B:299:LEU:C	1:B:367:ALA:HB2	2.32	0.47
1:A:446:ILE:HD11	1:B:494:TRP:HZ3	1.80	0.47
1:B:370:ALA:O	1:B:371:ALA:HB3	2.15	0.47
1:A:470:TYR:HE1	1:A:474:ARG:HH11	1.62	0.47
1:A:470:TYR:HE1	1:A:474:ARG:NE	2.12	0.47
1:B:325:GLU:CD	1:B:338:LYS:HE3	2.35	0.47
1:B:438:LEU:HD12	1:B:445:VAL:HG21	1.97	0.47
1:A:299:LEU:HD23	1:A:366:PHE:HA	1.98	0.46
1:A:517:ASP:HA	1:A:520:GLU:HB2	1.96	0.46
1:B:344:LEU:HA	1:B:347:ILE:HD12	1.97	0.46
1:B:591:ARG:O	1:B:592:GLN:CB	2.59	0.45
1:B:365:ARG:O	1:B:366:PHE:C	2.55	0.45
1:A:468:PRO:CD	1:A:469:MET:H	2.30	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:319:LYS:O	1:A:319:LYS:HG2	2.16	0.44
1:B:374:SER:HB2	1:B:450:LEU:HD12	1.98	0.44
1:A:543:GLN:HE21	1:A:547:ARG:HG3	1.82	0.44
1:B:390:ALA:HB1	1:B:438:LEU:CD2	2.48	0.44
1:A:295:GLN:HE22	1:A:371:ALA:HB3	1.83	0.44
1:A:379:SER:HB2	1:A:382:VAL:HG23	2.00	0.43
1:A:310:GLU:HG2	1:A:314:LYS:HE3	2.00	0.43
1:A:404:VAL:HG12	1:A:410:SER:HA	2.00	0.43
1:B:302:GLY:O	1:B:303:ASN:HB2	2.18	0.43
1:B:365:ARG:CD	1:B:454:ASP:CG	2.87	0.42
1:B:460:PRO:HB2	1:B:462:LYS:HG2	2.00	0.42
1:A:467:ASN:ND2	1:B:381:TYR:OH	2.53	0.42
1:B:590:ARG:O	1:B:590:ARG:HG2	2.18	0.42
1:A:377:ASN:ND2	1:A:444:PRO:O	2.53	0.41
1:B:379:SER:HB3	1:B:382:VAL:HG23	2.02	0.41
1:B:390:ALA:HB1	1:B:438:LEU:HD21	2.03	0.41
1:A:436:PHE:O	1:A:437:LEU:HD23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	297/326 (91%)	288 (97%)	9 (3%)	0	100	100
1	B	300/326 (92%)	290 (97%)	10 (3%)	0	100	100
All	All	597/652 (92%)	578 (97%)	19 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	264/286 (92%)	253 (96%)	11 (4%)	30	63
1	B	267/286 (93%)	257 (96%)	10 (4%)	34	65
All	All	531/572 (93%)	510 (96%)	21 (4%)	33	64

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	297	CYS
1	A	374	SER
1	A	384	ASN
1	A	433	GLU
1	A	459	LEU
1	A	479	ARG
1	A	513	LYS
1	A	517	ASP
1	A	523	MET
1	A	546	LEU
1	A	588	GLN
1	B	295	GLN
1	B	354	ASP
1	B	480	PHE
1	B	489	GLU
1	B	508	VAL
1	B	509	GLU
1	B	514[A]	ASP
1	B	514[B]	ASP
1	B	539	LEU
1	B	569	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	295	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	329	ASN
1	A	377	ASN
1	A	465	GLN
1	A	467	ASN
1	A	471	GLN
1	A	543	GLN
1	B	303	ASN
1	B	492	GLN
1	B	503	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.