



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 12:42 am BST

PDB ID : 3WJ5
Title : Crystal structure of PPARgamma ligand binding domain in complex with triphenyltin
Authors : Harada, S.; Hiromori, Y.; Fukakusa, S.; Kawahara, K.; Nakamura, S.; Noda, M.; Uchiyama, S.; Fukui, K.; Nishikawa, J.; Nagase, H.; Kobayashi, Y.; Ohkubo, T.; Yoshida, T.; Nakanishi, T.
Deposited on : 2013-10-04
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	<div> <div>6%</div> <div> <div></div> <div>78%</div> <div>14%</div> <div>• 7%</div> </div> </div>
1	B	276	<div> <div>7%</div> <div> <div></div> <div>82%</div> <div>9%</div> <div>• 7%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	T9T	A	501[A]	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4419 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Peroxisome proliferator-activated receptor gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	258	Total	C	N	O	S	0	1	0
			2062	1330	336	386	10			
1	B	258	Total	C	N	O	S	0	3	0
			2074	1342	340	382	10			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	202	GLY	-	EXPRESSION TAG	UNP P37231
A	203	PRO	-	EXPRESSION TAG	UNP P37231
A	204	LEU	-	EXPRESSION TAG	UNP P37231
A	205	GLY	-	EXPRESSION TAG	UNP P37231
A	206	SER	-	EXPRESSION TAG	UNP P37231
B	202	GLY	-	EXPRESSION TAG	UNP P37231
B	203	PRO	-	EXPRESSION TAG	UNP P37231
B	204	LEU	-	EXPRESSION TAG	UNP P37231
B	205	GLY	-	EXPRESSION TAG	UNP P37231
B	206	SER	-	EXPRESSION TAG	UNP P37231

- Molecule 2 is triphenylstannanyl (three-letter code: T9T) (formula: C₁₈H₁₅Sn).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	Sn	0	1
			38	36	2		
2	B	1	Total	C	Sn	0	0
			19	18	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	103	Total	O	0	0
			103	103		
3	B	123	Total	O	0	0
			123	123		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peroxisome proliferator-activated receptor gamma



- Molecule 1: Peroxisome proliferator-activated receptor gamma



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.52Å 88.49Å 57.94Å 90.00° 91.06° 90.00°	Depositor
Resolution (Å)	36.51 – 1.89 36.51 – 1.89	Depositor EDS
% Data completeness (in resolution range)	99.5 (36.51-1.89) 99.5 (36.51-1.89)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.74 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.203 , 0.247 0.201 , 0.246	Depositor DCC
R_{free} test set	2295 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	21.6	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.001 for l,k,-h 0.030 for h,-k,-l 0.019 for l,-k,h	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4419	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: T9T

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.78	0/2097	0.70	0/2821
1	B	0.78	0/2116	0.75	2/2843 (0.1%)
All	All	0.78	0/4213	0.72	2/5664 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	439	MET	CB-CG-SD	-5.13	97.02	112.40
1	B	234	ARG	NE-CZ-NH2	-5.08	117.76	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2062	0	2123	38	0
1	B	2074	0	2141	34	0
2	A	38	0	30	12	0
2	B	19	0	15	5	0
3	A	103	0	0	0	0
3	B	123	0	0	1	0
All	All	4419	0	4309	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:285[B]:CYS:SG	2:B:501:T9T:C1	2.60	0.90
1:A:448:GLU:O	1:A:452:LEU:HD13	1.75	0.87
1:A:327:TYR:HE1	2:A:501[A]:T9T:C17	1.91	0.83
1:A:360:PHE:HD1	2:A:501[A]:T9T:H4	1.44	0.83
1:B:285[B]:CYS:SG	2:B:501:T9T:C2	2.70	0.79
1:B:439:MET:HG2	3:B:637:HOH:O	1.84	0.77
1:A:279:ILE:CD1	1:A:357:ARG:HH21	1.99	0.75
1:A:279:ILE:HD11	1:A:357:ARG:NH2	2.02	0.73
1:B:363:PHE:HB3	1:B:364:MET:CE	2.19	0.73
1:A:279:ILE:O	1:A:283:GLN:HG2	1.87	0.73
1:A:441:ASP:O	1:A:445:ILE:HG12	1.92	0.69
1:B:363:PHE:HB3	1:B:364:MET:HE2	1.75	0.68
1:B:364:MET:HB3	2:B:501:T9T:C17	2.24	0.68
1:B:325:ILE:HD11	1:B:392:ILE:HG13	1.77	0.67
1:B:264:PHE:HE1	1:B:342:SER:HG	1.42	0.66
1:A:325:ILE:HG12	1:A:388:ILE:HG23	1.77	0.65
1:A:288:ARG:HH11	1:A:288:ARG:HG3	1.62	0.65
2:A:501[B]:T9T:H19	2:A:501[B]:T9T:H13	1.79	0.65
1:A:288:ARG:CG	1:A:288:ARG:HH11	2.11	0.64
1:B:443:ARG:O	1:B:447:THR:HG23	1.98	0.64
1:A:219:TYR:CZ	1:A:223:ILE:HD11	2.33	0.63
1:B:325:ILE:HD11	1:B:392:ILE:CG1	2.29	0.62
1:A:327:TYR:CE1	2:A:501[A]:T9T:H17	2.35	0.62
1:A:325:ILE:HG23	1:A:388:ILE:HD12	1.82	0.60
1:B:275:LYS:HE3	1:B:276:GLU:H	1.68	0.58
1:A:327:TYR:HE1	2:A:501[A]:T9T:H17	1.64	0.58
1:A:330:LEU:HD21	1:A:364:MET:HE1	1.84	0.58
1:B:466:HIS:HB2	1:B:467:PRO:HD3	1.86	0.58
1:A:287:PHE:HE2	1:A:291:GLU:OE2	1.87	0.58
1:A:325:ILE:HD11	1:A:392:ILE:HG13	1.86	0.57
1:A:318:LEU:O	1:A:322:VAL:HG13	2.06	0.56
1:B:264:PHE:HE1	1:B:342:SER:OG	1.88	0.55
1:A:327:TYR:CE1	2:A:501[A]:T9T:C17	2.80	0.55
1:A:279:ILE:CD1	1:A:357:ARG:NH2	2.65	0.55
1:B:466:HIS:CB	1:B:467:PRO:HD3	2.37	0.55
1:A:360:PHE:CD1	2:A:501[A]:T9T:H4	2.34	0.53
1:B:364:MET:HB3	2:B:501:T9T:C15	2.38	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:335:ASN:ND2	1:A:337:ASP:H	2.06	0.53
1:B:234:ARG:O	1:B:238:THR:HG23	2.08	0.53
1:B:363:PHE:HB3	1:B:364:MET:HE3	1.89	0.53
1:A:279:ILE:HD13	1:A:357:ARG:HH21	1.75	0.52
1:A:465:LEU:HD23	1:A:470:GLN:HG2	1.93	0.51
1:B:363:PHE:HE1	1:B:456:ILE:HD11	1.76	0.51
1:A:335:ASN:HD22	1:A:337:ASP:H	1.58	0.50
1:B:255:LEU:O	1:B:259:GLU:HG3	2.12	0.50
1:B:433:ALA:O	1:B:437:GLN:HG3	2.12	0.50
1:A:363:PHE:CZ	1:A:452:LEU:HB3	2.47	0.48
1:A:335:ASN:HD22	1:A:335:ASN:C	2.16	0.48
1:B:277:VAL:HG22	1:B:280:ARG:HH21	1.78	0.48
1:B:255:LEU:CD2	1:B:277:VAL:HG13	2.44	0.48
1:B:457:LYS:HD3	1:B:458:LYS:HE3	1.95	0.48
1:A:255:LEU:CD2	1:A:277:VAL:HG23	2.44	0.47
1:A:327:TYR:OH	2:A:501[B]:T9T:H6	2.15	0.47
1:B:466:HIS:CD2	1:B:466:HIS:N	2.83	0.46
1:B:430:GLN:O	1:B:434:LYS:HG3	2.15	0.46
1:B:255:LEU:CD2	1:B:277:VAL:CG1	2.94	0.46
1:A:285:CYS:SG	2:A:501[A]:T9T:C14	3.04	0.45
1:B:255:LEU:HD23	1:B:277:VAL:CG1	2.46	0.45
1:B:363:PHE:CE1	1:B:456:ILE:HD11	2.51	0.45
1:A:287:PHE:CE2	1:A:291:GLU:OE2	2.68	0.45
1:A:239:GLY:O	1:A:240:LYS:HB2	2.17	0.44
1:A:327:TYR:HE1	2:A:501[B]:T9T:C5	2.30	0.44
1:B:364:MET:HA	2:B:501:T9T:H15	2.00	0.44
1:B:469:LEU:HA	1:B:469:LEU:HD12	1.67	0.44
1:A:453:LEU:CD2	2:A:501[A]:T9T:H9	2.48	0.44
1:A:274:SER:OG	1:A:275:LYS:N	2.51	0.44
1:A:288:ARG:CG	1:A:288:ARG:NH1	2.78	0.43
1:B:466:HIS:CG	1:B:467:PRO:HD3	2.54	0.43
1:A:305:GLY:HA2	1:A:308:ASN:HD22	1.84	0.42
1:B:252:MET:SD	1:B:277:VAL:HG11	2.60	0.42
1:B:255:LEU:HD23	1:B:277:VAL:HG11	2.01	0.42
1:B:364:MET:HG2	1:B:364:MET:H	1.57	0.41
1:A:364:MET:SD	2:A:501[A]:T9T:H16	2.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	253/276 (92%)	250 (99%)	2 (1%)	1 (0%)	34	24
1	B	253/276 (92%)	250 (99%)	3 (1%)	0	100	100
All	All	506/552 (92%)	500 (99%)	5 (1%)	1 (0%)	47	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	461	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/247 (94%)	221 (96%)	10 (4%)	29	19
1	B	232/247 (94%)	222 (96%)	10 (4%)	29	19
All	All	463/494 (94%)	443 (96%)	20 (4%)	28	19

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	234	ARG
1	A	240	LYS
1	A	244	LYS
1	A	245	SER
1	A	262	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	288	ARG
1	A	335	ASN
1	A	358	LYS
1	A	441	ASP
1	A	464	SER
1	B	234	ARG
1	B	240	LYS
1	B	262	ILE
1	B	265	LYS
1	B	266	HIS
1	B	275	LYS
1	B	363	PHE
1	B	364	MET
1	B	466	HIS
1	B	469	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	308	ASN
1	A	314	GLN
1	A	335	ASN
1	A	430	GLN
1	B	412	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	T9T	B	501	-	21,21,21	0.73	0	18,27,27	1.11	2 (11%)
2	T9T	A	501[B]	-	21,21,21	0.70	0	18,27,27	1.49	3 (16%)
2	T9T	A	501[A]	-	21,21,21	0.69	0	18,27,27	1.46	6 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	T9T	B	501	-	-	-	0/3/3/3
2	T9T	A	501[B]	-	-	-	0/3/3/3
2	T9T	A	501[A]	-	-	-	0/3/3/3

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	501[B]	T9T	C3-C1-C2	3.15	120.87	116.54
2	A	501[A]	T9T	C19-C14-C16	2.99	120.64	116.54
2	A	501[B]	T9T	C19-C14-C16	2.60	120.11	116.54
2	A	501[A]	T9T	C15-C16-C14	-2.46	118.85	122.03
2	B	501	T9T	C19-C14-C16	2.34	119.75	116.54
2	A	501[A]	T9T	C13-C8-C11	2.33	119.75	116.54
2	A	501[B]	T9T	C13-C8-C11	2.28	119.67	116.54
2	A	501[A]	T9T	C9-C11-C8	-2.23	119.15	122.03
2	B	501	T9T	C18-C19-C14	-2.21	119.18	122.03
2	A	501[A]	T9T	C4-C2-C1	-2.07	119.36	122.03
2	A	501[A]	T9T	C3-C1-C2	2.01	119.30	116.54

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.


3 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	T9T	5	0
2	A	501[B]	T9T	3	0
2	A	501[A]	T9T	9	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	258/276 (93%)	0.29	17 (6%)	18 20	9, 21, 43, 51	0
1	B	258/276 (93%)	0.38	19 (7%)	14 16	8, 20, 52, 61	2 (0%)
All	All	516/552 (93%)	0.33	36 (6%)	16 18	8, 21, 45, 61	2 (0%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	262	ILE	5.1
1	B	459	THR	4.7
1	B	264	PHE	4.7
1	A	274	SER	4.6
1	B	363	PHE	4.6
1	A	206	SER	4.5
1	B	458	LYS	4.4
1	B	454	GLN	4.3
1	A	363	PHE	4.2
1	B	244	LYS	4.1
1	B	260	ASP	3.7
1	B	261	LYS	3.5
1	B	455	VAL	3.5
1	B	263	LYS	3.2
1	A	464	SER	3.2
1	A	275	LYS	3.1
1	B	239	GLY	3.1
1	A	205	GLY	3.1
1	B	240	LYS	3.0
1	B	475	ASP	2.8
1	B	457	LYS	2.8
1	B	245	SER	2.7
1	A	465	LEU	2.7
1	A	287	PHE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	456	ILE	2.4
1	A	462	ASP	2.4
1	A	277	VAL	2.3
1	A	283	GLN	2.3
1	A	244	LYS	2.3
1	B	466	HIS	2.3
1	A	475	ASP	2.2
1	B	327	TYR	2.2
1	B	259	GLU	2.1
1	A	280	ARG	2.1
1	A	207	GLU	2.0
1	A	260	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	T9T	B	501	19/19	0.87	0.31	49,54,55,55	0
2	T9T	A	501[B]	19/19	0.88	0.33	32,33,34,34	19
2	T9T	A	501[A]	19/19	0.88	0.33	42,45,46,46	19

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around T9T B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around T9T A 501 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around T9T A 501 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.