



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 06:17 am BST

PDB ID : 3WKM
Title : The periplasmic PDZ tandem fragment of the RseP homologue from *Aquifex aeolicus* in complex with the Fab fragment
Authors : Nogi, T.; Tabata, S.; Tamura-kawakami, K.; Takagi, J.
Deposited on : 2013-10-28
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	180	
1	B	180	
2	H	225	
2	I	225	
3	L	218	
3	M	218	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9781 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative zinc metalloprotease aq_1964.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	176	Total	C	N	O	S	0	6	0
			1433	928	249	255	1			
1	B	179	Total	C	N	O	S	0	3	0
			1427	922	245	259	1			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	113	GLY	-	EXPRESSION TAG	UNP O67776
A	114	SER	-	EXPRESSION TAG	UNP O67776
B	113	GLY	-	EXPRESSION TAG	UNP O67776
B	114	SER	-	EXPRESSION TAG	UNP O67776

- Molecule 2 is a protein called MOUSE IGG1-KAPPA FAB (HEAVY CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	216	Total	C	N	O	S	0	1	0
			1647	1042	271	327	7			
2	I	194	Total	C	N	O	S	0	3	0
			1502	947	250	299	6			

- Molecule 3 is a protein called MOUSE IGG1-KAPPA FAB (LIGHT CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	215	Total	C	N	O	S	0	3	0
			1692	1056	284	345	7			
3	M	214	Total	C	N	O	S	0	4	0
			1693	1057	283	346	7			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	67	Total 67	O 67	0	0
4	H	71	Total 71	O 71	0	0
4	L	87	Total 87	O 87	0	0
4	B	53	Total 53	O 53	0	0
4	I	49	Total 49	O 49	0	0
4	M	60	Total 60	O 60	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative zinc metalloprotease aq_1964




- Molecule 1: Putative zinc metalloprotease aq_1964




- Molecule 2: MOUSE IGG1-KAPPA FAB (HEAVY CHAIN)




- Molecule 2: MOUSE IGG1-KAPPA FAB (HEAVY CHAIN)







ASP
CYS

- Molecule 3: MOUSE IGG1-KAPPA FAB (LIGHT CHAIN)



ASN
GLU
CYS

- Molecule 3: MOUSE IGG1-KAPPA FAB (LIGHT CHAIN)



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	51.24Å 86.04Å 92.54Å 100.84° 96.92° 101.32°	Depositor
Resolution (Å)	47.44 – 2.20 47.44 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.2 (47.44-2.20) 98.2 (47.44-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.224 , 0.272 0.224 , 0.271	Depositor DCC
R_{free} test set	3771 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	32.0	Xtrriage
Anisotropy	0.459	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 36.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9781	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	1/1456 (0.1%)	0.58	0/1960
1	B	0.46	0/1450	0.59	0/1954
2	H	0.53	0/1690	0.59	0/2312
2	I	0.52	2/1538 (0.1%)	0.56	0/2097
3	L	0.52	3/1732 (0.2%)	0.63	0/2357
3	M	0.51	3/1733 (0.2%)	0.60	0/2358
All	All	0.50	9/9599 (0.1%)	0.59	0/13038

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	L	167	TRP	CD2-CE2	5.48	1.48	1.41
3	M	167	TRP	CD2-CE2	5.30	1.47	1.41
3	L	39	TRP	CD2-CE2	5.16	1.47	1.41
3	M	152	TRP	CD2-CE2	5.16	1.47	1.41
2	I	164	TRP	CD2-CE2	5.09	1.47	1.41
2	I	47	TRP	CD2-CE2	5.08	1.47	1.41
1	A	157	TRP	CD2-CE2	5.08	1.47	1.41
3	M	100	TRP	CD2-CE2	5.04	1.47	1.41
3	L	152	TRP	CD2-CE2	5.03	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1433	0	1547	36	0
1	B	1427	0	1528	31	0
2	H	1647	0	1600	14	0
2	I	1502	0	1447	10	0
3	L	1692	0	1610	11	0
3	M	1693	0	1609	17	0
4	A	67	0	0	2	0
4	B	53	0	0	1	0
4	H	71	0	0	1	0
4	I	49	0	0	0	0
4	L	87	0	0	0	0
4	M	60	0	0	4	0
All	All	9781	0	9341	107	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (107) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:138[B]:ILE:CD1	1:B:138[B]:ILE:HD12	1.76	1.15
1:A:138[B]:ILE:HD12	1:B:138[B]:ILE:CD1	1.83	1.08
2:H:23:THR:HG22	4:H:316:HOH:O	1.54	1.03
1:A:138[B]:ILE:HD12	1:B:138[B]:ILE:HD12	1.37	1.03
1:A:150:ASN:CG	1:A:151:GLY:H	1.65	0.96
1:B:150:ASN:CG	1:B:151:GLY:H	1.79	0.83
1:A:138[B]:ILE:HD11	1:B:138[B]:ILE:HD12	1.60	0.80
1:A:150:ASN:CG	1:A:151:GLY:N	2.35	0.79
4:A:2006:HOH:O	1:B:191:LYS:HE3	1.82	0.78
2:I:115[A]:GLN:O	2:I:115[A]:GLN:HG3	1.81	0.78
3:L:41:GLN:HB2	3:L:51:LEU:HD11	1.68	0.75
1:B:138[A]:ILE:HG23	4:B:328:HOH:O	1.86	0.75
1:A:150:ASN:HD21	1:A:152:TYR:HD1	1.34	0.74
1:A:138[A]:ILE:HD13	1:A:190:ILE:CG2	2.19	0.72
1:B:150:ASN:CG	1:B:151:GLY:N	2.42	0.71
1:B:150:ASN:HD21	1:B:152:TYR:HD1	1.36	0.70
1:A:138[A]:ILE:HD13	1:A:190:ILE:HG22	1.78	0.66
2:I:20:LEU:HD22	2:I:117:THR:HG21	1.78	0.64
3:L:112:ARG:HD3	3:L:113:ALA:O	1.98	0.64
1:B:149:ILE:O	1:B:150:ASN:ND2	2.31	0.63
1:A:179[B]:LEU:C	1:A:179[B]:LEU:HD12	2.18	0.63
1:A:149:ILE:O	1:A:150:ASN:ND2	2.31	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:146:ILE:HD12	1:B:177:LEU:HD21	1.82	0.62
1:B:138[A]:ILE:HD13	1:B:190:ILE:HG22	1.82	0.62
3:L:93:GLN:HG3	3:L:102:PHE:CE1	2.35	0.61
1:A:138[B]:ILE:CD1	1:B:138[B]:ILE:CD1	2.53	0.61
1:A:158:GLU:HG3	1:A:161[B]:ARG:NH2	2.16	0.59
1:A:235:ASN:HD21	1:A:257:ILE:HG22	1.66	0.59
3:M:95:ASN:HB2	3:M:100:TRP:CE2	2.38	0.59
1:A:150:ASN:OD1	1:A:151:GLY:N	2.21	0.59
1:B:122[B]:GLU:N	1:B:122[B]:GLU:OE2	2.36	0.58
1:B:138[B]:ILE:HG13	1:B:138[B]:ILE:O	2.03	0.58
3:L:63:PRO:HB2	3:L:65:ARG:HG2	1.86	0.58
3:M:111:LYS:HE2	4:M:321:HOH:O	2.03	0.58
1:B:138[A]:ILE:HD13	1:B:190:ILE:CG2	2.34	0.57
1:A:209:PRO:HA	1:A:242:TRP:HE3	1.71	0.56
1:A:138[B]:ILE:HD12	1:B:138[B]:ILE:HD11	1.85	0.54
2:H:30:LYS:HE3	2:H:74:THR:HG21	1.89	0.54
3:M:44:PRO:HA	4:M:305:HOH:O	2.06	0.54
3:M:87:ALA:HB2	3:M:110:ILE:HD12	1.89	0.54
3:L:185:LEU:HD13	3:L:190:TYR:HB2	1.89	0.53
2:H:67:LYS:NZ	2:H:67:LYS:HB2	2.21	0.53
1:A:150:ASN:ND2	1:A:167:LEU:HD21	2.24	0.53
1:B:150:ASN:OD1	1:B:151:GLY:N	2.28	0.53
1:B:233:GLU:HG2	1:B:238:LYS:HD2	1.89	0.53
4:A:2052:HOH:O	2:H:99:ARG:HD2	2.11	0.51
1:B:160:LEU:HD21	1:B:203:ILE:HD11	1.92	0.51
3:M:63:PRO:HB2	3:M:65:ARG:HG2	1.93	0.51
1:A:179[B]:LEU:O	1:A:179[B]:LEU:HD12	2.11	0.50
3:M:41:GLN:HB2	3:M:51:LEU:HD11	1.92	0.50
3:M:93:GLN:HG3	3:M:102:PHE:CE1	2.46	0.50
2:H:154:GLY:HA2	2:H:184:LEU:HB3	1.94	0.50
2:H:11:LEU:HD22	2:H:157:PRO:HD3	1.94	0.50
1:A:261:ILE:HD12	1:A:270:LYS:HG3	1.94	0.50
2:H:178:ALA:HA	2:H:187:LEU:HB3	1.93	0.50
2:I:91:THR:HG23	2:I:120:SER:HA	1.93	0.50
1:B:261:ILE:HD12	1:B:270:LYS:HG3	1.94	0.49
2:I:47:TRP:CD2	3:M:100:TRP:HB2	2.48	0.49
2:I:44:GLY:HA3	4:M:356:HOH:O	2.13	0.49
2:H:91:THR:HG23	2:H:120:SER:HA	1.94	0.48
1:A:233:GLU:HB3	1:A:260:LYS:HB2	1.95	0.48
1:A:153:GLU:OE1	1:A:155:ARG:NH2	2.47	0.47
1:A:246:VAL:HG21	2:H:106:TYR:CD1	2.49	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:117:PRO:HG3	3:L:148:ILE:HD11	1.96	0.47
3:M:185:LEU:HD13	3:M:190:TYR:HB2	1.96	0.47
1:A:150:ASN:ND2	1:A:152:TYR:CD1	2.64	0.47
1:B:279:LYS:HD2	3:M:36:PHE:CZ	2.49	0.47
1:A:166[A]:ARG:HH12	1:A:169:LEU:HB2	1.80	0.46
1:B:116:VAL:HG12	4:M:333:HOH:O	2.14	0.46
1:A:150:ASN:HD22	1:A:167:LEU:HD11	1.80	0.46
1:B:114:SER:HB3	3:M:69:SER:HA	1.97	0.46
3:M:132:GLY:HA2	3:M:187:LYS:HE3	1.98	0.46
2:I:115[A]:GLN:O	2:I:115[A]:GLN:CG	2.59	0.46
2:I:34:LEU:HD12	2:I:79:ALA:HB2	1.98	0.46
3:L:153:LYS:HB2	3:L:197:THR:HB	1.98	0.45
3:L:95:ASN:HB2	3:L:100:TRP:CE2	2.51	0.45
3:L:14:SER:HA	3:L:111:LYS:HB3	1.98	0.45
3:M:35:SER:HB2	3:M:55:ALA:HB2	1.98	0.45
3:M:14:SER:HA	3:M:111:LYS:HB3	1.99	0.45
1:A:124:VAL:HG11	1:A:160:LEU:HD22	1.98	0.44
2:H:134:LEU:HD12	2:H:149:GLY:HA3	1.99	0.44
3:M:187:LYS:HA	3:M:190:TYR:HB3	1.98	0.44
1:A:277:ASP:HA	1:A:278:PRO:HD3	1.85	0.43
2:H:33:TYR:CD2	2:H:50:ARG:HD2	2.53	0.43
1:A:247:GLU:O	1:A:251:LYS:HG3	2.19	0.43
2:H:198:TRP:HA	2:H:199:PRO:HA	1.85	0.43
1:A:122:GLU:O	1:A:156:THR:HA	2.19	0.43
1:B:235:ASN:HD21	1:B:257:ILE:HG22	1.83	0.43
1:B:216:LYS:HE2	1:B:217:GLY:N	2.34	0.43
1:B:150:ASN:HD22	1:B:167:LEU:HD11	1.85	0.42
1:B:150:ASN:ND2	1:B:167:LEU:HD11	2.34	0.42
3:M:197:THR:HG23	3:M:210:VAL:HG13	2.02	0.42
1:A:150:ASN:ND2	1:A:167:LEU:HD11	2.35	0.42
1:B:148:LYS:HE2	1:B:151:GLY:HA2	2.02	0.42
1:A:150:ASN:CG	1:A:167:LEU:HD21	2.40	0.42
1:A:215:LYS:HG3	3:L:32:TYR:CZ	2.54	0.41
1:A:138[B]:ILE:HG23	1:A:138[B]:ILE:O	2.20	0.41
2:I:12:VAL:HG11	2:I:18:VAL:HB	2.01	0.41
1:A:138[A]:ILE:HD11	1:A:140:ILE:HD11	2.02	0.41
1:B:246:VAL:HG21	2:I:106:TYR:CD1	2.55	0.41
1:A:257:ILE:HG13	1:A:272:LEU:HD23	2.03	0.41
2:I:51:ILE:O	2:I:53:PRO:HD3	2.21	0.41
3:L:179:MET:HE3	3:L:181:SER:HB2	2.03	0.40
2:H:156:PHE:HA	2:H:157:PRO:HA	1.79	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:150:ASN:HB3	1:B:175:THR:OG1	2.21	0.40
2:H:51:ILE:O	2:H:53:PRO:HD3	2.21	0.40
3:M:116:ALA:HA	3:M:117:PRO:HD2	1.96	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	180/180 (100%)	177 (98%)	3 (2%)	0	100	100
1	B	180/180 (100%)	174 (97%)	6 (3%)	0	100	100
2	H	213/225 (95%)	202 (95%)	11 (5%)	0	100	100
2	I	191/225 (85%)	181 (95%)	9 (5%)	1 (0%)	29	31
3	L	216/218 (99%)	209 (97%)	5 (2%)	2 (1%)	17	16
3	M	216/218 (99%)	202 (94%)	12 (6%)	2 (1%)	17	16
All	All	1196/1246 (96%)	1145 (96%)	46 (4%)	5 (0%)	41	37

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	L	203	LYS
3	L	72	ARG
2	I	213	SER
3	M	72[A]	ARG
3	M	72[B]	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	158/155 (102%)	154 (98%)	4 (2%)	47	60
1	B	157/155 (101%)	154 (98%)	3 (2%)	57	71
2	H	187/193 (97%)	181 (97%)	6 (3%)	39	50
2	I	168/193 (87%)	165 (98%)	3 (2%)	59	72
3	L	192/192 (100%)	188 (98%)	4 (2%)	53	67
3	M	192/192 (100%)	187 (97%)	5 (3%)	46	58
All	All	1054/1080 (98%)	1029 (98%)	25 (2%)	52	62

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	141	LYS
1	A	166[A]	ARG
1	A	166[B]	ARG
1	A	266	LYS
2	H	1	GLU
2	H	12	VAL
2	H	57	ASN
2	H	67	LYS
2	H	127	THR
2	H	206	ASN
3	L	112	ARG
3	L	160	GLN
3	L	185	LEU
3	L	212	SER
1	B	216	LYS
1	B	260	LYS
1	B	279	LYS
2	I	25	SER
2	I	217	ASP
2	I	218	LYS
3	M	82[A]	VAL
3	M	82[B]	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	M	147	ASP
3	M	185	LEU
3	M	193	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
3	L	96	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	176/180 (97%)	-0.11	1 (0%) 89 88	20, 30, 44, 59	0
1	B	179/180 (99%)	-0.04	3 (1%) 70 68	21, 31, 53, 78	0
2	H	216/225 (96%)	0.34	12 (5%) 24 23	18, 43, 67, 75	0
2	I	194/225 (86%)	0.92	44 (22%) 0 0	20, 47, 90, 102	0
3	L	215/218 (98%)	0.14	3 (1%) 75 73	16, 32, 56, 66	0
3	M	214/218 (98%)	0.93	46 (21%) 0 0	17, 35, 100, 110	0
All	All	1194/1246 (95%)	0.38	109 (9%) 9 8	16, 34, 85, 110	0

All (109) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	113	GLY	7.3
3	M	124	PRO	6.2
3	M	121	ILE	6.2
2	I	182	SER	6.1
3	M	123	PRO	5.9
3	M	131	SER	5.6
3	M	154	ILE	5.6
3	M	213	PHE	5.6
2	I	134	LEU	5.5
2	I	207	VAL	5.4
3	M	122	PHE	5.1
2	I	216	VAL	5.1
3	M	192	ARG	5.1
3	M	198	CYS	5.0
3	M	196	TYR	4.9
3	M	152	TRP	4.9
3	M	191	GLU	4.7
2	I	133	PRO	4.6
3	M	136	VAL	4.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	I	132	TYR	4.6
3	M	132	GLY	4.5
2	I	148	LEU	4.4
3	M	138	CYS	4.4
2	I	160	VAL	4.3
2	I	169	LEU	4.2
2	I	166	SER	4.2
3	M	159	ARG	4.1
3	M	184	THR	4.0
3	M	188	ASP	4.0
3	M	210	VAL	4.0
2	I	208	ALA	4.0
2	I	151	LEU	4.0
3	M	214	ASN	3.9
2	I	154	GLY	3.8
3	M	130	THR	3.8
2	H	135	ALA	3.8
3	M	157	SER	3.8
3	M	160	GLN	3.7
2	I	162	VAL	3.7
2	I	159	PRO	3.6
2	H	202	THR	3.6
3	M	125	SER	3.4
2	H	194	PRO	3.4
2	I	150	CYS	3.4
2	I	135	ALA	3.3
3	M	133	GLY	3.3
2	I	205	CYS	3.3
3	M	183	LEU	3.3
2	I	128	PRO	3.2
3	M	181	SER	3.2
3	M	185	LEU	3.1
3	M	211	LYS	3.1
2	I	170	SER	3.1
3	L	32	TYR	3.0
3	M	189	GLU	3.0
2	I	211	ALA	2.9
2	H	211	ALA	2.9
3	M	139	PHE	2.9
3	M	128	GLN	2.9
2	I	185	TYR	2.9
2	I	179	VAL	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	I	213	SER	2.9
2	I	206	ASN	2.9
2	I	157	PRO	2.9
2	H	146	VAL	2.8
3	M	134	ALA	2.8
2	I	181	GLN	2.8
3	M	190	TYR	2.8
3	M	150	VAL	2.8
2	I	87	THR	2.8
2	I	164	TRP	2.8
1	B	114	SER	2.8
3	M	32	TYR	2.7
2	I	152	VAL	2.7
3	M	129	LEU	2.7
2	H	87	THR	2.7
2	H	15	GLY	2.6
3	M	186	THR	2.5
3	M	187	LYS	2.5
2	I	12	VAL	2.5
2	I	131	VAL	2.5
2	H	221	VAL	2.4
3	M	197	THR	2.4
3	M	206	THR	2.4
2	I	217	ASP	2.4
2	I	122	SER	2.4
2	I	161	THR	2.3
3	M	163	VAL	2.3
3	M	156	GLY	2.3
2	I	163	THR	2.3
3	M	126	SER	2.3
3	M	162	GLY	2.3
2	I	212	SER	2.3
2	I	186	THR	2.2
2	H	164	TRP	2.2
2	I	123	SER	2.2
3	L	122	PHE	2.2
1	B	115	GLU	2.2
2	H	217	ASP	2.2
3	M	135	SER	2.2
2	I	155	TYR	2.1
2	I	174	HIS	2.1
1	A	116	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	I	218	LYS	2.1
3	L	188	ASP	2.1
2	I	149	GLY	2.1
2	I	180	LEU	2.0
2	H	205	CYS	2.0
2	H	220	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.