



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 25, 2023 – 04:56 PM EDT

PDB ID : 5WK4
Title : Crystal structure of an anti-idiotypic VLR
Authors : Collins, B.C.; Nakahara, H.; Cooper, M.D.; Herrin, B.R.; Wilson, I.A.
Deposited on : 2017-07-24
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	201	
1	B	201	
1	C	201	
1	D	201	
1	E	201	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	201	 <p>A horizontal bar chart representing the quality of the chain. The bar is divided into four segments: a small red segment at the beginning labeled '2%', a large green segment labeled '85%', a yellow segment labeled '10%', and a small grey segment at the end labeled '5%'.</p>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9612 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Variable lymphocyte receptor 39.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	D	192	Total 1459	C 918	N 259	O 274	S 8	0	0	0
1	A	190	Total 1443	C 911	N 254	O 270	S 8	0	0	0
1	B	192	Total 1452	C 915	N 259	O 270	S 8	0	0	0
1	C	192	Total 1447	C 911	N 256	O 272	S 8	0	0	0
1	E	191	Total 1438	C 910	N 250	O 270	S 8	0	0	0
1	F	191	Total 1426	C 903	N 250	O 265	S 8	0	0	0

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total 1	Mg 1	0	0
2	A	1	Total 1	Mg 1	0	0
2	E	1	Total 1	Mg 1	0	0
2	F	1	Total 1	Mg 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	175	Total 175	O 175	0	0
3	A	182	Total 182	O 182	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	180	Total 180	O 180	0	0
3	C	140	Total 140	O 140	0	0
3	E	142	Total 142	O 142	0	0
3	F	124	Total 124	O 124	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Variable lymphocyte receptor 39




- Molecule 1: Variable lymphocyte receptor 39




- Molecule 1: Variable lymphocyte receptor 39




- Molecule 1: Variable lymphocyte receptor 39




- Molecule 1: Variable lymphocyte receptor 39



- Molecule 1: Variable lymphocyte receptor 39

Chain F: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	60.41Å 61.38Å 80.56Å 110.09° 99.79° 93.52°	Depositor
Resolution (Å)	34.72 – 1.50 44.01 – 1.50	Depositor EDS
% Data completeness (in resolution range)	85.1 (34.72-1.50) 85.1 (44.01-1.50)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 1.50Å)	Xtrriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, R_{free}	0.193 , 0.223 0.193 , 0.223	Depositor DCC
R_{free} test set	7060 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	19.3	Xtrriage
Anisotropy	0.042	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 33.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9612	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/1477	0.60	0/2021
1	B	0.34	0/1486	0.59	0/2032
1	C	0.33	0/1481	0.58	0/2027
1	D	0.35	0/1493	0.61	0/2041
1	E	0.33	0/1473	0.56	0/2019
1	F	0.31	0/1461	0.55	0/2004
All	All	0.34	0/8871	0.58	0/12144

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1443	0	1417	9	0
1	B	1452	0	1428	5	0
1	C	1447	0	1412	6	0
1	D	1459	0	1434	11	0
1	E	1438	0	1397	10	0
1	F	1426	0	1384	12	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	182	0	0	2	0
3	B	180	0	0	2	0
3	C	140	0	0	2	0
3	D	175	0	0	0	0
3	E	142	0	0	3	0
3	F	124	0	0	1	0
All	All	9612	0	8472	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:108:GLN:NE2	3:E:402:HOH:O	2.21	0.73
1:B:210:LYS:NZ	3:B:301:HOH:O	2.22	0.72
1:C:110:THR:O	3:C:301:HOH:O	2.11	0.67
1:A:136:ILE:HG12	1:A:167:LEU:HB2	1.76	0.67
1:E:114:ALA:O	3:E:401:HOH:O	2.13	0.66
1:A:49:THR:HG22	1:A:71:SER:HB2	1.79	0.65
1:D:58:ASN:ND2	1:D:82:ASP:OD2	2.30	0.64
1:B:207:SER:HB3	1:B:210:LYS:HD2	1.78	0.64
1:B:138:ARG:NH2	3:B:303:HOH:O	2.31	0.62
1:F:58:ASN:ND2	1:F:82:ASP:OD2	2.32	0.62
1:F:48:PRO:O	1:F:51:ARG:HG2	2.04	0.58
1:A:210:LYS:NZ	3:A:403:HOH:O	2.37	0.57
1:A:80:VAL:HG22	1:A:104:LEU:HD23	1.88	0.56
1:D:175:ALA:O	1:D:178:VAL:HG12	2.06	0.55
1:E:39:SER:HB2	1:F:38:ARG:HG2	1.91	0.53
1:F:48:PRO:HB2	1:F:51:ARG:HD3	1.91	0.52
1:D:170:TRP:CH2	1:D:177:ILE:HD11	2.44	0.52
1:E:89:PRO:HG2	1:E:92:VAL:HB	1.92	0.52
1:E:160:GLU:HB3	1:E:210:LYS:HD3	1.93	0.51
1:F:136:ILE:HG12	1:F:167:LEU:HB2	1.93	0.50
1:A:136:ILE:HD13	1:A:167:LEU:HD13	1.94	0.50
1:C:89:PRO:HG2	1:C:92:VAL:HB	1.95	0.48
1:F:28:CYS:SG	1:F:33:VAL:HG22	2.53	0.48
1:E:136:ILE:HG12	1:E:167:LEU:HB2	1.96	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:136:ILE:HD13	1:D:141:PHE:CE2	2.49	0.48
1:A:199:ARG:NH1	3:A:408:HOH:O	2.47	0.47
1:F:33:VAL:CG2	1:F:51:ARG:HG3	2.44	0.47
1:D:80:VAL:HG22	1:D:104:LEU:HD23	1.97	0.46
1:A:72:LEU:HB3	1:A:75:LEU:HG	1.97	0.46
1:C:95:ARG:NH1	3:C:303:HOH:O	2.40	0.45
1:F:210:LYS:HB3	1:F:210:LYS:HE2	1.72	0.45
1:D:80:VAL:CG2	1:D:104:LEU:HD23	2.46	0.45
1:E:210:LYS:HE2	1:E:210:LYS:HB3	1.76	0.43
1:E:68:VAL:HG11	3:E:511:HOH:O	2.17	0.43
1:B:65:GLU:HB2	1:B:68:VAL:HG13	1.99	0.43
1:C:175:ALA:O	1:C:178:VAL:HG12	2.19	0.43
1:D:136:ILE:CG1	1:D:167:LEU:HB2	2.49	0.42
1:F:100:THR:HA	1:F:123:LEU:HA	2.01	0.42
1:D:136:ILE:HD12	1:D:166:TYR:CE2	2.55	0.42
1:D:55:TRP:CZ2	1:E:29:ARG:HG2	2.55	0.42
1:A:159:CYS:SG	1:A:198:VAL:HG22	2.60	0.42
1:A:170:TRP:CH2	1:A:177:ILE:HD11	2.55	0.42
1:E:40:LEU:HD13	1:E:44:PRO:HG3	2.02	0.42
1:F:25:GLN:NE2	3:F:408:HOH:O	2.53	0.42
1:C:80:VAL:CG2	1:C:104:LEU:HD23	2.50	0.42
1:D:72:LEU:HB3	1:D:75:LEU:HG	2.02	0.41
1:D:136:ILE:HG12	1:D:167:LEU:HB2	2.02	0.41
1:F:56:LEU:HB2	1:F:80:VAL:HG12	2.03	0.40
1:B:80:VAL:CG2	1:B:104:LEU:HD23	2.52	0.40
1:C:178:VAL:HG11	1:C:186:VAL:HG23	2.02	0.40
1:F:33:VAL:HG23	1:F:51:ARG:HG3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	188/201 (94%)	181 (96%)	7 (4%)	0	100	100
1	B	190/201 (94%)	183 (96%)	7 (4%)	0	100	100
1	C	190/201 (94%)	182 (96%)	8 (4%)	0	100	100
1	D	190/201 (94%)	183 (96%)	7 (4%)	0	100	100
1	E	189/201 (94%)	181 (96%)	8 (4%)	0	100	100
1	F	189/201 (94%)	182 (96%)	7 (4%)	0	100	100
All	All	1136/1206 (94%)	1092 (96%)	44 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	164/176 (93%)	161 (98%)	3 (2%)	59	30
1	B	164/176 (93%)	161 (98%)	3 (2%)	59	30
1	C	163/176 (93%)	159 (98%)	4 (2%)	47	18
1	D	166/176 (94%)	162 (98%)	4 (2%)	49	19
1	E	161/176 (92%)	155 (96%)	6 (4%)	34	8
1	F	159/176 (90%)	154 (97%)	5 (3%)	40	11
All	All	977/1056 (92%)	952 (97%)	25 (3%)	46	16

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	60	GLN
1	D	99	LEU
1	D	108	GLN
1	D	207	SER
1	A	99	LEU
1	A	168	LYS
1	A	210	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	99	LEU
1	B	178	VAL
1	B	210	LYS
1	C	99	LEU
1	C	147	LEU
1	C	168	LYS
1	C	210	LYS
1	E	70	ASP
1	E	99	LEU
1	E	108	GLN
1	E	147	LEU
1	E	161	CYS
1	E	199	ARG
1	F	60	GLN
1	F	84	GLN
1	F	99	LEU
1	F	108	GLN
1	F	168	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	143	ASN
1	B	60	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	190/201 (94%)	-0.00	3 (1%) 72 77	13, 20, 34, 52	0
1	B	192/201 (95%)	-0.08	1 (0%) 91 93	13, 20, 32, 40	0
1	C	192/201 (95%)	-0.05	1 (0%) 91 93	14, 22, 33, 44	0
1	D	192/201 (95%)	-0.14	2 (1%) 82 85	14, 19, 31, 41	0
1	E	191/201 (95%)	-0.09	3 (1%) 72 77	17, 24, 32, 47	0
1	F	191/201 (95%)	0.22	5 (2%) 56 61	16, 27, 39, 48	0
All	All	1148/1206 (95%)	-0.02	15 (1%) 77 81	13, 22, 35, 52	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	136	ILE	4.1
1	A	209	SER	3.9
1	A	211	CYS	3.6
1	D	136	ILE	2.9
1	E	114	ALA	2.9
1	E	212	PRO	2.9
1	F	212	PRO	2.8
1	D	211	CYS	2.4
1	F	94	ASP	2.3
1	C	209	SER	2.3
1	F	136	ILE	2.3
1	B	114	ALA	2.2
1	F	90	ILE	2.2
1	F	114	ALA	2.1
1	A	204	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MG	A	300	1/1	0.80	0.11	35,35,35,35	0
2	MG	D	300	1/1	0.91	0.14	36,36,36,36	0
2	MG	F	300	1/1	0.91	0.21	46,46,46,46	0
2	MG	E	300	1/1	0.99	0.14	31,31,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.