



# wwPDB X-ray Structure Validation Summary Report

Oct 10, 2023 – 02:00 PM EDT

PDB ID : 6WMW  
Title : GFRAL receptor bound with two antibody Fabs (3P10, 25M22)  
Authors : White, A.; Lakshminarasimhan, D.; Olland, A.; Suto, R.K.  
Deposited on : 2020-04-21  
Resolution : 2.91 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2307 (2.94-2.90)
Clashscore	141614	2531 (2.94-2.90)
Ramachandran outliers	138981	2462 (2.94-2.90)
Sidechain outliers	138945	2464 (2.94-2.90)
RSRZ outliers	127900	2248 (2.94-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	245	 3% 63% 12% 22%
2	H	225	 3% 68% 23% 6%
3	L	218	 2% 67% 29% 2%
4	M	232	 4% 75% 19% 2%
5	N	241	 3% 63% 23% 10%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8097 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GDNF family receptor alpha-like.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	192	1496	921	266	284	25	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	107	HIS	-	expression tag	UNP Q6UXV0
B	108	HIS	-	expression tag	UNP Q6UXV0
B	109	HIS	-	expression tag	UNP Q6UXV0
B	110	HIS	-	expression tag	UNP Q6UXV0
B	111	HIS	-	expression tag	UNP Q6UXV0
B	112	HIS	-	expression tag	UNP Q6UXV0
B	113	GLY	-	expression tag	UNP Q6UXV0
B	114	GLY	-	expression tag	UNP Q6UXV0

- Molecule 2 is a protein called FAB3P10 heavy chain fragment.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	212	1592	1015	258	314	5	0	0	0

- Molecule 3 is a protein called FAB3P10 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	218	1660	1040	278	334	8	0	0	0

- Molecule 4 is a protein called FAB25M22 heavy chain fragment.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	M	222	1671	1060	279	325	7	0	0	0


- Molecule 5 is a protein called FAB25M22 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	N	216	1678	1062	275	337	4	0	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GDNF family receptor alpha-like




- Molecule 2: FAB3P10 heavy chain fragment




- Molecule 3: FAB3P10 light chain




- Molecule 4: FAB25M22 heavy chain fragment

Chain M:  4% 75% 19%



• Molecule 5: FAB25M22 light chain

Chain N:  3% 63% 23% 10%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.50Å 116.05Å 227.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.70 – 2.91 47.65 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.4 (47.70-2.91) 99.3 (47.65-2.90)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.93 (at 2.91Å)	Xtrriage
Refinement program	REFMAC 5.8.0230	Depositor
R, $R_{free}$	0.222 , 0.290 0.225 , 0.288	Depositor DCC
$R_{free}$ test set	1591 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	64.6	Xtrriage
Anisotropy	0.091	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.26 , 29.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	8097	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	77.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.62% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	B	0.44	0/1523	0.67	0/2060
2	H	0.45	0/1633	0.69	0/2230
3	L	0.50	0/1698	0.70	0/2303
4	M	0.43	0/1713	0.67	0/2331
5	N	0.43	0/1715	0.65	0/2329
All	All	0.45	0/8282	0.68	0/11253

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	L	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	L	112	ARG	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1496	0	1434	22	0
2	H	1592	0	1548	33	0
3	L	1660	0	1596	39	0
4	M	1671	0	1642	24	0
5	N	1678	0	1633	36	0
All	All	8097	0	7853	145	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 145 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:M:104:LEU:O	4:M:105:ARG:HD3	1.69	0.92
4:M:87:THR:O	4:M:123:VAL:HG21	1.77	0.83
2:H:67:ARG:NH1	2:H:90:ASP:OD2	2.13	0.81
5:N:214:GLU:HG2	5:N:238:ARG:HH11	1.50	0.76
5:N:214:GLU:HG2	5:N:238:ARG:NH1	2.01	0.76

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	190/245 (78%)	177 (93%)	10 (5%)	3 (2%)	9 31
2	H	208/225 (92%)	166 (80%)	42 (20%)	0	100 100
3	L	216/218 (99%)	189 (88%)	24 (11%)	3 (1%)	11 34
4	M	218/232 (94%)	191 (88%)	25 (12%)	2 (1%)	17 46
5	N	214/241 (89%)	192 (90%)	20 (9%)	2 (1%)	17 46
All	All	1046/1161 (90%)	915 (88%)	121 (12%)	10 (1%)	15 43

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	319	TYR
5	N	95	GLY
5	N	141	SER
1	B	158	ASN
3	L	208	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	173/217 (80%)	159 (92%)	14 (8%)	11	32
2	H	176/191 (92%)	148 (84%)	28 (16%)	2	7
3	L	187/189 (99%)	166 (89%)	21 (11%)	6	17
4	M	185/196 (94%)	162 (88%)	23 (12%)	4	13
5	N	193/214 (90%)	177 (92%)	16 (8%)	11	30
All	All	914/1007 (91%)	812 (89%)	102 (11%)	6	17

5 of 102 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	L	166	SER
4	M	107	THR
5	N	226	GLN
3	L	192	LYS
4	M	37	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

#### 5.7 Other polymers [i](#)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	192/245 (78%)	-0.16	0 <b>100</b> <b>100</b>	35, 59, 94, 137	0
2	H	212/225 (94%)	0.10	7 (3%) 46 42	37, 82, 134, 151	0
3	L	218/218 (100%)	-0.03	5 (2%) 60 59	30, 60, 128, 160	0
4	M	222/232 (95%)	-0.04	10 (4%) 33 30	39, 74, 124, 143	0
5	N	216/241 (89%)	0.07	8 (3%) 41 38	42, 83, 157, 179	0
All	All	1060/1161 (91%)	-0.01	30 (2%) 53 50	30, 71, 135, 179	0

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	130	ALA	5.2
4	M	137	ALA	4.1
2	H	219	LYS	4.0
4	M	150	LEU	3.9
2	H	131	PRO	3.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.