Full wwPDB X-ray Structure Validation Report (i)

Dec 17, 2023 – 06:35 PM EST

PDB ID	:	4WNQ
Title	:	THE MOLECULAR BASES OF DELTA/ALPHA-BETA T-CELL MEDIATED ANTIGEN RECOGNITION
Authors	:	Pellicci, D.G.; Uldrich, A.P.; Le Nours, J.; Ross, F.; Chabrol, E.; Eckle, S.B.G.; de Boer, R.; Lim, R.T.; McPherson, K.; Besra, G.; Howell, A.R.; Moretta, L.; McCluskey, J.; Heemskerk, M.H.M.; Gras, S.; Rossjohn, J.; Godfrey, D.I.
Deposited on	:	2014-10-14
Resolution	:	1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	207	2%, 91%, 5%
1	C	207	90%, 6%
2	B	245	85%, 12%
2	D	245	%, 89%, 9%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 7549 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TCR Variable Delta 1 chain and TCR constant Alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	200	1583	1004	260	310	9	0	3	0
1	C	200	1593	1012	262	310	9	0	5	0

- Molecule 2 is a protein called TCR Variable Beta 2 (TRBV20) chain and TCR constant Beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	241	1914	1210	327	367	10	0	7	0
2	D	242	1914	1209	328	368	9	0	4	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	148	148	148	0	0
3	B	123	123	123	0	0
3	C	152	152	152	0	0
3	D	122	122	122	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: TCR Variable Delta 1 chain and TCR constant Alpha chain

Chain A:





• Molecule 1: TCR Variable Delta 1 chain and TCR constant Alpha chain

Chain C:

• Molecule 2: TCR Variable Beta 2 (TRBV20) chain and TCR constant Beta chain

Chain B:



• Molecule 2: TCR Variable Beta 2 (TRBV20) chain and TCR constant Beta chain

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	39.26Å 47.23Å 109.29Å 88.32° 79.78° 89.93°	Depositor
Resolution (Å)	47.21 – 1.80 47.21 – 1.80	Depositor EDS
% Data completeness (in resolution range)	97.6 (47.21-1.80) 97.6 (47.21-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.11 (at 1.79Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.193 , 0.229 0.203 , 0.241	Depositor DCC
R_{free} test set	3537 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å^2)	17.6	Xtriage
Anisotropy	0.220	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.30 , 22.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.46$, $< L^2 > = 0.29$	Xtriage
Estimated twinning fraction	0.036 for h,-k,h-l 0.135 for -h,k,-l 0.027 for -h,-k,-h+l	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	7549	wwPDB-VP
Average B, all atoms (Å^2)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.89% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.51	0/1627	0.65	0/2204
1	C	0.52	0/1643	0.65	0/2225
2	B	0.51	0/1985	0.65	0/2700
2	D	0.49	0/1976	0.64	0/2688
All	All	0.51	0/7231	0.65	0/9817

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1583	0	1542	6	0
1	C	1593	0	1563	8	0
2	B	1914	0	1851	24	0
2	D	1914	0	1845	15	0
3	A	148	0	0	0	0
3	B	123	0	0	2	0
3	C	152	0	0	0	0
3	D	122	0	0	1	0
All	All	7549	0	6801	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:116:GLN:HG2	3:B:307:HOH:O	1.88	0.72
2:B:95:HIS:HD2	2:B:97:GLU:H	1.39	0.70
1:A:144:LYS:HE2	2:B:157:THR:HG21	1.74	0.68
1:A:185:SER:OG	2:B:202:ARG:HD2	1.97	0.64
2:D:136:VAL:HG23	2:D:246:ALA:HB3	1.80	0.62
1:C:144:LYS:HE2	2:D:157:THR:HG21	1.81	0.61
1:C:75:ARG:NH2	1:C:98:ASP:OD2	2.33	0.60
2:D:200[A]:SER:OG	2:D:202:ARG:NH2	2.34	0.60
1:C:185:SER:OG	2:D:202:ARG:HD2	2.05	0.57
1:A:16:ARG:HH12	1:A:125:ASN:HD22	1.52	0.56
2:B:200:SER:OG	2:B:202:ARG:NH2	2.41	0.53
1:C:39:ILE:HG21	1:C:87:VAL:HG21	1.90	0.53
1:C:40:PHE:CE2	2:D:111:GLY:HA2	2.43	0.53
2:D:108:GLY:HA3	2:D:113:PHE:HE2	1.76	0.50
2:B:95:HIS:CD2	2:B:97:GLU:HB2	2.47	0.50
2:B:78:ILE:CD1	2:B:89:LEU:HD23	2.42	0.49
2:B:78:ILE:HD13	2:B:89:LEU:HD23	1.94	0.49
2:B:131:PRO:HD3	2:B:239:PRO:HB3	1.93	0.49
2:B:151:THR:CG2	2:B:202:ARG:HG2	2.43	0.49
2:D:95:HIS:ND1	2:D:97:GLU:HB2	2.28	0.48
1:A:39:ILE:HG21	1:A:87:VAL:HG21	1.94	0.48
1:A:40:PHE:CE2	2:B:111:GLY:HA2	2.49	0.47
1:A:27:THR:HB	1:A:106:LEU:HD21	1.98	0.46
2:B:108:GLY:HA3	2:B:113:PHE:CE2	2.50	0.45
2:B:14:LYS:NZ	2:B:126:LEU:H	2.15	0.45
2:B:187:LYS:HE2	2:B:195:SER:HB3	1.98	0.44
2:B:22:GLU:HG2	2:B:88[A]:THR:HG22	1.99	0.44
2:D:187:LYS:HE2	2:D:195:SER:HB3	1.99	0.44
2:B:151:THR:HG21	2:B:202:ARG:HG2	1.99	0.44
2:B:186:LEU:HD22	2:B:198:ALA:HB3	2.00	0.44
2:B:95:HIS:HD2	2:B:97:GLU:HB2	1.82	0.43
2:D:214:ARG:NH1	3:D:301:HOH:O	2.42	0.43
1:C:75:ARG:HH22	1:C:98:ASP:CG	2.22	0.43
2:D:186:LEU:HD22	2:D:198:ALA:HB3	2.01	0.43
2:B:187:LYS:HD3	2:B:190:PRO:HA	2.01	0.43
2:B:139:PRO:HD3	2:B:152:LEU:HG	2.01	0.42
2:D:220:GLN:HG3	2:D:243:ILE:HG23	2.01	0.42
2:B:220:GLN:HG3	2:B:243:ILE:HG23	2.01	0.42
2:B:220:GLN:HG3	2:B:243:ILE:CG2	2.50	0.42
2:D:220:GLN:HG3	2:D:243:ILE:CG2	2.50	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:124:GLU:HG2	2:B:128:ASN:OD1	2.20	0.41
1:C:27:THR:HB	1:C:106:LEU:HD21	2.02	0.41
2:D:139:PRO:HD3	2:D:152:LEU:HG	2.02	0.41
2:D:187:LYS:HD3	2:D:190:PRO:HA	2.02	0.41
1:C:29:TRP:CD1	1:C:108:VAL:HG11	2.56	0.41
2:D:131:PRO:HD3	2:D:239:PRO:HB3	2.02	0.40
2:B:214:ARG:NH1	3:B:301:HOH:O	2.47	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	201/207 (97%)	196 (98%)	4 (2%)	1 (0%)	29	15
1	C	203/207 (98%)	197 (97%)	5 (2%)	1 (0%)	29	15
2	B	246/245 (100%)	241 (98%)	5 (2%)	0	100	100
2	D	244/245 (100%)	237 (97%)	7 (3%)	0	100	100
All	All	894/904 (99%)	871 (97%)	21 (2%)	2 (0%)	47	33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	87	VAL
1	C	87	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/185 (98%)	179 (99%)	2 (1%)	73	68
1	C	183/185 (99%)	181 (99%)	2 (1%)	73	68
2	B	216/212 (102%)	211 (98%)	5 (2%)	50	37
2	D	214/212 (101%)	211 (99%)	3 (1%)	67	59
All	All	794/794 (100%)	782 (98%)	12 (2%)	65	56

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	137	ARG
1	A	194	ASP
2	B	89	LEU
2	B	116	GLN
2	B	140	SER
2	B	141	GLU
2	B	202	ARG
1	C	137	ARG
1	C	199	ASN
2	D	116	GLN
2	D	140	SER
2	D	202	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	125	ASN
2	B	95	HIS
2	B	193	ASN
2	B	222	GLN
2	D	222	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	200/207 (96%)	-0.50	4 (2%)	65	61	10, 19, 39, 52	0
1	C	200/207 (96%)	-0.46	1 (0%)	91	89	10, 19, 37, 54	0
2	B	241/245 (98%)	-0.31	1 (0%)	92	90	10, 22, 42, 87	0
2	D	242/245 (98%)	-0.31	2 (0%)	86	84	10, 23, 47, 84	0
All	All	883/904 (97%)	-0.39	8 (0%)	84	82	10, 21, 42, 87	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	109	THR	3.6
2	D	253	ASP	3.0
1	C	29	TRP	2.9
2	B	253	ASP	2.8
2	D	108	GLY	2.6
1	A	30	TRP	2.5
1	A	29	TRP	2.3
1	A	108	VAL	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.