Full wwPDB X-ray Structure Validation Report (i)

May 14, 2020 – 03:13 am BST

PDB ID	:	2WPN
Title	:	Structure of the oxidised, as-isolated NiFeSe hydrogenase from D. vulgaris Hildenborough
Authors	:	Marques, M.C.; Coelho, R.; De Lacey, A.L.; Pereira, I.A.C.; Matias, P.M.
Deposited on	:	2009-08-07
Resolution	:	2.04 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	317	79% 8% 13%
2	B	496	87% 10% •

2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 6257 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PERIPLASMIC [NIFESE] HYDROGENASE, SMALL SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	277	2088	1332	348	390	18	0	1	0

- Molecule 2 is a protein called PERIPLASMIC [NIFESE] HYDROGENASE, LARGE SUBUNIT, SELENOCYSTEINE-CONTAINING.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
2	B	481	3750	2390	654	684	19	3	0	1	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	489	PSW	SEC	microheterogeneity	UNP Q72AS3

- Molecule 3 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe_4S_4).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Fe	S		
3	A	1	8	4	4	0	0
			Total	Fe	S		
3	A	1	8	4	4	0	0
			Total	Fe	S		
3	A	1	8	4	4	0	1

- Molecule 4 is BIS-(MU-2-OXO),[(MU-3--SULFIDO)-BIS(MU-2--SULFIDO)-TRIS(CYS-S)-TRI-IRON] (AQUA)(GLU-O)IRON(II) (three-letter code: FSX) (formula: $Fe_4O_3S_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	Fe	O	S		
4	A	1	10	4	3	3	0	1

- Molecule 5 is 3-[DODECYL(DIMETHYL)AMMONIO]PROPANE-1-SULFONATE (three-letter code: SBY) (formula: $C_{17}H_{37}NO_3S$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C 11 11	0	0
5	A	1	Total C 12 12	0	0
5	B	1	Total C 11 11	0	0
5	B	1	Total C 10 10	0	0

- Molecule 6 is CARBONMONOXIDE-(DICYANO) IRON (three-letter code: FCO) (formula: C_3FeN_2O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
6	B	1	7	3	1	2	1	0	0

- Molecule 7 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ni		
7	B	1	1	1	0	0

- Molecule 8 is FE (II) ION (three-letter code: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Fe		
8	B	1	1	1	0	0

- Molecule 9 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
9	B	1	1	1	0	0

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
10	A	138	138	138	0	0
			Total	O		
10	B	193	193	193	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PERIPLASMIC [NIFESE] HYDROGENASE, SMALL SUBUNIT

Chain A:



• Molecule 2: PERIPLASMIC [NIFESE] HYDROGENASE, LARGE SUBUNIT, SELENOCYSTEINE-CONTAINING

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	60.60Å 91.22Å 66.75Å 90.00° 101.73° 90.00°	Depositor
Resolution (Å)	65.36 – 2.04 19.57 – 2.04	Depositor EDS
% Data completeness (in resolution range)	97.6 (65.36-2.04) 97.7 (19.57-2.04)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.78 (at 2.04Å)	Xtriage
Refinement program	REFMAC 5.5.0088	Depositor
R, R_{free}	0.144 , 0.201 0.162 , 0.214	Depositor DCC
R_{free} test set	2223 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å^2)	23.8	Xtriage
Anisotropy	0.086	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 38.0	EDS
L-test for twinning[2]	$< \|L\| > = 0.45$, $< L^2 > = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	6257	wwPDB-VP
Average B, all atoms (Å^2)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.12% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, CL, SF4, SBY, SEC, FSX, FE2, OCS, PSW, FCO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.79	0/2146	0.74	0/2917
2	B	0.76	1/3806 (0.0%)	0.79	3/5142 (0.1%)
All	All	0.77	1/5952 (0.0%)	0.77	3/8059 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	316	VAL	CB-CG2	6.17	1.65	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	487	ASP	CB-CG-OD1	5.30	123.07	118.30
2	B	481	ARG	NE-CZ-NH2	-5.12	117.74	120.30
2	B	458	ARG	NE-CZ-NH1	5.04	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2088	0	2028	18	0
2	B	3750	0	3745	32	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	24	0	0	0	0
4	A	10	0	0	2	0
5	A	23	0	41	1	0
5	B	21	0	34	0	0
6	B	7	0	0	1	0
7	B	1	0	0	0	0
8	B	1	0	0	0	0
9	B	1	0	0	0	0
10	A	138	0	0	1	0
10	B	193	0	0	1	0
All	All	6257	0	5848	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:100:ASN:HD21	2:B:103:ARG:HH21	1.14	0.95
2:B:101:ASN:HD21	2:B:397:LYS:H	1.32	0.75
2:B:100:ASN:ND2	2:B:103:ARG:HH21	1.83	0.73
4:A:1287[B]:FSX:O1	4:A:1287[B]:FSX:S4	2.50	0.70
2:B:482:LEU:HD13	2:B:482:LEU:C	2.18	0.64
1:A:116:LEU:HD12	1:A:116:LEU:N	2.13	0.63
2:B:489[A]:PSW:SE	6:B:1497:FCO:C1	3.00	0.60
1:A:30:HIS:CE1	1:A:32:SER:HB2	2.37	0.59
4:A:1287[B]:FSX:S2	4:A:1287[B]:FSX:O1	2.60	0.59
2:B:491:GLY:HA3	10:B:2011:HOH:O	2.03	0.58
2:B:320:ARG:HH12	2:B:450:ASN:ND2	2.03	0.56
1:A:170:VAL:HG22	5:A:1289:SBY:H42	1.88	0.56
2:B:249:GLY:HA2	2:B:262:PHE:O	2.08	0.54
2:B:214:ARG:NH1	2:B:390:GLU:HG3	2.22	0.54
2:B:214:ARG:HG3	2:B:386:LEU:HD13	1.91	0.53
2:B:240:GLN:HG3	2:B:369:PHE:O	2.09	0.52
1:A:80:VAL:HG21	1:A:105:LEU:CD1	2.41	0.51
1:A:200:THR:N	1:A:201:PRO:CD	2.74	0.50
1:A:216:LYS:HB3	1:A:221:VAL:HG13	1.93	0.50
1:A:199:PRO:C	1:A:201:PRO:HD2	2.32	0.50
2:B:28:GLU:HB3	2:B:489[A]:PSW:CB	2.42	0.50
2:B:210:PHE:CE2	2:B:390:GLU:HG2	2.47	0.50
2:B:240:GLN:HA	2:B:369:PHE:O	2.12	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:80:VAL:HG21	1:A:105:LEU:HD12	1.94	0.49
2:B:432:LYS:HB2	2:B:437:ASP:HB2	1.96	0.47
2:B:422:ARG:HD2	2:B:489[B]:PSW:SD	2.55	0.46
1:A:8:PRO:HG2	1:A:41:ILE:HG22	1.98	0.46
2:B:319:PRO:HG3	2:B:446:LEU:HG	1.98	0.46
2:B:348:LEU:HD12	2:B:348:LEU:HA	1.79	0.44
2:B:359:LYS:O	2:B:362:ASP:HB2	2.19	0.43
1:A:127:ILE:HG12	1:A:274:PRO:HG3	2.00	0.43
1:A:48:THR:O	2:B:26:ARG:HA	2.17	0.43
2:B:28:GLU:HB3	2:B:489[B]:PSW:SE	2.68	0.43
2:B:197:ILE:HD12	2:B:401:ASP:HA	2.00	0.43
1:A:106:LYS:HE2	10:A:2078:HOH:O	2.18	0.42
1:A:188:GLU:O	1:A:191:LYS:HB2	2.19	0.42
1:A:200:THR:N	1:A:201:PRO:HD2	2.34	0.42
2:B:449:CYS:SG	2:B:488:PRO:HB3	2.60	0.42
2:B:188:GLY:HA2	2:B:195:THR:OG1	2.19	0.42
2:B:482:LEU:CD1	2:B:482:LEU:C	2.89	0.41
1:A:36:VAL:HA	1:A:40:VAL:HB	2.03	0.41
2:B:100:ASN:HD22	2:B:255:ASN:HD21	1.69	0.41
2:B:28:GLU:HB3	2:B:489[B]:PSW:HA	2.01	0.41
2:B:210:PHE:HE2	2:B:390:GLU:HG2	1.86	0.41
1:A:258:TRP:CH2	1:A:260:VAL:HB	2.56	0.41
2:B:480:ALA:O	2:B:484:ARG:HG3	2.21	0.41
2:B:74:ILE:HD11	2:B:441:ILE:HD13	2.03	0.41
1:A:52:TRP:CE2	1:A:56:HIS:HB3	2.57	0.40
1:A:40:VAL:HG12	1:A:41:ILE:HG23	2.03	0.40
2:B:103:ARG:HE	2:B:255:ASN:ND2	2.19	0.40
2:B:26:ARG:HD2	2:B:125:LEU:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	276/317 (87%)	268 (97%)	7 (2%)	1 (0%)	34	24
2	B	477/496 (96%)	463 (97%)	14 (3%)	0	100	100
All	All	753/813 (93%)	731 (97%)	21 (3%)	1 (0%)	51	45

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	263	ALA

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/253 (88%)	221 (99%)	2 (1%)	78	79
2	B	390/398 (98%)	382 (98%)	8 (2%)	53	48
All	All	613/651 (94%)	603 (98%)	10 (2%)	62	59

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	52	TRP
1	A	261	GLN
2	B	16	ARG
2	B	80	THR
2	B	116	GLN
2	B	162	TYR
2	B	224	VAL
2	B	273	MET
2	B	348	LEU
2	B	403	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	180	ASN
2	B	100	ASN
2	B	101	ASN
2	B	185	HIS
2	B	255	ASN
2	B	450	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	OCS	B	75	2,7	3,7,9	1.13	0	1,8,13	2.01	1 (100%)
2	PSW	B	489[A]	2,7	3,6,7	0.89	0	0,6,8	0.00	-
2	PSW	B	489[B]	2,7	3,6,7	0.82	0	0,6,8	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OCS	B	75	2,7	-	0/2/6/9	-
2	PSW	B	489[A]	2,7	-	0/0/5/7	-
2	PSW	B	489[B]	2,7	-	0/0/5/7	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	75	OCS	OD1-SG-CB	2.01	109.36	105.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	489[A]	PSW	2	0
2	B	489[B]	PSW	3	0

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 3 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	SBY	A	1288	-	10,10,21	0.38	0	9,9,26	0.55	0
4	FSX	A	1287[B]	1	0,14,14	0.00	-	-		
3	SF4	A	1284	1	0,12,12	0.00	-	-		
6	FCO	B	1497	2,7	0,6,6	0.00	-	-		
3	SF4	A	1286[A]	1	0,12,12	0.00	-	-		
5	SBY	B	1502	-	9,9,21	0.46	0	8,8,26	0.40	0
5	SBY	A	1289	-	11,11,21	0.41	0	10,10,26	0.41	0
5	SBY	B	1501	-	10,10,21	0.39	0	9,9,26	0.39	0
3	SF4	A	1285	1	0,12,12	0.00	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	SBY	A	1288	-	-	2/8/8/21	-
3	SF4	A	1286[A]	1	-	-	0/6/5/5
3	SF4	A	1284	1	-	-	0/6/5/5
4	FSX	A	1287[B]	1	-	-	0/4/5/5
5	SBY	B	1502	-	-	7/7/7/21	-
5	SBY	A	1289	-	-	2/9/9/21	-
5	SBY	B	1501	-	-	2/8/8/21	-
3	SF4	A	1285	1	-	-	0/6/5/5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1288	SBY	C3-C4-C5-C6
5	B	1502	SBY	C7-C8-C9-C10
5	A	1288	SBY	C2-C3-C4-C5
5	B	1502	SBY	C11-C10-C9-C8
5	B	1502	SBY	C4-C5-C6-C7
5	B	1502	SBY	C9-C10-C11-C12
5	A	1289	SBY	C5-C6-C7-C8
5	A	1289	SBY	C1-C2-C3-C4
5	B	1501	SBY	C7-C8-C9-C10
5	B	1502	SBY	C6-C7-C8-C9
5	B	1502	SBY	C5-C6-C7-C8
5	B	1502	SBY	C3-C4-C5-C6
5	B	1501	SBY	C5-C6-C7-C8

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1287[B]	FSX	2	0
6	B	1497	FCO	1	0
5	A	1289	SBY	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand SBY A 1288



Ligand SBY B 1502



Ligand SBY A 1289



Ligand SBY B 1501



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	277/317 (87%)	-0.25	1 (0%)	92	93	15, 20, 26, 33	5 (1%)
2	B	479/496 (96%)	-0.40	1 (0%)	95	95	13, 20, 32, 50	5 (1%)
All	All	756/813 (92%)	-0.35	2 (0%)	94	94	13, 20, 30, 50	10 (1%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	283	ALA	2.7
2	B	256	SER	2.4

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	SEC	B	489[C]	6/7	0.95	0.12	10,18,22,23	6
2	PSW	B	489[B]	7/8	0.98	0.08	15,22,23,82	7
2	PSW	B	489[A]	7/8	0.98	0.08	15,20,23,24	7
2	OCS	B	75	8/10	0.99	0.07	16,20,23,23	0

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	SBY	B	1502	10/22	0.87	0.15	34,40,58,58	0
5	SBY	A	1289	12/22	0.87	0.14	32,47,48,61	0
5	SBY	B	1501	11/22	0.88	0.16	28,35,38,47	0
5	SBY	A	1288	11/22	0.95	0.13	32,38,45,48	0
3	SF4	A	1284	8/8	0.96	0.04	16,20,23,24	0
3	SF4	A	1285	8/8	0.97	0.06	17,18,22,23	0
3	SF4	A	1286[A]	8/8	0.98	0.04	17,20,22,24	8
7	NI	B	1498	1/1	0.98	0.03	24,24,24,24	0
9	CL	B	1500	1/1	0.99	0.07	25,25,25,25	0
8	FE2	B	1499	1/1	0.99	0.07	27,27,27,27	1
6	FCO	B	1497	7/7	0.99	0.08	15,19,23,28	0
4	FSX	A	1287[B]	10/10	0.99	0.05	14,21,25,25	10

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around SBY B 1502:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around SBY A 1289:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around SBY B 1501:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around SBY A 1288:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.