

# Full wwPDB NMR Structure Validation Report (i)

#### Aug 21, 2022 – 08:01 AM EDT

PDB ID : 1WTU

Title: TRANSCRIPTION FACTOR 1, NMR, MINIMIZED AVERAGE STRUC-

TURE

Authors: Jia, X.; Grove, A.; Ivancic, M.; Hsu, V.L.; Geiduschek, E.P.; Kearns, D.R.

Deposited on : 1996-07-29

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

ShiftChecker : 2.29

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.29

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $SOLUTION\ NMR$ 

The overall completeness of chemical shifts assignment was not calculated.

There are no overall percentile quality scores available for this entry.

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	99	100%
1	В	99	100%



# 2 Ensemble composition and analysis (i)

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.



# 3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 3096 atoms, of which 1588 are hydrogens and 0 are deuteriums.

 $\bullet$  Molecule 1 is a protein called TRANSCRIPTION FACTOR 1.

Mol	Chain	Residues		Atoms							
1	Λ	99	Total	С	Н	N	О	S	0		
1	А	99	1548	477	794	127	148	2			
1	D	99	Total	С	Н	N	О	S	0		
1	Б	99	1548	477	794	127	148	2			



# 4 Residue-property plots (i)

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: TRANSCRIPTION FACTOR 1

Chain	A:														10	0%																				
M1 N2 K3 T4 E5	L6 17 K8	A9 110	Q12 D13	T14	L16	T17 Q18	V19	V21	\$22 K23	M24	L25 A26	S27	F28	K30	131	T32	E34	T35	V36	K38	G39	D40 K41	V42	043	T45	G46	r4/ 1.48	N49	150	N51	V53	A54	R55	A57	R58	099 090
F61 N62 P63 Q64 T65	066 E67 A68	E70	A72 P73	S74	676 G76	V77 S78	677	P81	G82 E83	S84	L85 K86	K87	A88	H89	G9.1	L92 K93	Y94	E95	D96	A98	K99															
• Mole	ecule	1:	TR	kA.	NS	SC.	RI	P.	ΓΙ	IC	N	FA	40	СΊ	Ю	R	1																			
Chain	В:														10	0%																				
M1 N2 K3 T4 E5	L6 17 K8	110	Q12 D13	T14	E15 L16	T17 Q18	V19	V21	\$22 K23	M24	L25 A26	S27	F28	K30	131	T32	E34	T35	V36	K38	G39	D40 K41	V42	Q43	T45	G46	1.48	N49	150	N51	V53	A54	R55	A57	R58	K59
F61 N62 P63 Q64 T65	Q66 E67 A68	E70	A72 P73	S74	076 G76	V77 S78	67V	n80 P81	G82 E83	S84	L85	K87	A88	A89 E90	G91	L92 K93	Y94	E95	D96 F97	A98	K99															



#### Refinement protocol and experimental data overview (i) 5



Of the ? calculated structures, 1 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	

No chemical shift data was provided.



## 6 Model quality (i)

## 6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

#### 6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	0	0	0	0
1	В	0	0	0	0
All	All	0	0	0	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

## 6.3 Torsion angles (i)

#### 6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	0	-	-	-	-
1	В	0	-	-	-	-
All	All	0	-	-	-	-

There are no Ramachandran outliers.



#### 6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	0	-	-	-
1	В	0	-	-	-
All	All	0	-	-	-

There are no protein residues with a non-rotameric sidechain to report.

#### 6.3.3 RNA (i)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.6 Ligand geometry (i)

There are no ligands in this entry.

## 6.7 Other polymers (i)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 7 Chemical shift validation (i)

No chemical shift data were provided

