



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 11:38 am BST

PDB ID : 3WTC
Title : Crystal structure of Gox2036
Authors : Yuan, Y.A.; Lin, J.P.
Deposited on : 2014-04-09
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	262	
1	B	262	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 4163 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative oxidoreductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	259	1898	1197	321	371	9	0	0	0
1	B	259	1898	1197	321	371	9	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP Q5FPC4
A	-1	SER	-	EXPRESSION TAG	UNP Q5FPC4
A	0	HIS	-	EXPRESSION TAG	UNP Q5FPC4
B	-2	GLY	-	EXPRESSION TAG	UNP Q5FPC4
B	-1	SER	-	EXPRESSION TAG	UNP Q5FPC4
B	0	HIS	-	EXPRESSION TAG	UNP Q5FPC4

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	192	Total 192	O 192	0	0
2	B	175	Total 175	O 175	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	117.84Å 58.10Å 67.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.22 – 1.65 35.24 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.8 (41.22-1.65) 99.8 (35.24-1.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.60 (at 1.65Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.148 , 0.174 0.162 , 0.189	Depositor DCC
R_{free} test set	2847 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	17.5	Xtrriage
Anisotropy	0.062	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4163	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.05	2/1919 (0.1%)	1.52	8/2594 (0.3%)
1	B	1.00	4/1919 (0.2%)	1.10	11/2594 (0.4%)
All	All	1.02	6/3838 (0.2%)	1.33	19/5188 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	24	ARG	CZ-NH2	-9.06	1.21	1.33
1	A	24	ARG	CD-NE	-6.43	1.35	1.46
1	B	71	SER	CB-OG	-6.34	1.34	1.42
1	B	240	SER	CB-OG	-5.39	1.35	1.42
1	B	225	GLU	CD-OE2	5.28	1.31	1.25
1	B	105	GLU	CD-OE1	5.07	1.31	1.25

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	24	ARG	NE-CZ-NH2	-39.87	100.36	120.30
1	A	24	ARG	NE-CZ-NH1	33.92	137.26	120.30
1	A	163	ARG	NE-CZ-NH1	15.65	128.13	120.30
1	B	54	ARG	NE-CZ-NH2	-12.62	113.99	120.30
1	A	163	ARG	NE-CZ-NH2	-11.34	114.63	120.30
1	A	24	ARG	CD-NE-CZ	10.25	137.95	123.60
1	A	121	MET	CG-SD-CE	-9.90	84.36	100.20
1	B	54	ARG	CG-CD-NE	-8.04	94.91	111.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	54	ARG	NE-CZ-NH1	7.52	124.06	120.30
1	B	197	ASP	CB-CG-OD2	7.41	124.97	118.30
1	A	163	ARG	CD-NE-CZ	6.88	133.24	123.60
1	A	55	ARG	NE-CZ-NH2	-6.24	117.18	120.30
1	B	69	PHE	CB-CG-CD1	5.52	124.66	120.80
1	B	42	LEU	CA-CB-CG	5.37	127.65	115.30
1	B	66	ARG	CD-NE-CZ	5.36	131.10	123.60
1	B	228	ASP	CB-CG-OD1	5.25	123.03	118.30
1	B	243	ASP	CB-CG-OD1	5.19	122.97	118.30
1	B	223	ARG	NE-CZ-NH1	5.12	122.86	120.30
1	B	67	ASP	CB-CG-OD1	5.07	122.86	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	24	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1898	0	1950	16	0
1	B	1898	0	1950	8	0
2	A	192	0	0	3	1
2	B	175	0	0	3	1
All	All	4163	0	3900	24	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:GLY:O	1:A:163:ARG:CD	2.14	0.96

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:GLY:O	1:A:163:ARG:HD3	1.69	0.90
1:A:62:ASP:H	1:A:68:GLN:HE22	1.24	0.82
1:A:133:THR:HG21	2:A:355:HOH:O	1.80	0.82
1:B:208:ILE:HG22	2:B:455:HOH:O	1.83	0.78
1:A:145:GLY:O	1:A:163:ARG:HD2	1.87	0.74
1:B:15:GLN:HG2	2:B:356:HOH:O	1.94	0.68
1:B:181:ASN:HD22	1:B:248:GLN:H	1.45	0.65
1:A:62:ASP:H	1:A:68:GLN:NE2	1.94	0.64
1:A:181:ASN:HD22	1:A:248:GLN:H	1.46	0.62
1:B:95:VAL:O	1:B:199:ARG:NH1	2.34	0.60
1:A:141:CYS:SG	1:A:163:ARG:HG3	2.42	0.59
1:B:1:MET:N	2:B:302:HOH:O	2.43	0.51
1:A:190:THR:HG21	1:A:212:TYR:OH	2.12	0.50
1:A:131:LYS:HG2	1:A:133:THR:HG22	1.95	0.48
1:A:38:LYS:HG2	1:A:40:ASP:OD1	2.13	0.48
1:A:25:LEU:HD23	2:A:464:HOH:O	2.16	0.45
1:B:88:ASN:HD21	1:B:120:GLY:HA3	1.83	0.43
1:A:190:THR:CG2	1:A:212:TYR:OH	2.67	0.43
1:A:88:ASN:HD21	1:A:120:GLY:HA3	1.84	0.41
1:B:197:ASP:HB2	1:B:212:TYR:HB2	2.02	0.41
1:A:3:LEU:HD11	2:A:464:HOH:O	2.21	0.40
1:B:36:ASP:O	1:B:60:THR:HA	2.20	0.40
1:A:128:PHE:HA	1:A:133:THR:HG23	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:491:HOH:O	2:B:453:HOH:O[3_656]	1.23	0.97

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/262 (98%)	250 (97%)	7 (3%)	0	100	100
1	B	257/262 (98%)	252 (98%)	5 (2%)	0	100	100
All	All	514/524 (98%)	502 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/196 (99%)	186 (96%)	8 (4%)	30	8
1	B	194/196 (99%)	191 (98%)	3 (2%)	65	44
All	All	388/392 (99%)	377 (97%)	11 (3%)	43	18

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	24	ARG
1	A	42	LEU
1	A	69	PHE
1	A	133	THR
1	A	151	LEU
1	A	203	ILE
1	A	213	LYS
1	B	42	LEU
1	B	69	PHE
1	B	192	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	68	GLN
1	A	88	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	181	ASN
1	A	248	GLN
1	B	88	ASN
1	B	89	ASN
1	B	181	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	259/262 (98%)	0.07	22 (8%) 10 10	10, 17, 44, 71	0
1	B	259/262 (98%)	-0.11	13 (5%) 28 27	11, 18, 42, 71	0
All	All	518/524 (98%)	-0.02	35 (6%) 17 16	10, 18, 42, 71	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	208	ILE	8.0
1	A	206	THR	7.6
1	A	203	ILE	5.2
1	B	192	MET	5.1
1	A	211	THR	4.9
1	A	194	VAL	4.5
1	A	199	ARG	4.5
1	A	209	GLY	4.5
1	B	191	ASP	4.3
1	B	194	VAL	4.0
1	A	193	TRP	3.7
1	A	207	GLU	3.6
1	A	205	GLY	3.5
1	A	191	ASP	3.5
1	A	195	THR	3.4
1	B	195	THR	3.4
1	A	204	THR	3.3
1	A	210	ALA	3.2
1	B	189	GLY	3.1
1	B	198	LYS	3.1
1	A	189	GLY	3.1
1	A	190	THR	3.0
1	B	204	THR	2.8
1	A	201	ALA	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	202	GLU	2.6
1	A	200	MET	2.5
1	B	203	ILE	2.5
1	A	197	ASP	2.4
1	B	206	THR	2.2
1	B	205	GLY	2.2
1	B	199	ARG	2.1
1	B	215	TYR	2.1
1	A	213	LYS	2.1
1	B	190	THR	2.0
1	A	198	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.