

# Full wwPDB X-ray Structure Validation Report (i)

May 15, 2020 – 07:38 pm BST

PDB ID : 1WVX

Title : Crystal structures of kinase domain of DAP kinase in complex with small

molecular inhibitors

Authors: Ueda, Y.; Ogata, H.; Yamakawa, A.; Higuchi, Y.

Deposited on : 2004-12-27

Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED

EDS: NOT EXECUTED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mo	l Chain	Length	Quality of chain				
1	A	278	72%	24%			



## 2 Entry composition (i)


There are 3 unique types of molecules in this entry. The entry contains 2329 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Death-associated protein kinase 1.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	275	Total	С	N	О	S	0	0	0
1	Λ	210	2231	1433	371	423	4	0	0	

• Molecule 2 is 6-(3-AMINOPROPYL)-4,9-DIMETHYLPYRROLO[3,4-C]CARBAZOLE-1,3(2H,6H)-DIONE (three-letter code: BD4) (formula: C<sub>19</sub>H<sub>19</sub>N<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 24	C 19	N 3	O 2	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	74	Total O 74 74	0	0




## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Death-associated protein kinase 1





# 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	P 21 21 21	Depositor	
Cell constants	77.34Å 108.87Å 50.19Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor	
Resolution (Å)	10.00 - 2.60	Depositor	
% Data completeness	89.8 (10.00-2.60)	Depositor	
(in resolution range)	03.0 (10.00-2.00)	Depositor	
$R_{merge}$	0.09	Depositor	
$R_{sym}$	(Not available)	Depositor	
Refinement program	CNS 1.1	Depositor	
$R, R_{free}$	0.225 , $0.270$	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	2329	wwPDB-VP	
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP	



# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BD4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

1	Tal	Chain	Bond	lengths	Bond angles		
10.	101	l Chain	RMSZ	# Z >5	RMSZ	# Z  > 5	
	1	A	0.39	0/2276	0.60	0/3073	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2231	0	2233	55	0
2	A	24	0	19	1	0
3	A	74	0	0	0	0
All	All	2329	0	2252	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	$egin{array}{c}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{array}$
1:A:85:ASN:HD21	1:A:88:ASP:H	1.07	0.96

Continued on next page...



Continued from previous page...

Continued from prev		Interatomic	Clash
Atom-1	Atom-2	${ m distance}({ m \AA})$	overlap (Å)
1:A:85:ASN:ND2	1:A:88:ASP:H	1.68	0.91
1:A:120:LEU:HD21	1:A:207:THR:HG22	1.54	0.90
1:A:8:ASN:HD21	1:A:10:ASP:HB2	1.46	0.80
1:A:168:ILE:HD12	1:A:196:LEU:HD13	1.69	0.72
1:A:85:ASN:C	1:A:85:ASN:HD22	1.94	0.71
1:A:112:THR:HG23	1:A:115:GLU:H	1.53	0.70
1:A:8:ASN:ND2	1:A:10:ASP:HB2	2.06	0.69
1:A:203:ILE:O	1:A:207:THR:HG23	1.97	0.65
1:A:120:LEU:CD2	1:A:207:THR:HG22	2.25	0.64
2:A:1001:BD4:H6	2:A:1001:BD4:O24	1.98	0.63
1:A:85:ASN:HD21	1:A:88:ASP:N	1.89	0.63
1:A:77:ILE:HD11	1:A:94:GLU:O	1.99	0.63
1:A:118:GLU:OE2	1:A:154:LYS:HD2	1.99	0.61
1:A:71:ILE:HG21	1:A:76:VAL:HG11	1.83	0.60
1:A:47:ARG:O	1:A:49:THR:HG22	2.04	0.57
1:A:112:THR:HG22	1:A:115:GLU:OE1	2.04	0.56
1:A:212:SER:HB2	1:A:240:TYR:HB3	1.87	0.55
1:A:192:GLU:HB3	1:A:193:PRO:HD2	1.90	0.53
1:A:177:ILE:HG23	1:A:177:ILE:O	2.09	0.52
1:A:82:VAL:HG13	1:A:91:LEU:HD23	1.93	0.51
1:A:49:THR:HG23	1:A:52:SER:HB3	1.92	0.51
1:A:71:ILE:HG21	1:A:76:VAL:CG1	2.41	0.51
1:A:85:ASN:ND2	1:A:87:THR:H	2.09	0.51
1:A:231:ALA:O	1:A:232:VAL:HG12	2.12	0.50
1:A:112:THR:CG2	1:A:115:GLU:HG3	2.42	0.50
1:A:15:THR:CG2	1:A:28:LYS:HD3	2.42	0.49
1:A:58:ARG:HG2	1:A:58:ARG:HH21	1.77	0.49
1:A:73:HIS:HE1	1:A:75:ASN:HD22	1.61	0.49
1:A:157:ILE:HG13	1:A:157:ILE:O	2.13	0.48
1:A:168:ILE:HD13	1:A:195:GLY:HA2	1.96	0.48
1:A:73:HIS:ND1	1:A:74:PRO:HD2	2.28	0.47
1:A:185:ALA:HB1	1:A:186:PRO:HD2	1.97	0.47
1:A:102:PHE:HE2	1:A:209:ILE:HG22	1.79	0.47
1:A:112:THR:HG22	1:A:115:GLU:CG	2.44	0.47
1:A:146:MET:HG3	1:A:160:ILE:HD13	1.98	0.46
1:A:4:PHE:HB2	1:A:84:GLU:HG2	1.97	0.46
1:A:73:HIS:CE1	1:A:74:PRO:HD2	2.50	0.46
1:A:232:VAL:HG22	1:A:232:VAL:O	2.16	0.45
1:A:181:PRO:HB3	1:A:226:LEU:HD21	1.99	0.45
1:A:90:ILE:N	1:A:90:ILE:HD12	2.32	0.45
1:A:272:HIS:CG	1:A:273:PRO:HD2	2.51	0.44

Continued on next page...



 $Continued\ from\ previous\ page...$ 

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${f distance} ({f \AA})$	overlap (Å)
1:A:27:VAL:HA	1:A:41:ALA:O	2.17	0.44
1:A:112:THR:HG22	1:A:115:GLU:HG3	1.99	0.43
1:A:73:HIS:CG	1:A:74:PRO:HD2	2.54	0.42
1:A:178:PHE:HB3	1:A:188:ILE:HG21	2.00	0.42
1:A:234:TYR:O	1:A:235:GLU:HG3	2.20	0.42
1:A:85:ASN:C	1:A:85:ASN:ND2	2.67	0.42
1:A:164:LEU:O	1:A:166:HIS:HD2	2.03	0.42
1:A:185:ALA:HB1	1:A:187:GLU:OE1	2.21	0.41
1:A:46:LYS:HG2	1:A:89:VAL:HG23	2.02	0.41
1:A:129:TYR:O	1:A:133:LEU:HD23	2.19	0.41
1:A:145:ILE:HG23	1:A:157:ILE:HD13	2.02	0.41
1:A:180:THR:HA	1:A:181:PRO:HD3	1.87	0.41
1:A:270:LEU:HD13	1:A:270:LEU:HA	1.76	0.41
1:A:181:PRO:HB3	1:A:226:LEU:CD2	2.51	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	273/278 (98%)	251 (92%)	18 (7%)	4 (2%)	10 21

#### All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	177	ILE
1	A	17	GLU
1	A	49	THR
1	A	161	ASP



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/248 (99%)	232 (95%)	13 (5%)	22 45	

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	LYS
1	A	49	THR
1	A	58	ARG
1	A	62	GLU
1	A	85	ASN
1	A	148	LEU
1	A	156	ARG
1	A	157	ILE
1	A	161	ASP
1	A	170	PHE
1	A	180	THR
1	A	270	LEU
1	A	271	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	GLN
1	A	8	ASN
1	A	75	ASN
1	A	80	HIS
1	A	85	ASN
1	A	166	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

#### 5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res Link		Bo	ond leng	hs	В	ond ang	les
IVIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BD4	A	1001	-	23,27,27	1.29	3 (13%)	21,41,41	1.44	5 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	${f Res}$	Link	Chirals	Torsions	Rings
2	BD4	A	1001	-	-	0/4/4/4	0/4/4/4

#### All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(\mathbf{\mathring{A}})$	$\operatorname{Ideal}( ext{\AA})$
2	A	1001	BD4	C11-C12	-2.81	1.39	1.44
2	A	1001	BD4	C9-C8	-2.58	1.39	1.42
2	A	1001	BD4	C22-C13	-2.03	1.39	1.43

#### All (5) bond angle outliers are listed below:

Mol	Chain	$\operatorname{Res}$	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\mathbf{Ideal}(^o)$
2	A	1001	BD4	C1-C6-C5	-3.35	118.09	122.11

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	${f Atoms}$	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
2	A	1001	BD4	C6-C5-C9	-2.72	129.85	134.24
2	A	1001	BD4	C15-C11-C12	-2.22	120.29	124.37
2	A	1001	BD4	C9-C5-C4	2.16	108.73	106.37
2	A	1001	BD4	C5-C9-C8	2.05	108.33	106.09

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	BD4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

