



Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 05:25 pm BST

PDB ID : 3WVI
Title : Time-Resolved Crystal Structure of HindIII with 40 sec soaking
Authors : Kawamura, T.; Kobayashi, T.; Watanabe, N.
Deposited on : 2014-05-21
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	
1	B	300	
1	C	300	
1	D	300	
2	E	12	
2	F	12	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	12	 83% 17%
2	H	12	 92% 8%
2	I	12	 75% 25%
2	J	12	 83% 8% 8%
2	K	12	 100%
2	L	12	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	B	302	-	X	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 12168 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Type-2 restriction enzyme HindIII.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	298	Total 2458	C 1584	N 406	O 465	S 3	0	1	0
1	B	298	Total 2452	C 1580	N 405	O 464	S 3	0	0	0
1	C	298	Total 2452	C 1580	N 405	O 464	S 3	0	0	0
1	D	298	Total 2452	C 1580	N 405	O 464	S 3	0	0	0


- Molecule 2 is a DNA chain called DNA (5'-D(*GP*CP*CP*AP*AP*GP*CP*TP*TP*GP*GP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	F	12	Total 243	C 116	N 46	O 70	P 11	0	0	0
2	E	12	Total 243	C 116	N 46	O 70	P 11	0	0	0
2	H	12	Total 243	C 116	N 46	O 70	P 11	0	0	0
2	G	12	Total 243	C 116	N 46	O 70	P 11	0	0	0
2	K	12	Total 243	C 116	N 46	O 70	P 11	0	0	0
2	L	12	Total 243	C 116	N 46	O 70	P 11	0	0	0
2	I	12	Total 243	C 116	N 46	O 70	P 11	0	0	0
2	J	12	Total 243	C 116	N 46	O 70	P 11	0	0	0

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	G	1	Total Mn 1 1	0	0
3	D	2	Total Mn 2 2	0	0
3	E	1	Total Mn 1 1	0	0
3	B	1	Total Mn 1 1	0	0
3	C	1	Total Mn 1 1	0	0
3	A	1	Total Mn 1 1	0	0
3	F	1	Total Mn 1 1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	87	Total O 87 87	0	0
5	B	84	Total O 84 84	0	0
5	C	68	Total O 68 68	0	0
5	D	78	Total O 78 78	0	0
5	F	11	Total O 11 11	0	0
5	E	19	Total O 19 19	0	0
5	H	15	Total O 15 15	0	0
5	G	16	Total O 16 16	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Type-2 restriction enzyme HindIII




- Molecule 1: Type-2 restriction enzyme HindIII




- Molecule 1: Type-2 restriction enzyme HindIII




- Molecule 1: Type-2 restriction enzyme HindIII





- Molecule 2: DNA (5'-D(*GP*CP*CP*AP*AP*GP*CP*TP*TP*GP*GP*C)-3')

Chain F: 75% 25%



- Molecule 2: DNA (5'-D(*GP*CP*CP*AP*AP*GP*CP*TP*TP*GP*GP*C)-3')

Chain E: 92% 8%



- Molecule 2: DNA (5'-D(*GP*CP*CP*AP*AP*GP*CP*TP*TP*GP*GP*C)-3')

Chain H: 92% 8%



- Molecule 2: DNA (5'-D(*GP*CP*CP*AP*AP*GP*CP*TP*TP*GP*GP*C)-3')

Chain G: 83% 17%



- Molecule 2: DNA (5'-D(*GP*CP*CP*AP*AP*GP*CP*TP*TP*GP*GP*C)-3')

Chain K: 100%

There are no outlier residues recorded for this chain.


- Molecule 2: DNA (5'-D(*GP*CP*CP*AP*AP*GP*CP*TP*TP*GP*GP*C)-3')

Chain L: 100%


There are no outlier residues recorded for this chain.

- Molecule 2: DNA (5'-D(*GP*CP*CP*AP*AP*GP*CP*TP*TP*GP*GP*C)-3')

Chain I: 75% 25%



- Molecule 2: DNA (5'-D(*GP*CP*CP*AP*AP*GP*CP*TP*TP*GP*GP*C)-3')

Chain J:  83% 8% 8%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.61Å 143.24Å 94.36Å 90.00° 113.44° 90.00°	Depositor
Resolution (Å)	32.23 – 2.55 32.23 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.7 (32.23-2.55) 99.7 (32.23-2.55)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 2.54Å)	Xtrriage
Refinement program	REFMAC 5.8.0069	Depositor
R, R_{free}	0.173 , 0.234 0.181 , 0.236	Depositor DCC
R_{free} test set	3249 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	29.7	Xtrriage
Anisotropy	0.195	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.024 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12168	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.73	0/2503	0.80	0/3363
1	B	0.74	0/2494	0.81	3/3351 (0.1%)
1	C	0.70	0/2494	0.82	2/3351 (0.1%)
1	D	0.71	0/2494	0.81	0/3351
2	E	0.67	0/272	0.96	0/418
2	F	0.63	0/272	0.87	0/418
2	G	0.63	0/272	0.86	0/418
2	H	0.60	0/272	1.28	2/418 (0.5%)
2	I	0.66	0/272	0.94	0/418
2	J	0.57	0/272	1.06	1/418 (0.2%)
2	K	0.60	0/272	0.99	0/418
2	L	0.56	0/272	0.97	0/418
All	All	0.70	0/12161	0.85	8/16760 (0.0%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	4	DA	O5'-P-OP2	12.68	125.92	110.70
2	H	4	DA	O5'-P-OP1	-12.54	94.42	105.70
1	C	113	ARG	NE-CZ-NH1	8.31	124.46	120.30
1	B	36	ARG	NE-CZ-NH1	6.23	123.41	120.30
1	C	113	ARG	NE-CZ-NH2	-6.03	117.28	120.30
1	B	133	ARG	NE-CZ-NH2	-5.64	117.48	120.30
1	B	36	ARG	NE-CZ-NH2	-5.59	117.50	120.30
2	J	6	DG	C1'-O4'-C4'	-5.21	104.89	110.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2458	0	2510	18	0
1	B	2452	0	2502	21	0
1	C	2452	0	2502	12	0
1	D	2452	0	2502	14	0
2	E	243	0	136	1	0
2	F	243	0	136	2	0
2	G	243	0	136	1	0
2	H	243	0	136	0	0
2	I	243	0	136	2	0
2	J	243	0	136	1	0
2	K	243	0	136	0	0
2	L	243	0	136	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	2	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
4	B	12	0	16	4	0
4	C	12	0	16	0	0
5	A	87	0	0	2	0
5	B	84	0	0	3	0
5	C	68	0	0	0	0
5	D	78	0	0	4	0
5	E	19	0	0	2	0
5	F	11	0	0	0	0
5	G	16	0	0	0	0
5	H	15	0	0	0	0
All	All	12168	0	11136	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:60:GLU:OE2	1:C:113:ARG:NH2	2.09	0.85
1:D:87:GLN:O	5:D:466:HOH:O	2.00	0.78
1:B:59:GLN:HG3	5:B:466:HOH:O	1.85	0.75
1:B:146:PHE:HA	4:B:302:GOL:H11	1.70	0.74
1:B:145:PHE:O	4:B:302:GOL:H12	1.88	0.73
1:B:148:TYR:O	4:B:302:GOL:O1	2.07	0.72
1:A:252:TRP:CE3	1:B:277:MET:HG2	2.30	0.66
2:J:5:DA:H2''	2:J:6:DG:OP2	1.94	0.65
1:C:259:ILE:HG23	1:D:271:LEU:HD23	1.79	0.65
1:D:235:GLN:HG3	5:D:436:HOH:O	2.00	0.61
1:D:253:LYS:HB2	5:D:445:HOH:O	2.02	0.60
1:C:299:LEU:O	1:D:213:ARG:NH1	2.37	0.57
1:B:146:PHE:HA	4:B:302:GOL:C1	2.33	0.57
1:B:226:ASP:OD1	1:B:229:THR:HG23	2.07	0.55
1:D:110:LYS:HE2	5:D:468:HOH:O	2.07	0.54
2:F:5:DA:H2'	2:F:6:DG:O4'	2.09	0.52
1:A:110:LYS:HE3	1:A:124:PHE:HA	1.90	0.52
2:E:12:DC:OP2	2:I:1:DG:O5'	2.25	0.52
1:D:263:THR:O	1:D:265:GLU:N	2.44	0.51
1:B:107:ALA:HA	1:B:140:VAL:O	2.11	0.51
1:C:253:LYS:O	1:C:256:ILE:HG22	2.11	0.50
1:A:20:PHE:CD2	2:F:11:DG:H5''	2.46	0.50
2:G:5:DA:H2'	2:G:6:DG:O4'	2.11	0.50
1:A:110:LYS:HE2	5:E:204:HOH:O	2.13	0.49
1:D:263:THR:HG23	1:D:266:GLU:OE1	2.14	0.47
1:C:230:LEU:HD23	1:D:299:LEU:HD21	1.96	0.47
1:B:105:LEU:C	1:B:105:LEU:HD12	2.35	0.47
1:C:259:ILE:CG2	1:D:271:LEU:HD23	2.44	0.47
1:C:53:ILE:O	1:C:113:ARG:NH1	2.47	0.47
2:I:4:DA:H1'	2:I:5:DA:C8	2.50	0.47
1:B:106:VAL:HG21	1:B:133:ARG:HA	1.97	0.46
1:B:288:ILE:HG22	1:B:292:LYS:HE2	1.97	0.46
1:D:95:ILE:HD12	1:D:136:LYS:HE3	1.97	0.46
1:A:157:LYS:HD3	5:A:450:HOH:O	2.16	0.45
1:B:143:ALA:O	1:B:168:SER:HA	2.15	0.45
1:C:44:ARG:HD3	1:C:237:GLU:OE2	2.17	0.45
1:A:160:LEU:HB3	1:A:199:SER:HB3	1.98	0.44
1:A:24:THR:O	1:A:28:ILE:HG13	2.17	0.44
1:B:272:LEU:HD23	1:B:277:MET:HE1	2.00	0.44
1:C:156:PHE:O	1:C:160:LEU:HG	2.18	0.43
1:A:234:LEU:HD23	1:A:234:LEU:HA	1.83	0.43
1:A:89:GLY:HA3	5:E:209:HOH:O	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:46:ILE:HG12	1:A:67:GLY:HA2	2.01	0.43
1:C:213:ARG:NH1	1:D:299:LEU:O	2.52	0.43
1:B:42:LEU:HD12	1:B:42:LEU:HA	1.65	0.42
1:A:194:PHE:N	1:A:195:PRO:CD	2.82	0.42
1:A:143:ALA:O	1:A:168:SER:HA	2.18	0.42
1:C:252:TRP:CE2	1:D:280:LYS:HD2	2.54	0.42
1:B:276:ASN:OD1	1:B:279:SER:HB2	2.19	0.42
1:A:280:LYS:HD2	1:B:252:TRP:CZ2	2.55	0.42
1:C:105:LEU:C	1:C:105:LEU:HD12	2.40	0.42
1:D:257:ASN:O	1:D:258:ILE:C	2.58	0.42
1:A:11:LEU:O	1:A:15:LEU:HG	2.21	0.41
1:B:200:LYS:HA	1:B:200:LYS:HD2	1.93	0.41
1:B:277:MET:O	1:B:278:SER:C	2.59	0.41
1:A:235:GLN:HE21	1:B:297:LEU:HD11	1.85	0.41
1:A:263:THR:O	1:A:264:ARG:C	2.59	0.41
1:B:36:ARG:HD3	5:B:453:HOH:O	2.21	0.41
1:A:42:LEU:HA	1:A:42:LEU:HD12	1.96	0.41
1:B:110:LYS:HE2	5:B:475:HOH:O	2.20	0.40
1:A:244:ARG:HD3	5:A:435:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	297/300 (99%)	284 (96%)	13 (4%)	0	100	100
1	B	296/300 (99%)	286 (97%)	9 (3%)	1 (0%)	41	51
1	C	296/300 (99%)	289 (98%)	7 (2%)	0	100	100
1	D	296/300 (99%)	284 (96%)	11 (4%)	1 (0%)	41	51
All	All	1185/1200 (99%)	1143 (96%)	40 (3%)	2 (0%)	47	60

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	264	ARG
1	B	117	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	275/276 (100%)	268 (98%)	7 (2%)	47 62
1	B	274/276 (99%)	264 (96%)	10 (4%)	35 47
1	C	274/276 (99%)	265 (97%)	9 (3%)	38 51
1	D	274/276 (99%)	256 (93%)	18 (7%)	16 22
All	All	1097/1104 (99%)	1053 (96%)	44 (4%)	31 43

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	SER
1	A	49	LEU
1	A	70	VAL
1	A	200	LYS
1	A	201	LYS
1	A	211	PHE
1	A	269	GLU
1	B	10	SER
1	B	13	GLU
1	B	200	LYS
1	B	205	SER
1	B	211	PHE
1	B	221	ASP
1	B	229	THR
1	B	278	SER
1	B	286	SER
1	B	295	ASP
1	C	22	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	42	LEU
1	C	86	GLU
1	C	211	PHE
1	C	221	ASP
1	C	224	LYS
1	C	262	PHE
1	C	273	LYS
1	C	278	SER
1	D	13	GLU
1	D	21	LYS
1	D	22	GLN
1	D	26	SER
1	D	29	SER
1	D	45	SER
1	D	49	LEU
1	D	54	LYS
1	D	83	LYS
1	D	179	ASP
1	D	197	LYS
1	D	211	PHE
1	D	224	LYS
1	D	228	ASP
1	D	243	GLU
1	D	260	LYS
1	D	262	PHE
1	D	265	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	GLN
1	A	154	GLN
1	A	235	GLN
1	C	22	GLN
1	C	87	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	C	302	-	5,5,5	0.73	0	5,5,5	0.84	0
4	GOL	B	301	-	5,5,5	0.36	0	5,5,5	0.30	0
4	GOL	B	302	-	5,5,5	0.94	0	5,5,5	1.80	2 (40%)
4	GOL	C	301	-	5,5,5	0.37	0	5,5,5	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	C	302	-	-	1/4/4/4	-
4	GOL	B	301	-	-	1/4/4/4	-
4	GOL	B	302	-	-	4/4/4/4	-
4	GOL	C	301	-	-	0/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	B	302	GOL	C3-C2-C1	-2.64	101.43	111.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	302	GOL	O2-C2-C3	2.54	120.30	109.12

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	302	GOL	C1-C2-C3-O3
4	B	302	GOL	O2-C2-C3-O3
4	B	301	GOL	C1-C2-C3-O3
4	B	302	GOL	O1-C1-C2-C3
4	B	302	GOL	O1-C1-C2-O2
4	C	302	GOL	O1-C1-C2-C3

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	302	GOL	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	298/300 (99%)	-0.19	3 (1%) 82 86	14, 27, 49, 66	0
1	B	298/300 (99%)	-0.21	1 (0%) 94 96	15, 28, 50, 65	0
1	C	298/300 (99%)	-0.05	7 (2%) 60 67	15, 29, 66, 101	0
1	D	298/300 (99%)	0.00	13 (4%) 34 41	14, 29, 72, 95	0
2	E	12/12 (100%)	-0.69	0 100 100	16, 20, 24, 25	0
2	F	12/12 (100%)	-0.73	0 100 100	16, 18, 22, 25	0
2	G	12/12 (100%)	-0.75	0 100 100	17, 20, 25, 25	0
2	H	12/12 (100%)	-0.74	0 100 100	17, 19, 24, 26	0
2	I	12/12 (100%)	0.50	0 100 100	25, 40, 48, 50	2 (16%)
2	J	12/12 (100%)	0.51	0 100 100	24, 39, 47, 49	6 (50%)
2	K	12/12 (100%)	0.44	0 100 100	22, 36, 44, 44	7 (58%)
2	L	12/12 (100%)	0.56	0 100 100	23, 36, 49, 51	6 (50%)
All	All	1288/1296 (99%)	-0.11	24 (1%) 66 73	14, 28, 54, 101	21 (1%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	259	ILE	3.5
1	D	263	THR	3.5
1	B	17	ASN	3.2
1	D	270	ALA	3.0
1	D	262	PHE	2.9
1	C	265	GLU	2.7
1	C	261	ASN	2.7
1	D	17	ASN	2.7
1	D	265	GLU	2.6
1	D	261	ASN	2.6
1	D	273	LYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	278	SER	2.4
1	C	269	GLU	2.4
1	C	263	THR	2.3
1	C	2	LYS	2.3
1	D	267	ALA	2.2
1	A	142	THR	2.2
1	A	273	LYS	2.2
1	D	2	LYS	2.1
1	D	257	ASN	2.1
1	A	33	LYS	2.1
1	C	273	LYS	2.0
1	C	240	PHE	2.0
1	D	274	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	B	302	6/6	0.93	0.16	31,33,35,43	0
4	GOL	C	302	6/6	0.95	0.15	33,36,38,40	0
3	MN	G	101	1/1	0.97	0.07	34,34,34,34	1
4	GOL	C	301	6/6	0.97	0.13	34,40,45,46	0
3	MN	D	302	1/1	0.97	0.08	31,31,31,31	1
3	MN	F	101	1/1	0.97	0.08	29,29,29,29	1
4	GOL	B	301	6/6	0.98	0.13	28,32,36,39	0
3	MN	D	301	1/1	0.99	0.11	23,23,23,23	0
3	MN	B	303	1/1	0.99	0.09	23,23,23,23	0
3	MN	C	303	1/1	0.99	0.10	25,25,25,25	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MN	A	301	1/1	1.00	0.12	24,24,24,24	0
3	MN	E	101	1/1	1.00	0.12	35,35,35,35	1

6.5 Other polymers [i](#)

There are no such residues in this entry.