

Full wwPDB X-ray Structure Validation Report (i)

Oct 9, 2023 - 08:06 AM EDT

PDB ID	:	6X0T
Title	:	Structure of human plasma factor XIIa in complex with (2S)-1-(N,3-dicycloh
		exyl-D-alanyl)-4-[(4R,5S)-4-methyl-5-phenyl-4,5-dihydro-1,3-oxazol-2-yl]-N-[(
		thiophen-2-yl)methyl]piperazine-2-carboxamide (compound 8h)
Authors	:	Orth, P.
Deposited on	:	2020-05-17
Resolution	:	1.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R _{free}	130704	2907 (1.40-1.36)
Clashscore	141614	3037 (1.40-1.36)
Ramachandran outliers	138981	2970 (1.40-1.36)
Sidechain outliers	138945	2969 (1.40-1.36)
RSRZ outliers	127900	2846 (1.40-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	259	14%	۹%	7%
		200	15%	970	770
1	В	259	83%	8%	8%
1	С	259	86%	7%	7%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 5816 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Δ	941	Total	С	Ν	Ο	\mathbf{S}	0	0	0
1	A	241	1796	1132	319	330	15	0	0	0
1	р	028	Total	С	Ν	0	S	0	0	0
1	D	230	1784	1125	317	327	15	0	0	0
1	С	240	Total	С	Ν	0	S	0	0	0
		240	1796	1131	316	334	15	0	U	0

• Molecule 1 is a protein called Coagulation factor XII.

• Molecule 2 is (2S)-1-(N,3-dicyclohexyl-D-alanyl)-4-[(4R,5S)-4-methyl-5-phenyl-4,5-dihydro-1,3-oxazol-2-yl]-N-[(thiophen-2-yl)methyl]piperazine-2-carboxamide (three-letter code: UJ7) (formula: C₃₅H₄₉N₅O₃S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	Δ	1	Total	С	Ν	0	\mathbf{S}	0	0	
2	11	1	44	35	5	3	1	0	0	
9	В	1	Total	С	Ν	Ο	\mathbf{S}	0	0	
	D		44	35	5	3	1	0	0	



Continued from previous page...

Mol	Chain	Residues		Ato	\mathbf{ms}			ZeroOcc	AltConf
2	С	1	Total 44	C 35	N 5	0 3	S 1	0	0

• Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	А	1	Total 5	0 4	S 1	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	93	Total O 93 93	0	0
4	В	105	Total O 105 105	0	0
4	С	105	Total O 105 105	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Coagulation factor XII



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	77.12Å 133.41Å 88.86Å	Depositor
a, b, c, α , β , γ	90.00° 104.40° 90.00°	Depositor
$\mathbf{Posolution} \left(\overset{\circ}{\mathbf{A}} \right)$	47.26 - 1.39	Depositor
Resolution (A)	47.26 - 1.39	EDS
% Data completeness	63.5 (47.26-1.39)	Depositor
(in resolution range)	63.5(47.26-1.39)	EDS
R_{merge}	0.09	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.46 (at 1.39 \text{\AA})$	Xtriage
Refinement program	BUSTER 2.11.7 (6-FEB-2020)	Depositor
P. P.	0.289 , 0.311	Depositor
n, n_{free}	0.290 , 0.316	DCC
R_{free} test set	2229 reflections (2.01%)	wwPDB-VP
Wilson B-factor $(Å^2)$	13.7	Xtriage
Anisotropy	0.344	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.37, 37.7	EDS
L-test for twinning ²	$ < L >=0.47, < L^2>=0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	5816	wwPDB-VP
Average B, all atoms $(Å^2)$	10.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.84% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, $\rm UJ7$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.42	0/1841	0.62	0/2510	
1	В	0.43	0/1828	0.62	0/2491	
1	С	0.43	0/1841	0.63	0/2510	
All	All	0.42	0/5510	0.62	0/7511	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1796	0	1696	10	0
1	В	1784	0	1692	10	0
1	С	1796	0	1695	9	0
2	А	44	0	0	0	0
2	В	44	0	0	0	0
2	С	44	0	0	0	0
3	А	5	0	0	0	0
4	А	93	0	0	0	0
4	В	105	0	0	0	0
4	С	105	0	0	0	0
All	All	5816	0	5083	26	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:431:ARG:HH21	1:A:433:ASN:HD21	1.29	0.81
1:B:501:GLN:HE21	1:B:523:GLN:NE2	1.84	0.74
1:C:431:ARG:HH21	1:C:433:ASN:HD21	1.49	0.59
1:B:431:ARG:HH21	1:B:433:ASN:HD21	1.51	0.58
1:B:488:PRO:HD2	1:B:606:LEU:HD21	1.96	0.47
1:A:488:PRO:HD2	1:A:606:LEU:HD21	1.98	0.45
1:A:502:VAL:HG23	1:A:584:ILE:HD11	1.98	0.45
1:C:392:TRP:HB3	1:C:423:LEU:HD23	1.99	0.45
1:C:431:ARG:HH21	1:C:433:ASN:ND2	2.13	0.45
1:A:510:GLU:OE1	1:A:589:GLY:HA2	2.17	0.44
1:C:502:VAL:HG23	1:C:584:ILE:HD11	1.99	0.44
1:A:582:GLY:HA2	1:A:600:THR:O	2.18	0.43
1:B:392:TRP:HB3	1:B:423:LEU:HD23	2.00	0.43
1:B:432:ARG:HG3	1:B:505:TRP:CD1	2.53	0.43
1:B:582:GLY:HA2	1:B:600:THR:O	2.18	0.43
1:B:562:ASP:HB3	1:B:584:ILE:HD13	2.01	0.42
1:C:562:ASP:HB3	1:C:584:ILE:HD13	2.01	0.42
1:A:392:TRP:HB3	1:A:423:LEU:HD23	2.01	0.42
1:A:562:ASP:HB3	1:A:584:ILE:HD13	2.01	0.42
1:C:532:CYS:SG	1:C:538:HIS:HD2	2.43	0.42
1:A:417:ARG:CZ	1:C:452:ALA:HB2	2.50	0.41
1:A:532:CYS:SG	1:A:538:HIS:HD2	2.44	0.41
1:A:452:ALA:HB2	1:B:417:ARG:CZ	2.52	0.40
1:B:452:ALA:HB2	1:C:417:ARG:CZ	2.51	0.40
1:B:532:CYS:SG	1:B:538:HIS:HD2	2.43	0.40
1:C:386:TYR:CG	1:C:485:VAL:HB	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	235/259~(91%)	227~(97%)	7 (3%)	1 (0%)	34 12
1	В	230/259~(89%)	222 (96%)	7 (3%)	1 (0%)	34 12
1	С	234/259~(90%)	226 (97%)	7 (3%)	1 (0%)	34 12
All	All	699/777~(90%)	675~(97%)	21 (3%)	3~(0%)	34 12

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	537	VAL
1	В	537	VAL
1	С	537	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	А	184/207~(89%)	178~(97%)	6 (3%)	38 8		
1	В	185/207~(89%)	182 (98%)	3(2%)	62 33		
1	С	186/207~(90%)	183 (98%)	3(2%)	62 33		
All	All	555/621~(89%)	543 (98%)	12 (2%)	52 19		

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	364	ARG
1	А	435	SER
1	А	436	CYS
1	А	499	LEU
1	А	501	GLN
1	А	523	GLN
1	В	364	ARG



Continued from previous page...

Mol	Chain	Res	Type
1	В	436	CYS
1	В	499	LEU
1	С	364	ARG
1	С	436	CYS
1	С	568	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	523	GLN
1	А	594	ASN
1	В	523	GLN
1	В	594	ASN
1	С	572	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mal	Mal Trung Chain Bas		Dec	Tinle	Bo	Bond lengths			Bond angles		
WIOI	туре	Unain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2	
3	SO4	А	702	-	4,4,4	0.34	0	6,6,6	0.59	0	
2	UJ7	С	701	-	48,49,49	0.92	4 (8%)	50,67,67	1.51	5 (10%)	
2	UJ7	В	701	-	48,49,49	1.01	3 (6%)	50,67,67	1.56	5 (10%)	
2	UJ7	А	701	-	48,49,49	1.02	5 (10%)	50,67,67	1.60	2 (4%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UJ7	С	701	-	-	1/31/74/74	0/6/6/6
2	UJ7	В	701	-	-	1/31/74/74	0/6/6/6
2	UJ7	А	701	-	-	1/31/74/74	0/6/6/6

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	В	701	UJ7	C10-C11	-4.25	1.46	1.51
2	А	701	UJ7	C10-C11	-3.47	1.47	1.51
2	С	701	UJ7	C10-C11	-2.86	1.48	1.51
2	А	701	UJ7	C36-N37	-2.43	1.44	1.48
2	С	701	UJ7	C36-N37	-2.34	1.44	1.48
2	А	701	UJ7	C16-N6	2.21	1.39	1.34
2	В	701	UJ7	O34-C35	-2.20	1.43	1.46
2	А	701	UJ7	C36-C35	-2.17	1.52	1.55
2	А	701	UJ7	C7-N8	2.11	1.38	1.33
2	В	701	UJ7	C36-N37	-2.08	1.45	1.48
2	С	701	UJ7	C7-N8	2.06	1.38	1.33
2	С	701	UJ7	O34-C35	-2.04	1.43	1.46

All (12) bond length outliers are listed below:

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
2	А	701	UJ7	O34-C33-N37	-8.37	113.36	117.80
2	В	701	UJ7	O34-C33-N37	-7.83	113.65	117.80
2	С	701	UJ7	O34-C33-N37	-7.37	113.89	117.80
2	С	701	UJ7	C3-N1-C2	4.25	121.57	113.06
2	В	701	UJ7	C3-N1-C2	3.97	121.01	113.06
2	А	701	UJ7	C3-N1-C2	3.69	120.45	113.06
2	В	701	UJ7	C7-C4-N6	-2.74	104.58	111.13



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	В	701	UJ7	C4-C7-N8	-2.32	111.00	116.11
2	С	701	UJ7	C5-N6-C4	2.27	120.58	115.80
2	С	701	UJ7	C7-C4-N6	-2.23	105.79	111.13
2	В	701	UJ7	C5-N6-C4	2.19	120.41	115.80
2	С	701	UJ7	C10-N8-C7	-2.11	119.30	122.34

Continued from previous page...

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	С	701	UJ7	C16-C17-N20-C21
2	А	701	UJ7	C16-C17-N20-C21
2	В	701	UJ7	C16-C17-N20-C21

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.













5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2		$OWAB(Å^2)$	Q<0.9
1	А	241/259~(93%)	1.26	35~(14%) 2	2	3, 9, 22, 29	0
1	В	238/259~(91%)	1.28	38 (15%) 1	1	3, 9, 20, 28	0
1	С	240/259~(92%)	1.21	31 (12%) 3	3	3, 9, 21, 24	0
All	All	719/777~(92%)	1.25	104 (14%) 2	2	3, 9, 21, 29	0

All (104) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ	
1	А	357	LEU	9.0	
1	В	614	VAL	7.3	
1	А	471	ALA	6.5	
1	С	499	LEU	6.1	
1	В	498	THR	5.8	
1	В	499	LEU	5.2	
1	А	614	VAL	4.9	
1	В	532	CYS	4.8	
1	А	472	ASP	4.3	
1	С	529	LEU	4.3	
1	С	532	CYS	4.3	
1	А	394	HIS	4.2	
1	А	499	LEU	4.0	
1	С	489	SER	3.9	
1	В	573	ALA	3.9	
1	В	515	TYR	3.9	
1	А	358	SER	3.8	
1	А	473	GLY	3.8	
1	С	614	VAL	3.6	
1	С	515	TYR	3.6	
1	В	526	PHE	3.6	
1	С	394	HIS	3.6	
1	С	606	LEU	3.6	



6X0T

Mol	Chain	Res	Type	RSRZ
1	А	477	LEU	3.5
1	А	511	GLY	3.5
1	А	532	CYS	3.4
1	В	397	CYS	3.4
1	С	397	CYS	3.4
1	В	511	GLY	3.3
1	А	529	LEU	3.3
1	А	396	PHE	3.2
1	В	489	SER	3.2
1	А	470	ASP	3.2
1	В	358	SER	3.2
1	А	435	SER	3.2
1	А	526	PHE	3.1
1	В	470	ASP	3.1
1	С	435	SER	3.1
1	А	481	TYR	3.1
1	В	531	ARG	3.0
1	А	515	TYR	3.0
1	А	397	CYS	3.0
1	А	536	ASP	3.0
1	В	394	HIS	2.9
1	А	609	ILE	2.9
1	В	529	LEU	2.9
1	С	477	LEU	2.9
1	А	489	SER	2.9
1	В	609	ILE	2.9
1	С	358	SER	2.9
1	В	359	CYS	2.8
1	А	570	GLU	2.8
1	В	536	ASP	2.8
1	С	396	PHE	2.8
1	С	577	ARG	2.8
1	А	436	CYS	2.7
1	С	573	ALA	2.7
1	В	575	GLU	2.6
1	С	471	ALA	2.6
1	В	519	LEU	2.6
1	А	590	CYS	2.6
1	А	552	LEU	2.6
1	В	551	PHE	2.6
1	А	500	CYS	2.6
1	В	610	ARG	2.6

Continued from previous page...



Mol	Chain	Res	Type	RSRZ	
1	C	572	GLN	2.6	
1	В	509	PHE	2.6	
1	С	475	CYS	2.5	
1	В	436	CYS	2.5	
1	В	465	LEU	2.5	
1	С	509	PHE	2.5	
1	С	590	CYS	2.4	
1	В	606	LEU	2.4	
1	А	531	ARG	2.4	
1	С	406	TRP	2.4	
1	А	513	GLU	2.4	
1	А	509	PHE	2.4	
1	С	608	TRP	2.3	
1	В	439	CYS	2.3	
1	В	500	CYS	2.3	
1	А	519	LEU	2.3	
1	В	481	TYR	2.3	
1	С	575	GLU	2.3	
1	В	414	LEU	2.3	
1	С	536	ASP	2.3	
1	В	572	GLN	2.2	
1	В	613	THR	2.2	
1	С	481	TYR	2.2	
1	А	551	PHE	2.2	
1	В	558	ALA	2.2	
1	А	527	LEU	2.2	
1	С	552	LEU	2.2	
1	С	551	PHE	2.1	
1	A	424	THR	2.1	
1	С	526	PHE	2.1	
1	A	512	ALA	2.1	
1	В	527	LEU	2.1	
1	В	458	TYR	2.1	
1	С	574	ALA	2.1	
1	С	470	ASP	2.0	
1	В	444	VAL	2.0	
1	С	407	VAL	2.0	
1	В	475	CYS	2.0	
1	В	408	LEU	2.0	

Continued from previous page...



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B} ext{-factors}({ m \AA}^2)$	Q < 0.9
2	UJ7	С	701	44/44	0.87	0.14	$9,\!11,\!16,\!16$	0
2	UJ7	В	701	44/44	0.89	0.13	7,12,14,14	0
2	UJ7	А	701	44/44	0.89	0.12	8,10,13,13	0
3	SO4	А	702	5/5	0.97	0.12	8,8,9,9	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.













6.5 Other polymers (i)

There are no such residues in this entry.

