

Full wwPDB X-ray Structure Validation Report (i)

Oct 10, 2023 – 04:21 AM EDT

PDB ID : 6X1Z

Title: Mre11 dimer in complex with small molecule modulator PFMJ

Authors: Arvai, A.S.; Moiani, D.; Tainer, J.A.

Deposited on : 2020-05-19

Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	Similar resolution $(\# \text{Entries, resolution range}(\text{\AA}))$
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	A	336	7% 83%	14%		
1	В	336	78%	16%		



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 5694 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Nuclease SbcCD subunit D.

	\mathbf{Mol}	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
Ī	1	Λ	329	Total	С	N	О	S	0	0	0
	1	Λ	329	2653	1704	456	487	6	0	U	
	1	B	326	Total	С	N	О	S	0	1	0
	1	D	320	2636	1693	452	485	6		1	U

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	initiating methionine	UNP Q9X1X0
A	-10	GLY	-	expression tag	UNP Q9X1X0
A	-9	SER	-	expression tag	UNP Q9X1X0
A	-8	ASP	_	expression tag	UNP Q9X1X0
A	-7	LYS	-	expression tag	UNP Q9X1X0
A	-6	ILE	-	expression tag	UNP Q9X1X0
A	-5	HIS	-	expression tag	UNP Q9X1X0
A	-4	HIS	-	expression tag	UNP Q9X1X0
A	-3	HIS	-	expression tag	UNP Q9X1X0
A	-2	HIS	-	expression tag	UNP Q9X1X0
A	-1	HIS	-	expression tag	UNP Q9X1X0
A	0	HIS	-	expression tag	UNP Q9X1X0
A	1	VAL	-	expression tag	UNP Q9X1X0
В	-11	MET	-	initiating methionine	UNP Q9X1X0
В	-10	GLY	-	expression tag	UNP Q9X1X0
В	-9	SER	-	expression tag	UNP Q9X1X0
В	-8	ASP	-	expression tag	UNP Q9X1X0
В	-7	LYS	-	expression tag	UNP Q9X1X0
В	-6	ILE	-	expression tag	UNP Q9X1X0
В	-5	HIS	-	expression tag	UNP Q9X1X0
В	-4	HIS	-	expression tag	UNP Q9X1X0
В	-3	HIS	-	expression tag	UNP Q9X1X0
В	-2	HIS	-	expression tag	UNP Q9X1X0
В	-1	HIS	-	expression tag	UNP Q9X1X0
В	0	HIS	-	expression tag	UNP Q9X1X0



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	1	VAL	-	expression tag	UNP Q9X1X0

• Molecule 2 is (5Z)-5-[(3,4-dimethoxyphenyl)methylidene]-2-sulfanylidene-1,3-thiazolidin-4-one (three-letter code: UL1) (formula: $C_{12}H_{11}NO_3S_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
9	Λ	1	Total	С	N	О	S	0	0
	A	1	18	12	1	3	2	0	0
9	D	1	Total	С	N	О	S	0	0
	Б	1	18	12	1	3	2	0	0

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	В	1	Total Mg 1 1	0	0

• Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: $C_6H_{13}NO_4S$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf			
4	Λ	1	Total	С	N	О	S	0	0	
4	A	1	12	6	1	4	1	0		
1	D	1	Total	С	N	О	S	0	0	
4	Б	1	12	6	1	4	1		U	

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	178	Total O 178 178	0	0
5	В	165	Total O 165 165	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	47.26Å 109.44Å 75.65Å	Donositon
a, b, c, α , β , γ	90.00° 99.97° 90.00°	Depositor
Resolution (Å)	37.25 - 1.90	Depositor
Resolution (A)	37.25 - 1.90	EDS
% Data completeness	85.9 (37.25-1.90)	Depositor
(in resolution range)	85.9 (37.25-1.90)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.97 (at 1.91Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
D D.	0.200 , 0.239	Depositor
R, R_{free}	0.200 , 0.238	DCC
R_{free} test set	2000 reflections (3.53%)	wwPDB-VP
Wilson B-factor (Å ²)	28.4	Xtriage
Anisotropy	0.216	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33, 48.5	EDS
L-test for twinning ²	$ < L > = 0.55, < L^2> = 0.39$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5694	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 49.75 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.1357e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: UL1, MES, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain		nd lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z >5	
1	A	0.44	0/2715	0.60	3/3672 (0.1%)	
1	В	0.41	1/2698 (0.0%)	0.58	1/3647 (0.0%)	
All	All	0.43	1/5413 (0.0%)	0.59	4/7319 (0.1%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	В	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$[Ideal(ext{Å})]$
1	В	194	GLN	CB-CG	-5.45	1.37	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	256	ARG	NE-CZ-NH1	-6.35	117.12	120.30
1	A	5	LYS	CD-CE-NZ	5.45	124.22	111.70
1	В	199	ILE	CB-CG1-CD1	-5.24	99.22	113.90
1	A	5	LYS	CB-CG-CD	-5.07	98.41	111.60

There are no chirality outliers.

All (2) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	A	193	GLU	Peptide
1	В	142	GLU	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	2653	0	2651	28	0
1	В	2636	0	2654	57	0
2	A	18	0	0	0	0
2	В	18	0	0	0	0
3	A	1	0	0	0	0
3	В	1	0	0	0	0
4	A	12	0	12	1	0
4	В	12	0	12	1	0
5	A	178	0	0	3	1
5	В	165	0	0	5	1
All	All	5694	0	5329	85	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)
1:A:5:LYS:HD2	1:A:256:ARG:HD3	1.46	0.96
1:B:194:GLN:HE22	1:B:199:ILE:HD11	1.29	0.94
1:B:194:GLN:NE2	1:B:199:ILE:HD11	1.94	0.83
1:B:143:SER:HA	1:B:146:LEU:HD22	1.67	0.76
1:A:147:ARG:HG3	1:A:153:PHE:HB2	1.67	0.76
1:B:144:GLU:O	1:B:147:ARG:HB2	1.86	0.74
1:B:2:ILE:HG23	1:B:4:LEU:HB2	1.70	0.73
1:B:142:GLU:HA	1:B:144:GLU:HG3	1.71	0.72
1:B:194:GLN:HE22	1:B:199:ILE:CD1	2.04	0.71
1:B:142:GLU:HA	1:B:142:GLU:OE1	1.89	0.70
1:B:187:ALA:O	1:B:221:ARG:NH2	2.24	0.69
1:A:65:ASN:O	5:A:701:HOH:O	2.09	0.69



 $Continued\ from\ previous\ page...$

Continuea from previo		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:147:ARG:NH2	1:B:150:GLU:OE1	2.26	0.69
1:A:142:GLU:HG3	1:A:145:ALA:HB3	1.74	0.68
1:B:65:ASN:O	5:B:701:HOH:O	2.11	0.68
1:A:192:ILE:C	1:A:193:GLU:HG2	2.15	0.67
1:B:63:ARG:HB3	4:B:603:MES:H62	1.77	0.66
1:A:316:GLU:OE1	5:A:702:HOH:O	2.13	0.66
1:B:1:VAL:H	1:B:2:ILE:HA	1.62	0.65
1:B:279:LYS:NZ	1:B:306:ASP:O	2.31	0.64
1:B:4:LEU:HD11	1:B:173:ASP:OD2	2.00	0.61
1:A:10:LEU:HD22	1:A:45:ALA:HB2	1.84	0.59
1:B:100:LYS:NZ	1:B:141:ASP:OD2	2.34	0.59
1:A:63:ARG:HB3	4:A:603:MES:H62	1.85	0.58
1:B:149:ASN:N	1:B:149:ASN:OD1	2.36	0.58
1:B:3:ASN:OD1	1:B:128:ARG:NH1	2.36	0.58
1:B:311:PRO:O	1:B:313:LEU:N	2.37	0.58
1:B:142:GLU:OE1	1:B:145:ALA:N	2.37	0.57
1:B:131:LYS:HB2	1:B:173:ASP:OD2	2.05	0.57
1:B:278:LYS:O	1:B:305:GLU:HB2	2.05	0.56
1:A:273:LYS:HG2	1:A:291:PHE:CZ	2.41	0.56
1:A:140:PRO:HG2	1:A:144:GLU:HB3	1.90	0.54
1:B:26:ARG:NH2	1:B:317:ILE:O	2.39	0.54
1:A:100:LYS:HA	1:A:117:MET:HE1	1.90	0.54
1:B:141:ASP:C	1:B:143:SER:H	2.12	0.52
1:B:2:ILE:CG2	1:B:4:LEU:HB2	2.38	0.52
1:B:104:ASN:HB2	5:B:830:HOH:O	2.08	0.51
1:B:140:PRO:O	1:B:144:GLU:HG2	2.11	0.51
1:A:26:ARG:NH2	1:A:314:MET:O	2.44	0.50
1:A:103:GLY:HA3	1:A:117:MET:HE2	1.94	0.50
1:A:278:LYS:HG3	1:A:279:LYS:N	2.27	0.50
1:B:5:LYS:O	1:B:255:LYS:HA	2.12	0.50
1:B:273:LYS:HG2	1:B:291:PHE:CZ	2.47	0.49
1:B:142:GLU:CD	1:B:145:ALA:HB2	2.32	0.49
1:B:190:ALA:O	5:B:702:HOH:O	2.19	0.49
1:A:166:GLU:OE1	5:A:703:HOH:O	2.20	0.49
1:A:13:SER:OG	1:A:14:ASP:OD2	2.20	0.48
1:B:172:GLU:HG2	1:B:173:ASP:OD1	2.13	0.48
1:B:142:GLU:OE1	1:B:144:GLU:HG3	2.12	0.48
1:B:140:PRO:O	1:B:142:GLU:N	2.46	0.48
1:B:186:LEU:HD23	1:B:186:LEU:HA	1.72	0.48
1:A:189:TYR:CE2	1:A:191:GLY:HA2	2.49	0.47
1:B:194:GLN:NE2	1:B:199:ILE:CD1	2.69	0.47



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${ m distance}({ m \AA})$	overlap (Å)
1:B:253:GLU:OE1	1:B:261[B]:ARG:NH2	2.46	0.47
1:B:261[A]:ARG:NH2	5:B:710:HOH:O	2.45	0.47
1:A:207:SER:HA	1:A:227:PRO:HD3	1.97	0.47
1:B:125:GLU:OE1	1:B:131:LYS:NZ	2.47	0.47
1:B:4:LEU:HA	1:B:4:LEU:HD12	1.48	0.46
1:B:140:PRO:HB2	1:B:144:GLU:HG2	1.98	0.45
1:B:1:VAL:N	1:B:2:ILE:HA	2.27	0.45
1:A:125:GLU:OE2	1:A:131:LYS:NZ	2.48	0.45
1:B:2:ILE:HD11	1:B:129:GLY:O	2.17	0.45
1:A:186:LEU:HD23	1:A:186:LEU:HA	1.75	0.44
1:B:186:LEU:HD22	1:B:221:ARG:NH1	2.33	0.44
1:B:243:GLU:HG3	1:B:272:LEU:HB2	2.00	0.44
1:B:142:GLU:O	1:B:146:LEU:HD13	2.18	0.44
1:A:168:ALA:O	1:A:171:LYS:HB2	2.18	0.43
1:A:320:LEU:HD21	1:A:323:ILE:HD11	1.99	0.43
1:A:133:ARG:NH1	1:A:167:GLU:OE2	2.51	0.43
1:A:103:GLY:HA3	1:A:117:MET:CE	2.48	0.43
1:B:207:SER:HA	1:B:227:PRO:HD3	2.01	0.43
1:A:5:LYS:CD	1:A:256:ARG:HD3	2.32	0.43
1:B:13:SER:OG	1:B:14:ASP:OD2	2.35	0.43
1:A:243:GLU:HG3	1:A:272:LEU:HB2	2.01	0.42
1:B:141:ASP:C	1:B:143:SER:N	2.73	0.42
1:A:182:THR:O	1:A:199:ILE:HA	2.19	0.42
1:B:11:HIS:CE1	1:B:213:ALA:HB1	2.55	0.42
1:B:100:LYS:HE3	1:B:100:LYS:HB2	1.75	0.41
1:B:199:ILE:HG21	1:B:199:ILE:HD13	1.77	0.41
1:B:97:LYS:O	5:B:703:HOH:O	2.22	0.41
1:B:1:VAL:HB	1:B:2:ILE:C	2.40	0.41
1:B:219:SER:HB3	1:B:245:ASP:CG	2.42	0.40
1:B:189:TYR:CE2	1:B:191:GLY:HA2	2.56	0.40
1:A:226:GLN:HA	1:A:227:PRO:HA	1.85	0.40
1:B:131:LYS:O	1:B:173:ASP:HB2	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-1 Atom-2		$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
5:A:857:HOH:O	5:B:731:HOH:O[2_658]	2.14	0.06



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers	Percentiles	\mathbf{s}
1	A	327/336~(97%)	309 (94%)	17 (5%)	1 (0%)	41 31	
1	В	325/336~(97%)	311 (96%)	12 (4%)	2 (1%)	25 15	
All	All	$652/672 \ (97\%)$	620 (95%)	29 (4%)	3 (0%)	29 18	

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	216	HIS
1	В	216	HIS
1	В	142	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	288/298 (97%)	281 (98%)	7 (2%)	49	43
1	В	288/298 (97%)	275 (96%)	13 (4%)	27	18
All	All	576/596 (97%)	556 (96%)	20 (4%)	36	27

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	\mathbf{Type}
1	A	-3	HIS
1	A	139	TYR
1	A	141	ASP



Continued from previous page...

Mol	Chain	Res	Type
1	A	146	LEU
1	A	147	ARG
1	A	148	LYS
1	A	193	GLU
1	В	-1	HIS
1	В	3	ASN
1	В	139	TYR
1	В	141	ASP
1	В	142	GLU
1	В	143	SER
1	В	144	GLU
1	В	146	LEU
1	В	147	ARG
1	В	148	LYS
1	В	149	ASN
1	В	152	ASP
1	В	282	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	В	194	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type Ch	Chain	Chain Dag	s Link	Bond lengths			Bond angles			
MIOI	Type	Chain	Res	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	UL1	A	601	-	19,19,19	2.08	4 (21%)	26,26,26	3.11	9 (34%)
4	MES	В	603	-	12,12,12	2.22	1 (8%)	14,16,16	1.85	3 (21%)
4	MES	A	603	-	12,12,12	2.28	1 (8%)	14,16,16	1.83	3 (21%)
2	UL1	В	601	-	19,19,19	2.08	4 (21%)	26,26,26	2.40	8 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UL1	A	601	_	-	0/8/20/20	0/2/2/2
4	MES	В	603	-	-	2/6/14/14	0/1/1/1
4	MES	A	603	-	-	2/6/14/14	0/1/1/1
2	UL1	В	601	-	-	1/8/20/20	0/2/2/2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	Ideal(A)
4	A	603	MES	C8-S	-7.66	1.66	1.77
4	В	603	MES	C8-S	-7.39	1.67	1.77
2	A	601	UL1	CAE-CAR	5.26	1.40	1.34
2	В	601	UL1	CAE-CAR	4.85	1.40	1.34
2	A	601	UL1	CAQ-SAL	-4.84	1.68	1.74
2	В	601	UL1	CAQ-SAL	-4.51	1.68	1.74
2	В	601	UL1	CAP-CAR	-4.23	1.41	1.48
2	A	601	UL1	CAP-CAR	-3.86	1.42	1.48
2	В	601	UL1	CAM-CAE	-2.90	1.41	1.46
2	A	601	UL1	CAM-CAE	-2.54	1.41	1.46

All (23) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	${f Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}(^{o})$
2	A	601	UL1	CAR-CAP-NAI	8.11	116.98	110.22
2	В	601	UL1	CAR-CAP-NAI	6.50	115.64	110.22
2	A	601	UL1	CAP-CAR-SAL	-6.40	105.06	109.84
2	A	601	UL1	CAQ-NAI-CAP	-6.39	110.81	117.77
2	A	601	UL1	CAE-CAR-CAP	6.29	125.44	120.47
2	В	601	UL1	CAQ-NAI-CAP	-4.85	112.49	117.77
2	В	601	UL1	CAP-CAR-SAL	-4.67	106.35	109.84
4	A	603	MES	C5-N4-C3	4.21	118.31	108.83
4	В	603	MES	C5-N4-C3	4.15	118.16	108.83
2	В	601	UL1	CAE-CAR-CAP	4.09	123.70	120.47
2	A	601	UL1	CAQ-SAL-CAR	3.85	94.42	92.42
4	В	603	MES	O1S-S-C8	3.27	110.85	106.92
2	A	601	UL1	OAC-CAP-CAR	-3.23	121.93	126.01
4	A	603	MES	O3S-S-C8	3.21	110.96	105.77
2	В	601	UL1	OAC-CAP-CAR	-3.17	122.00	126.01
2	В	601	UL1	OAJ-CAN-CAO	2.83	119.35	115.41
4	A	603	MES	C7-N4-C5	2.76	118.29	111.23
2	A	601	UL1	SAL-CAQ-SAD	-2.74	119.44	124.39
4	В	603	MES	C7-N4-C5	2.72	118.18	111.23
2	A	601	UL1	OAJ-CAN-CAO	2.44	118.81	115.41
2	В	601	UL1	CAQ-SAL-CAR	2.39	93.66	92.42
2	В	601	UL1	CAM-CAE-CAR	-2.25	127.86	130.94
2	A	601	UL1	CAB-OAK-CAO	-2.12	114.32	117.53

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	603	MES	C8-C7-N4-C5
4	В	603	MES	C8-C7-N4-C5
4	A	603	MES	C8-C7-N4-C3
4	В	603	MES	C8-C7-N4-C3
2	В	601	UL1	CAO-CAN-OAJ-CAA

There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	603	MES	1	0
4	A	603	MES	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,



bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	329/336~(97%)	0.19	22 (6%) 17 20	20, 34, 100, 185	0
1	В	326/336~(97%)	0.51	39 (11%) 4 4	24, 41, 139, 207	0
All	All	655/672~(97%)	0.35	61 (9%) 8 10	20, 38, 126, 207	0

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	2	ILE	20.3
1	В	1	VAL	12.9
1	В	194	GLN	9.6
1	В	148	LYS	8.5
1	В	-1	HIS	8.4
1	В	3	ASN	7.5
1	A	193	GLU	7.0
1	A	148	LYS	6.7
1	A	146	LEU	6.6
1	A	194	GLN	6.4
1	В	145	ALA	6.4
1	В	191	GLY	6.4
1	A	-4	HIS	6.2
1	В	0	HIS	5.9
1	В	4	LEU	5.8
1	A	145	ALA	5.6
1	В	143	SER	5.5
1	A	143	SER	5.5
1	В	192	ILE	5.4
1	В	309	ILE	5.3
1	В	189	TYR	4.7
1	В	193	GLU	4.6
1	A	142	GLU	4.5
1	В	313	LEU	3.9



Continued from previous page...

Mol	nuea fron Chain	Res	Type	RSRZ
1	В	149	ASN	3.8
1	В	282	THR	3.6
1	A	278	LYS	3.6
1	В	278	LYS	3.5
1	В	146	LEU	3.5
1	A	147	ARG	3.5
1	В	144	GLU	3.4
1	В	276	TYR	3.4
1	В	314	MET	3.3
1	A	144	GLU	3.3
1	В	312	ASP	3.2
1	В	305	GLU	3.2
1	В	150	GLU	3.1
1	A	283	SER	3.0
1	В	147	ARG	3.0
1	A	195	GLY	3.0
1	В	195	GLY	3.0
1	В	280	ILE	2.9
1	A	257	GLY	2.9
1	В	257	GLY	2.9
1	В	285	LEU	2.8
1	В	226	GLN	2.7
1	В	196	ARG	2.7
1	В	142	GLU	2.6
1	В	190	ALA	2.5
1	В	283	SER	2.5
1	В	277	TYR	2.5
1	A	256	ARG	2.4
1	A	3	ASN	2.4
1	A	191	GLY	2.4
1	В	281	ASP	2.4
1	A	276	TYR	2.3
1	В	256	ARG	2.2
1	A	190	ALA	2.1
1	A	192	ILE	2.1
1	A	279	LYS	2.0
1	A	282	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
4	MES	A	603	12/12	0.89	0.15	74,75,77,79	0
4	MES	В	603	12/12	0.90	0.13	70,71,75,77	0
2	UL1	A	601	18/18	0.93	0.08	28,32,38,40	0
2	UL1	В	601	18/18	0.94	0.09	33,39,42,42	0
3	MG	A	602	1/1	0.95	0.07	49,49,49,49	0
3	MG	В	602	1/1	0.98	0.12	50,50,50,50	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









6.5 Other polymers (i)

There are no such residues in this entry.

