



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 30, 2023 – 05:23 PM JST

PDB ID : 4X42  
Title : Crystal structure of DEN4 ED3 mutant with epitope two residues substituted from DEN3 ED3  
Authors : Kulkarni, M.R.; Islam, M.M.; Numoto, N.; Elahi, M.M.; Ito, N.; Kuroda, Y.  
Deposited on : 2014-12-02  
Resolution : 2.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4107 (2.80-2.76)
Clashscore	141614	4575 (2.80-2.76)
Ramachandran outliers	138981	4487 (2.80-2.76)
Sidechain outliers	138945	4489 (2.80-2.76)
RSRZ outliers	127900	4027 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	107	 88% 5% 7%
1	B	107	 93% 7%
1	C	107	 83% 7% 9%
1	D	107	 91% 5% 5%
1	E	107	 86% 10%
1	F	107	 90% 7%

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4575 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope protein E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	99	Total 756	C 478	N 127	O 147	S 4	0	0	0
1	B	100	Total 764	C 483	N 128	O 148	S 5	0	0	0
1	C	97	Total 746	C 473	N 125	O 144	S 4	0	0	0
1	D	102	Total 774	C 488	N 130	O 151	S 5	0	0	0
1	E	96	Total 740	C 470	N 124	O 142	S 4	0	0	0
1	F	100	Total 764	C 483	N 128	O 148	S 5	0	0	0

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	573	GLY	-	expression tag	UNP P09866
A	574	SER	-	expression tag	UNP P09866
A	663	ASP	ASN	engineered mutation	UNP P09866
A	664	LYS	SER	engineered mutation	UNP P09866
A	667	LYS	THR	engineered mutation	UNP P09866
A	669	ASN	HIS	engineered mutation	UNP P09866
B	573	GLY	-	expression tag	UNP P09866
B	574	SER	-	expression tag	UNP P09866
B	663	ASP	ASN	engineered mutation	UNP P09866
B	664	LYS	SER	engineered mutation	UNP P09866
B	667	LYS	THR	engineered mutation	UNP P09866
B	669	ASN	HIS	engineered mutation	UNP P09866
C	573	GLY	-	expression tag	UNP P09866
C	574	SER	-	expression tag	UNP P09866
C	663	ASP	ASN	engineered mutation	UNP P09866
C	664	LYS	SER	engineered mutation	UNP P09866
C	667	LYS	THR	engineered mutation	UNP P09866

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	669	ASN	HIS	engineered mutation	UNP P09866
D	573	GLY	-	expression tag	UNP P09866
D	574	SER	-	expression tag	UNP P09866
D	663	ASP	ASN	engineered mutation	UNP P09866
D	664	LYS	SER	engineered mutation	UNP P09866
D	667	LYS	THR	engineered mutation	UNP P09866
D	669	ASN	HIS	engineered mutation	UNP P09866
E	573	GLY	-	expression tag	UNP P09866
E	574	SER	-	expression tag	UNP P09866
E	663	ASP	ASN	engineered mutation	UNP P09866
E	664	LYS	SER	engineered mutation	UNP P09866
E	667	LYS	THR	engineered mutation	UNP P09866
E	669	ASN	HIS	engineered mutation	UNP P09866
F	573	GLY	-	expression tag	UNP P09866
F	574	SER	-	expression tag	UNP P09866
F	663	ASP	ASN	engineered mutation	UNP P09866
F	664	LYS	SER	engineered mutation	UNP P09866
F	667	LYS	THR	engineered mutation	UNP P09866
F	669	ASN	HIS	engineered mutation	UNP P09866

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O	0	0
			1	1		
3	B	2	Total	O	0	0
			2	2		
3	C	1	Total	O	0	0
			1	1		
3	D	4	Total	O	0	0
			4	4		
3	E	1	Total	O	0	0
			1	1		
3	F	2	Total	O	0	0
			2	2		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Envelope protein E

Chain A: 




- Molecule 1: Envelope protein E

Chain B: 



- Molecule 1: Envelope protein E

Chain C: 




- Molecule 1: Envelope protein E

Chain D: 




- Molecule 1: Envelope protein E

Chain E: 



- Molecule 1: Envelope protein E



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	124.58Å 124.58Å 86.13Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.03 – 2.78 40.00 – 2.78	Depositor EDS
% Data completeness (in resolution range)	99.8 (40.03-2.78) 99.8 (40.00-2.78)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.56 (at 2.77Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.234 , 0.273 0.234 , 0.271	Depositor DCC
$R_{free}$ test set	938 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	51.1	Xtrriage
Anisotropy	0.037	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 25.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.039 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	4575	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.67% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.67	0/769	0.84	0/1038
1	B	0.66	0/777	0.82	0/1048
1	C	0.63	0/759	0.85	1/1025 (0.1%)
1	D	0.60	0/787	0.81	0/1061
1	E	0.59	0/753	0.79	0/1017
1	F	0.70	0/777	0.85	0/1048
All	All	0.64	0/4622	0.83	1/6237 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1
1	F	0	1
All	All	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	601	VAL	CB-CA-C	-5.99	100.03	111.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	633	SER	Peptide

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	F	633	SER	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	756	0	765	3	0
1	B	764	0	774	0	0
1	C	746	0	757	9	0
1	D	774	0	782	4	0
1	E	740	0	752	4	0
1	F	764	0	774	2	0
2	A	10	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
3	A	1	0	0	0	0
3	B	2	0	0	0	0
3	C	1	0	0	0	0
3	D	4	0	0	0	0
3	E	1	0	0	0	0
3	F	2	0	0	0	0
All	All	4575	0	4604	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:649:GLU:O	1:F:649:GLU:HG3	2.05	0.57
1:C:591:MET:HE3	1:C:601:VAL:CG2	2.36	0.55
1:E:649:GLU:O	1:E:649:GLU:HG3	2.07	0.54
1:C:589:LYS:HB3	1:C:602:LYS:HB2	1.91	0.52
1:A:621:VAL:HG12	1:A:654:ASP:HB2	1.91	0.52
1:F:621:VAL:HA	1:F:656:TYR:HE1	1.74	0.51
1:C:619:ARG:HB3	1:C:623:LYS:HA	1.93	0.51

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:602:LYS:HE2	1:A:645:ASN:OD1	2.12	0.50
1:C:591:MET:HE3	1:C:601:VAL:HG21	1.93	0.50
1:A:649:GLU:HG3	1:A:649:GLU:O	2.12	0.49
1:C:601:VAL:HG12	1:C:602:LYS:N	2.28	0.48
1:D:611:PRO:HG3	1:E:587:ILE:HG13	1.96	0.48
1:C:591:MET:CE	1:C:601:VAL:CG2	2.93	0.47
1:D:578:TYR:OH	1:E:590:GLU:HG3	2.16	0.46
1:C:601:VAL:CG1	1:C:602:LYS:N	2.82	0.42
1:D:633:SER:O	1:D:634:THR:C	2.58	0.41
1:C:602:LYS:HE3	1:C:645:ASN:OD1	2.20	0.41
1:D:577:SER:HB3	1:E:669:ASN:ND2	2.35	0.41
1:C:591:MET:HE1	1:C:657:ILE:HD13	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	97/107 (91%)	96 (99%)	1 (1%)	0	100	100
1	B	98/107 (92%)	97 (99%)	1 (1%)	0	100	100
1	C	95/107 (89%)	95 (100%)	0	0	100	100
1	D	100/107 (94%)	98 (98%)	2 (2%)	0	100	100
1	E	94/107 (88%)	94 (100%)	0	0	100	100
1	F	98/107 (92%)	98 (100%)	0	0	100	100
All	All	582/642 (91%)	578 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/90 (94%)	85 (100%)	0	100	100
1	B	86/90 (96%)	86 (100%)	0	100	100
1	C	84/90 (93%)	84 (100%)	0	100	100
1	D	87/90 (97%)	87 (100%)	0	100	100
1	E	83/90 (92%)	83 (100%)	0	100	100
1	F	86/90 (96%)	86 (100%)	0	100	100
All	All	511/540 (95%)	511 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	596	HIS
1	E	595	GLN
1	E	669	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	C	701	-	4,4,4	0.43	0	6,6,6	0.22	0
2	SO4	B	701	-	4,4,4	0.63	0	6,6,6	0.34	0
2	SO4	A	701	-	4,4,4	0.48	0	6,6,6	0.49	0
2	SO4	A	702	-	4,4,4	0.40	0	6,6,6	0.28	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	99/107 (92%)	-0.12	0 <a href="#">100</a> <a href="#">100</a>	24, 41, 63, 75	0
1	B	100/107 (93%)	0.12	1 (1%) <a href="#">82</a> <a href="#">80</a>	24, 46, 78, 89	0
1	C	97/107 (90%)	0.17	1 (1%) <a href="#">82</a> <a href="#">80</a>	40, 62, 87, 98	0
1	D	102/107 (95%)	0.20	1 (0%) <a href="#">82</a> <a href="#">80</a>	37, 62, 85, 100	0
1	E	96/107 (89%)	-0.04	0 <a href="#">100</a> <a href="#">100</a>	39, 56, 81, 116	0
1	F	100/107 (93%)	-0.11	0 <a href="#">100</a> <a href="#">100</a>	28, 42, 67, 76	0
All	All	594/642 (92%)	0.04	3 (0%) <a href="#">91</a> <a href="#">90</a>	24, 50, 83, 116	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	643	VAL	3.3
1	D	621	VAL	2.2
1	B	638	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	A	702	5/5	0.90	0.15	69,70,85,88	0
2	SO4	C	701	5/5	0.91	0.13	69,83,87,88	0
2	SO4	A	701	5/5	0.94	0.14	65,70,75,78	0
2	SO4	B	701	5/5	0.96	0.10	58,62,71,75	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.