



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2023 – 05:24 PM EDT

PDB ID : 6X4A  
Title : Crystal Structure of HIV-1 Reverse Transcriptase (Y181C) Variant in Complex with 5-chloro-7-(2-(2-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)ethoxy)phenoxy)-8-methyl-2-naphthonitrile (JLJ651), a Non-nucleoside Inhibitor  
Authors : Chan, A.H.; Duong, V.N.; Ippolito, J.A.; Jorgensen, W.L.; Anderson, K.S.  
Deposited on : 2020-05-22  
Resolution : 2.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	557	 13% 89% 9%
2	B	428	 13% 82% 11% 6%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7818 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reverse transcriptase/ribonuclease H.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	547	4413	2858	729	817	9	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	expression tag	UNP P03366
A	0	VAL	-	expression tag	UNP P03366
A	172	ALA	LYS	engineered mutation	UNP P03366
A	173	ALA	LYS	engineered mutation	UNP P03366
A	181	CYS	TYR	engineered mutation	UNP P03366
A	280	SER	CYS	engineered mutation	UNP P03366


- Molecule 2 is a protein called p51 RT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	401	3319	2160	550	603	6	0	0	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	280	SER	CYS	engineered mutation	UNP P03366

- Molecule 3 is 5-chloro-7-(2-(2-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)ethoxy)phenoxy)-8-methyl-2-naphthonitrile (three-letter code: 7AY) (formula: C<sub>24</sub>H<sub>18</sub>ClN<sub>3</sub>O<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Cl	N			O
3	A	1	32	24	1	3	4	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O S		
4	A	1	5	4 1	0	0
4	A	1	5	4 1	0	0


- Molecule 5 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	A	21	Total 21	O 21	0	0
5	B	23	Total 23	O 23	0	0


### 3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Reverse transcriptase/ribonuclease H



- Molecule 2: p51 RT



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	161.77Å 73.98Å 107.90Å 90.00° 100.05° 90.00°	Depositor
Resolution (Å)	38.13 – 2.54 43.14 – 2.54	Depositor EDS
% Data completeness (in resolution range)	99.4 (38.13-2.54) 99.4 (43.14-2.54)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.70 (at 2.54Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.232 , 0.269 0.234 , 0.271	Depositor DCC
$R_{free}$ test set	2000 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	57.2	Xtrriage
Anisotropy	0.282	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 49.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	7818	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	73.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.67% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 7AY, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.22	0/4529	0.39	0/6166
2	B	0.22	0/3413	0.39	0/4635
All	All	0.22	0/7942	0.39	0/10801

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4413	0	4426	29	0
2	B	3319	0	3341	31	0
3	A	32	0	0	0	0
4	A	10	0	0	0	0
5	A	21	0	0	0	0
5	B	23	0	0	1	0
All	All	7818	0	7767	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:544:GLY:HA2	2:B:286:THR:HG22	1.79	0.64
2:B:72:ARG:NH2	2:B:151:GLN:OE1	2.31	0.64
1:A:28:GLU:HG3	1:A:135:ILE:HD12	1.79	0.63
2:B:73:LYS:NZ	2:B:146:TYR:OH	2.32	0.62
1:A:90:VAL:HG11	2:B:140:PRO:HB3	1.82	0.61
2:B:169:GLU:OE1	2:B:173:LYS:NZ	2.34	0.61
1:A:317:VAL:HG23	1:A:349:LEU:HD23	1.85	0.58
1:A:88:TRP:CD1	2:B:143:ARG:HD2	2.40	0.55
2:B:275:LYS:HD3	2:B:276:VAL:N	2.22	0.55
1:A:369:THR:O	1:A:373:GLN:HG2	2.08	0.54
2:B:21:VAL:HB	2:B:59:PRO:HD3	1.89	0.54
2:B:425:LEU:HD23	2:B:428:GLN:HB3	1.89	0.53
1:A:543:GLY:HA3	2:B:284:ARG:HA	1.91	0.52
1:A:28:GLU:HG2	1:A:32:LYS:HE3	1.93	0.51
2:B:275:LYS:HD3	2:B:276:VAL:H	1.75	0.51
2:B:203:GLU:HA	2:B:206:ARG:HG2	1.93	0.50
1:A:405:TYR:CE2	1:A:407:GLN:HB2	2.47	0.50
1:A:454:LYS:HZ2	1:A:552:VAL:HB	1.77	0.49
2:B:46:LYS:HE2	2:B:116:PHE:HB3	1.95	0.49
2:B:308:GLU:HA	2:B:311:LYS:HE2	1.94	0.49
1:A:111:VAL:HG21	1:A:160:PHE:HZ	1.78	0.48
1:A:503:LEU:HD22	1:A:535:TRP:HB2	1.96	0.47
1:A:443:ASP:HB3	1:A:550:LYS:HD3	1.97	0.47
2:B:167:ILE:HG12	2:B:212:TRP:CE3	2.50	0.46
2:B:281:LYS:HG3	2:B:284:ARG:NH1	2.30	0.46
1:A:342:TYR:HB3	1:A:348:ASN:HA	1.98	0.46
2:B:271:TYR:HA	2:B:272:PRO:HD3	1.77	0.46
1:A:21:VAL:HB	1:A:59:PRO:HD3	1.98	0.45
2:B:168:LEU:HD13	2:B:180:ILE:HG21	1.97	0.45
2:B:84:THR:HB	2:B:154:LYS:HE2	1.99	0.44
1:A:366:LYS:HE2	1:A:405:TYR:OH	2.17	0.44
1:A:442:VAL:HB	1:A:481:ALA:HB1	2.00	0.44
2:B:315:HIS:O	2:B:347:LYS:NZ	2.49	0.44
1:A:363:ASN:HD21	1:A:366:LYS:HE3	1.83	0.44
2:B:101:LYS:HD3	2:B:382:ILE:HG23	2.00	0.44
1:A:90:VAL:HG21	2:B:141:GLY:H	1.83	0.43
2:B:163:SER:O	2:B:167:ILE:HG13	2.18	0.43
2:B:281:LYS:HG3	2:B:284:ARG:HH11	1.84	0.43
1:A:295:LEU:HD13	1:A:300:GLU:HB2	2.01	0.42
1:A:475:GLN:HB3	1:A:501:TYR:CE2	2.54	0.42
1:A:80:LEU:O	1:A:84:THR:OG1	2.26	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:513:SER:N	1:A:519:ASN:OD1	2.52	0.41
2:B:194:GLU:O	2:B:198:HIS:N	2.38	0.41
2:B:82:LYS:NZ	5:B:504:HOH:O	2.51	0.41
2:B:104:LYS:HB2	2:B:192:ASP:HA	2.02	0.41
1:A:244:ILE:H	1:A:244:ILE:HG13	1.66	0.41
1:A:540:LYS:HD3	1:A:540:LYS:HA	1.93	0.41
2:B:169:GLU:HB3	2:B:170:PRO:HD3	2.02	0.41
1:A:312:GLU:HA	1:A:313:PRO:HD3	1.85	0.41
2:B:270:ILE:HG12	2:B:346:PHE:HB3	2.03	0.41
2:B:420:PRO:HA	2:B:421:PRO:HD2	1.98	0.41
1:A:122:GLU:HA	1:A:125:ARG:HG3	2.03	0.40
1:A:142:ILE:HD12	1:A:142:ILE:H	1.85	0.40
1:A:363:ASN:ND2	1:A:366:LYS:HE3	2.36	0.40
2:B:198:HIS:O	2:B:202:ILE:HG12	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	543/557 (98%)	526 (97%)	17 (3%)	0	100 100
2	B	395/428 (92%)	381 (96%)	14 (4%)	0	100 100
All	All	938/985 (95%)	907 (97%)	31 (3%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	480/495 (97%)	474 (99%)	6 (1%)	69	80
2	B	364/390 (93%)	361 (99%)	3 (1%)	81	88
All	All	844/885 (95%)	835 (99%)	9 (1%)	73	83

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	91	GLN
1	A	92	LEU
1	A	286	THR
1	A	373	GLN
1	A	546	GLU
1	A	551	LEU
2	B	232	TYR
2	B	275	LYS
2	B	417	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	147	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	7AY	A	601	-	35,35,35	1.50	5 (14%)	49,49,49	1.48	5 (10%)
4	SO4	A	603	-	4,4,4	0.14	0	6,6,6	0.07	0
4	SO4	A	602	-	4,4,4	0.14	0	6,6,6	0.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	7AY	A	601	-	-	1/12/12/12	0/4/4/4

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	7AY	C0P-N0H	3.86	1.42	1.37
3	A	601	7AY	CBA-CBC	3.35	1.52	1.44
3	A	601	7AY	O0A-CAL	2.82	1.45	1.39
3	A	601	7AY	CAJ-CAI	-2.68	1.38	1.43
3	A	601	7AY	CAZ-CBA	2.45	1.44	1.39

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	7AY	C0E-N0H-C0K	5.49	124.33	118.54
3	A	601	7AY	C0E-N0H-C0P	-3.46	114.48	119.75
3	A	601	7AY	C0D-O0B-C05	3.17	125.44	117.69
3	A	601	7AY	CAM-CAL-CAK	-3.01	118.93	122.45
3	A	601	7AY	C0O-C0P-N0H	-2.16	120.42	122.44

There are no chirality outliers.


All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	601	7AY	C05-C04-O0A-CAL

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	547/557 (98%)	0.79	75 (13%) 3 3	36, 73, 122, 145	0
2	B	401/428 (93%)	0.78	54 (13%) 3 4	39, 65, 119, 135	0
All	All	948/985 (96%)	0.79	129 (13%) 3 3	36, 69, 120, 145	0

All (129) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	292	VAL	9.9
1	A	254	VAL	7.4
1	A	293	ILE	6.5
1	A	218	ASP	6.3
1	A	246	LEU	6.2
2	B	356	ARG	6.2
1	A	222	GLN	6.1
1	A	303	LEU	6.1
2	B	87	PHE	6.0
1	A	551	LEU	5.9
1	A	255	ASN	5.9
1	A	18	GLY	5.8
2	B	237	ASP	5.4
2	B	209	LEU	5.4
2	B	238	LYS	5.3
2	B	168	LEU	5.3
2	B	207	GLN	5.3
1	A	286	THR	5.2
2	B	241	VAL	5.1
1	A	289	LEU	5.1
1	A	283	LEU	5.0
2	B	67	ASP	4.8
2	B	202	ILE	4.8
1	A	247	PRO	4.7

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	311	LYS	4.6
1	A	288	ALA	4.6
2	B	359	GLY	4.5
1	A	552	VAL	4.4
1	A	261	VAL	4.4
1	A	294	PRO	4.4
1	A	548	VAL	4.3
1	A	24	TRP	4.2
2	B	14	PRO	4.2
2	B	357	MET	4.2
2	B	195	ILE	4.1
2	B	239	TRP	4.1
1	A	291	GLU	4.0
1	A	2	ILE	3.9
2	B	358	ARG	3.9
1	A	221	HIS	3.9
2	B	206	ARG	3.9
1	A	287	LYS	3.8
1	A	61	PHE	3.8
2	B	88	TRP	3.7
1	A	259	LYS	3.7
1	A	17	ASP	3.7
2	B	191	SER	3.7
1	A	295	LEU	3.6
1	A	245	VAL	3.5
2	B	199	ARG	3.5
2	B	193	LEU	3.5
1	A	263	LYS	3.5
1	A	251	SER	3.5
1	A	92	LEU	3.4
1	A	300	GLU	3.4
1	A	38	CYS	3.4
2	B	69	THR	3.3
2	B	196	GLY	3.3
2	B	197	GLN	3.3
1	A	297	GLU	3.2
1	A	59	PRO	3.2
2	B	284	ARG	3.2
1	A	244	ILE	3.2
2	B	236	PRO	3.2
1	A	12	LEU	3.2
1	A	84	THR	3.1

*Continued on next page...*



*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	130	PHE	3.1
1	A	74	LEU	3.1
1	A	56	TYR	3.1
2	B	301	LEU	3.0
1	A	257	ILE	3.0
1	A	193	LEU	3.0
1	A	252	TRP	3.0
1	A	21	VAL	2.9
1	A	82	LYS	2.9
2	B	200	THR	2.9
2	B	178	ILE	2.9
2	B	204	GLU	2.9
2	B	240	THR	2.9
2	B	94	ILE	2.8
1	A	279	LEU	2.8
2	B	13	LYS	2.8
2	B	232	TYR	2.8
1	A	127	TYR	2.8
2	B	425	LEU	2.8
2	B	208	HIS	2.7
1	A	543	GLY	2.7
2	B	95	PRO	2.7
1	A	91	GLN	2.7
1	A	15	GLY	2.6
1	A	77	PHE	2.6
2	B	372	VAL	2.6
1	A	16	MET	2.6
1	A	73	LYS	2.6
1	A	309	ILE	2.6
1	A	14	PRO	2.5
2	B	205	LEU	2.5
2	B	203	GLU	2.5
2	B	210	LEU	2.5
2	B	169	GLU	2.5
1	A	63	ILE	2.4
2	B	166	LYS	2.4
1	A	19	PRO	2.4
1	A	4	PRO	2.3
2	B	194	GLU	2.3
1	A	144	TYR	2.3
2	B	172	LYS	2.2
1	A	-1	MET	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	93	GLY	2.2
2	B	173	LYS	2.2
1	A	264	LEU	2.2
2	B	170	PRO	2.2
1	A	549	ASP	2.2
1	A	248	GLU	2.2
1	A	220	LYS	2.2
2	B	212	TRP	2.1
1	A	25	PRO	2.1
2	B	171	PHE	2.1
2	B	83	ARG	2.1
1	A	301	LEU	2.1
1	A	50	ILE	2.1
1	A	22	LYS	2.1
2	B	15	GLY	2.1
1	A	90	VAL	2.1
2	B	119	PRO	2.0
1	A	116	PHE	2.0
1	A	253	THR	2.0
1	A	290	THR	2.0
2	B	198	HIS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	SO4	A	602	5/5	0.86	0.15	85,85,93,99	0
4	SO4	A	603	5/5	0.93	0.23	100,101,111,112	0

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	7AY	A	601	32/32	0.94	0.18	53,59,65,69	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.