



# wwPDB NMR Structure Validation Summary Report ⓘ

May 29, 2020 – 01:58 am BST

PDB ID : 2XDF  
Title : Solution Structure of the Enzyme I Dimer Complexed with HPr Using Residual Dipolar Couplings and Small Angle X-Ray Scattering  
Authors : Schwieters, C.D.; Suh, J.-Y.; Grishaev, A.; Guirlando, R.; Takayama, Y.; Clore, G.M.  
Deposited on : 2010-04-30

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : 2.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11

# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:  
*SOLUTION NMR, SOLUTION SCATTERING*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	573	38% (green), 49% (yellow), 12% (orange)
1	B	573	37% (green), 51% (yellow), 12% (orange)
2	C	85	65% (green), 33% (yellow), .. (orange)
2	D	85	64% (green), 34% (yellow), .. (orange)

## 2 Ensemble composition and analysis

This entry contains 2 models. Identification of well-defined residues and clustering analysis are not possible.

### 3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 20502 atoms, of which 10338 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called PHOSPHOENOLPYRUVATE-PROTEIN PHOSPHO-TRANSFERASE.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	573	8956	2790	4514	757	874	21	0
1	B	573	8956	2790	4514	757	874	21	0

- Molecule 2 is a protein called PHOSPHOCARRIER PROTEIN HPR.


Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
2	C	85	1295	401	655	107	130	2	0
2	D	85	1295	401	655	107	130	2	0

## 4 Residue-property plots


### 4.1 Average score per residue in the NMR ensemble


These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: PHOSPHOENOLPYRUVATE-PROTEIN PHOSPHOTRANSFERASE



- Molecule 1: PHOSPHOENOLPYRUVATE-PROTEIN PHOSPHOTRANSFERASE





- Molecule 2: PHOSPHOCARRIER PROTEIN HPR

Chain C: 65% 33%



- Molecule 2: PHOSPHOCARRIER PROTEIN HPR

Chain D: 64% 34%





## 4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.


- Molecule 1: PHOSPHOENOLPYRUVATE-PROTEIN PHOSPHOTRANSFERASE

Chain A: 51% 39% 10%





● Molecule 1: PHOSPHOENOLPYRUVATE-PROTEIN PHOSPHOTRANSFERASE



● Molecule 2: PHOSPHOCARRIER PROTEIN HPR



● Molecule 2: PHOSPHOCARRIER PROTEIN HPR



## 5 Refinement protocol and experimental data overview [i](#)

The models were refined using the following method: *simulated annealing*.

Of the 120 calculated structures, 2 were deposited, based on the following criterion: *BEST EXPERIMENT FIT, AND THEN LOWEST ENERGY*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Xplor-NIH	refinement	2.25
Xplor-NIH	structure solution	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

### 5.1 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	4434	4510	4502	486±7
1	B	4425	4497	4489	486±5
2	C	640	655	650	38±5
2	D	640	655	650	38±4
All	All	20278	20634	20582	1948

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 48.

5 of 1730 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:126:ARG:HH22	2:C:619:ALA:CB	1.61	1.04	1	1
1:B:126:ARG:HH22	2:D:619:ALA:CB	1.55	1.04	1	1
1:B:314:PHE:CE2	1:B:376:ALA:HA	1.49	1.41	1	1
1:A:314:PHE:CE2	1:A:376:ALA:HA	1.48	1.40	1	1
1:B:562:MET:SD	1:B:562:MET:CE	1.48	2.01	2	1

## 5.2 Torsion angles [i](#)

### 5.2.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	570/573 (99%)	538±1 (94±0%)	26±1 (4±0%)	7±0 (1±0%)	17	64
1	B	569/573 (99%)	536±1 (94±0%)	26±1 (5±0%)	7±0 (1±0%)	17	64
2	C	83/85 (98%)	76±0 (92±0%)	6±0 (7±0%)	1±0 (1±0%)	17	64
2	D	83/85 (98%)	76±0 (92±0%)	6±0 (7±0%)	1±0 (1±0%)	17	64
All	All	2610/2632 (99%)	2451 (94%)	127 (5%)	32 (1%)	17	64

5 of 19 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	183	ALA	2
1	B	183	ALA	2
1	A	260	PRO	2
1	A	6	LEU	2
2	D	651	GLN	2

### 5.2.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	473/474 (100%)	409±5 (86±1%)	65±5 (14±1%)	7	47
1	B	472/474 (100%)	406±4 (86±1%)	66±4 (14±1%)	6	46
2	C	70/70 (100%)	63±2 (90±3%)	7±2 (10±3%)	11	56
2	D	70/70 (100%)	64±2 (91±2%)	7±2 (9±2%)	12	59
All	All	2170/2176 (100%)	1882 (87%)	288 (13%)	7	48

5 of 227 unique residues with a non-rotameric sidechain are listed below. They are sorted by the

frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	B	404	LYS	2
1	A	58	LYS	2
1	A	27	ASP	2
1	B	307	LEU	2
1	A	474	TYR	2

### 5.2.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.3 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.4 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.5 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.6 Other polymers [i](#)

There are no such molecules in this entry.

### 5.7 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Chemical shift validation

No chemical shift data were provided