



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 6, 2023 – 02:29 AM EST

PDB ID : 4XD6
Title : Phosphotriesterase Variant E2a
Authors : Jackson, C.J.; Campbell, E.; Kaltenbach, M.; Tokuriki, N.
Deposited on : 2014-12-19
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	333	 5% 87% 11% ..
1	G	333	 7% 86% 12% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5868 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphotriesterase variant PTE-E2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	330	Total 2585	C 1630	N 465	O 482	S 8	0	8	0
1	G	329	Total 2547	C 1609	N 453	O 477	S 8	0	5	0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total 2	Zn 2	0	0
2	G	2	Total 2	Zn 2	0	0

- Molecule 3 is CACODYLATE ION (three-letter code: CAC) (formula: C₂H₆AsO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	As	C	O		
3	A	1	5	1	2	2	0	0
3	G	1	5	1	2	2	0	0

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	8	6	2	0	0
4	A	1	8	6	2	0	0
4	G	1	8	6	2	0	0
4	G	1	8	6	2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	370	370	370	0	0
5	G	320	320	320	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphotriesterase variant PTE-E2



- Molecule 1: Phosphotriesterase variant PTE-E2



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	85.74Å 85.76Å 88.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.28 – 1.75 44.19 – 1.75	Depositor EDS
% Data completeness (in resolution range)	89.4 (39.28-1.75) 83.2 (44.19-1.75)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 1.75Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.9_1692)	Depositor
R, R_{free}	0.214 , 0.269 0.216 , 0.270	Depositor DCC
R_{free} test set	3157 reflections (5.26%)	wwPDB-VP
Wilson B-factor (Å ²)	12.9	Xtriage
Anisotropy	0.359	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 46.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.025 for -h,l,k 0.025 for -l,-k,-h 0.027 for k,h,-l 0.016 for k,l,h 0.016 for l,h,k	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5868	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CAC, KCX, ZN, MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/2620	0.55	0/3554
1	G	0.33	0/2582	0.54	0/3505
All	All	0.38	0/5202	0.55	0/7059

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2585	0	2600	28	0
1	G	2547	0	2563	28	0
2	A	2	0	0	0	0
2	G	2	0	0	0	0
3	A	5	0	0	0	0
3	G	5	0	0	0	0
4	A	16	0	28	2	0
4	G	16	0	28	2	0
5	A	370	0	0	10	1
5	G	320	0	0	6	0
All	All	5868	0	5219	55	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:343:GLN:HE22	4:G:405:MPD:H12	1.39	0.85
1:G:301:ASP:OD2	5:G:815:HOH:O	1.98	0.82
1:A:301:ASP:OD2	5:A:859:HOH:O	1.99	0.81
1:A:118[B]:ARG:NH2	5:A:844:HOH:O	2.20	0.73
1:G:181:GLU:OE2	5:G:803:HOH:O	2.06	0.73
1:A:343:GLN:HE22	4:A:404:MPD:H12	1.54	0.71
1:A:118[A]:ARG:NH1	5:A:501:HOH:O	2.27	0.67
1:G:285:LYS:NZ	1:G:289:ASP:OD2	2.32	0.62
1:G:254:HIS:NE2	5:G:815:HOH:O	2.31	0.61
1:A:89[B]:ARG:HD3	5:A:695:HOH:O	2.01	0.60
1:A:89[B]:ARG:HG2	5:A:525:HOH:O	2.03	0.59
1:A:57:HIS:HB2	1:A:303:LEU:HB3	1.90	0.54
1:G:67:ARG:NH2	1:G:111[A]:SER:OG	2.40	0.54
1:G:46[B]:ILE:HG21	1:G:359[B]:SER:OG	2.09	0.52
1:A:46:ILE:HD11	1:A:355:ALA:HB1	1.91	0.52
1:G:192:LEU:O	1:G:362:LEU:HD11	2.10	0.52
1:G:57:HIS:HB2	1:G:303:LEU:HB3	1.90	0.52
1:G:77:LYS:NZ	1:G:81:GLU:OE2	2.37	0.51
1:A:202:THR:HG21	5:A:631:HOH:O	2.10	0.51
4:A:405:MPD:O4	4:A:405:MPD:O2	2.28	0.51
1:G:343:GLN:NE2	4:G:405:MPD:H12	2.17	0.50
1:A:259:ALA:HB2	1:A:272:MET:SD	2.51	0.50
1:G:300:ASN:OD1	1:G:328:ILE:HG12	2.12	0.49
1:A:41:ARG:NH2	5:A:502:HOH:O	2.29	0.49
1:A:344:GLU:CD	1:A:344:GLU:H	2.17	0.48
1:G:54:THR:HG21	1:G:297:LEU:HB3	1.96	0.48
1:A:114:ALA:O	1:A:118[B]:ARG:HG2	2.14	0.47
1:G:167:ILE:HG13	1:G:197:PRO:HG2	1.96	0.47
1:A:149:PHE:O	1:A:152:ARG:HG3	2.15	0.47
1:A:71:GLU:H	1:A:71:GLU:CD	2.20	0.46
1:G:172:THR:HG22	1:G:180:GLN:HB3	1.97	0.45
1:G:202:THR:HG22	1:G:212:GLN:NE2	2.31	0.45
1:A:86:GLY:HA3	1:A:304:PHE:CE1	2.52	0.45
1:A:254:HIS:NE2	5:A:859:HOH:O	2.31	0.45
1:A:71:GLU:OE1	1:G:152:ARG:NE	2.41	0.45
1:G:51:PHE:CE2	1:G:329:PRO:HB3	2.52	0.45
1:A:57:HIS:O	1:A:303:LEU:HA	2.18	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:302:TRP:CH2	1:G:321:ASN:HB3	2.53	0.44
1:G:40:VAL:HA	1:G:163:ILE:HG23	2.00	0.44
1:A:333:ILE:HB	1:A:334:PRO:HD3	2.00	0.43
1:G:336:LEU:HB3	1:G:341:ILE:HD12	2.00	0.43
1:A:323:ASP:O	5:A:695:HOH:O	2.22	0.43
1:G:362:LEU:HD12	1:G:362:LEU:HA	1.79	0.42
1:A:131:TRP:CG	1:A:132:LEU:N	2.88	0.41
1:A:118[B]:ARG:HG3	5:A:602:HOH:O	2.19	0.41
1:A:249:LEU:HD13	1:A:295:GLN:HA	2.02	0.41
1:G:131:TRP:CG	1:G:132:LEU:N	2.89	0.41
1:A:308:SER:O	1:G:140:LEU:HD22	2.21	0.41
1:G:57:HIS:O	1:G:303:LEU:HA	2.19	0.41
1:A:243:LEU:HD23	1:A:243:LEU:HA	1.93	0.41
1:A:328:ILE:HD13	1:A:328:ILE:HA	1.88	0.41
1:G:175:LYS:HG3	5:G:748:HOH:O	2.21	0.41
1:G:356:ARG:HD2	5:G:561:HOH:O	2.20	0.41
1:G:315:ASP:OD1	5:G:693:HOH:O	2.21	0.40
1:A:330:LEU:O	1:A:334:PRO:HG2	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:545:HOH:O	5:A:549:HOH:O[3_555]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	335/333 (101%)	322 (96%)	12 (4%)	1 (0%)	41 22
1	G	331/333 (99%)	318 (96%)	12 (4%)	1 (0%)	41 22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	666/666 (100%)	640 (96%)	24 (4%)	2 (0%)	41 22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	174	GLY
1	A	174	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	271/266 (102%)	261 (96%)	10 (4%)	34 12
1	G	268/266 (101%)	264 (98%)	4 (2%)	65 49
All	All	539/532 (101%)	525 (97%)	14 (3%)	44 23

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	PHE
1	A	152	ARG
1	A	202	THR
1	A	264	ASP
1	A	271	PHE
1	A	293	MET
1	A	299	SER
1	A	303	LEU
1	A	344	GLU
1	A	356	ARG
1	G	51	PHE
1	G	54	THR
1	G	299	SER
1	G	303	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such

sidechains are listed below:

Mol	Chain	Res	Type
1	G	343	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	KCX	G	169	1,2	9,11,12	1.10	1 (11%)	5,12,14	2.27	2 (40%)
1	KCX	A	169	1,2	9,11,12	0.94	0	5,12,14	1.50	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	G	169	1,2	-	1/9/10/12	-
1	KCX	A	169	1,2	-	0/9/10/12	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	169	KCX	OQ1-CX	2.42	1.26	1.21

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	169	KCX	OQ1-CX-NZ	-3.63	119.33	124.96
1	G	169	KCX	CE-NZ-CX	-3.41	116.41	121.89
1	A	169	KCX	OQ1-CX-NZ	-2.69	120.79	124.96

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	G	169	KCX	C-CA-CB-CG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	MPD	A	404	-	7,7,7	0.27	0	9,10,10	0.74	0
3	CAC	G	402	2	0,4,4	-	-	0,6,6	-	-
4	MPD	G	404	-	7,7,7	0.28	0	9,10,10	0.29	0
4	MPD	A	405	-	7,7,7	0.32	0	9,10,10	0.18	0
3	CAC	A	403	2	0,4,4	-	-	0,6,6	-	-
4	MPD	G	405	-	7,7,7	0.29	0	9,10,10	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MPD	G	404	-	-	1/5/5/5	-
4	MPD	G	405	-	-	2/5/5/5	-
4	MPD	A	404	-	-	5/5/5/5	-
4	MPD	A	405	-	-	1/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	404	MPD	C1-C2-C3-C4
4	A	404	MPD	C2-C3-C4-C5
4	G	404	MPD	C2-C3-C4-C5
4	G	405	MPD	C2-C3-C4-C5
4	A	404	MPD	CM-C2-C3-C4
4	A	405	MPD	C1-C2-C3-C4
4	A	404	MPD	O2-C2-C3-C4
4	G	405	MPD	O2-C2-C3-C4
4	A	404	MPD	C2-C3-C4-O4

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	404	MPD	1	0
4	A	405	MPD	1	0
4	G	405	MPD	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	329/333 (98%)	0.62	17 (5%) 27 33	5, 14, 50, 138	0
1	G	328/333 (98%)	0.57	24 (7%) 15 20	9, 22, 56, 95	0
All	All	657/666 (98%)	0.59	41 (6%) 20 26	5, 18, 56, 138	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	273	GLY	24.6
1	A	260	ILE	24.4
1	A	262	LEU	19.0
1	A	266	ALA	18.2
1	A	265	ASN	17.6
1	A	271	PHE	14.7
1	A	261	GLY	13.8
1	A	270	ALA	13.6
1	A	268	ALA	11.0
1	A	269	SER	10.9
1	A	264	ASP	8.4
1	G	273	GLY	7.4
1	G	274	ILE	7.2
1	A	267	SER	7.2
1	G	260	ILE	6.8
1	G	265	ASN	6.3
1	G	271	PHE	6.2
1	G	263	GLU	5.5
1	G	261	GLY	5.4
1	A	263	GLU	5.4
1	G	270	ALA	5.4
1	G	266	ALA	4.9
1	A	259	ALA	4.9
1	G	262	LEU	4.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	264	ASP	4.6
1	G	362	LEU	4.5
1	A	272	MET	4.5
1	A	274	ILE	4.4
1	G	269	SER	4.0
1	G	268	ALA	3.9
1	A	34	GLY	3.0
1	G	267	SER	3.0
1	G	275	ARG	2.6
1	G	259	ALA	2.5
1	G	335	PHE	2.5
1	G	204	ALA	2.4
1	G	272	MET	2.4
1	G	35	ASP	2.3
1	G	276	SER	2.2
1	G	37	ILE	2.1
1	G	258	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	G	169	12/13	0.94	0.10	11,16,18,20	0
1	KCX	A	169	12/13	0.95	0.10	3,7,9,11	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	MPD	A	405	8/8	0.75	0.21	18,41,44,48	0
4	MPD	G	404	8/8	0.76	0.24	25,39,47,50	0
4	MPD	G	405	8/8	0.81	0.26	34,46,49,49	0
4	MPD	A	404	8/8	0.83	0.18	23,33,42,50	0
3	CAC	A	403	5/5	0.90	0.20	15,33,56,79	0
3	CAC	G	402	5/5	0.90	0.26	19,28,63,72	0
2	ZN	G	401	1/1	0.99	0.04	22,22,22,22	0
2	ZN	G	403	1/1	0.99	0.05	18,18,18,18	0
2	ZN	A	401	1/1	1.00	0.06	8,8,8,8	0
2	ZN	A	402	1/1	1.00	0.05	12,12,12,12	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.