

Full wwPDB X-ray Structure Validation Report (i)

Aug 15, 2023 – 10:42 PM EDT

PDB ID : 1XJD

Title : Crystal Structure of PKC-theta complexed with Staurosporine at 2A resolution

Authors : Xu, Z.B. Deposited on : 2004-09-23

Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED

EDS : NOT EXECUTED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution	
Metric	$(\# \mathrm{Entries})$	(#Entries, resolution range(Å)	
Clashscore	141614	9178 (2.00-2.00)	
Ramachandran outliers	138981	9054 (2.00-2.00)	
Sidechain outliers	138945	9053 (2.00-2.00)	

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain					
1	A	345	63%	15%		18%		



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2502 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


• Molecule 1 is a protein called Protein kinase C, theta type.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace			
1	A	282	Total 2352	C 1525	N 388	O 421	P 2	S 16	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	538	TPO	THR	modified residue	UNP Q04759
A	695	SEP	SER	modified residue	UNP Q04759

• Molecule 2 is STAUROSPORINE (three-letter code: STU) (formula: C₂₈H₂₆N₄O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	С	N	0	0	0
			35	28	4	3		

• Molecule 3 is water.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	115	Total O 115 115	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Protein kinase C, theta type





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	139.57Å 42.40Å 67.68Å	Depositor
a, b, c, α , β , γ	90.00° 116.24° 90.00°	Depositor
Resolution (Å)	19.93 - 2.00	Depositor
% Data completeness	83.2 (19.93-2.00)	Depositor
(in resolution range)	09.2 (13.35-2.00)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	0.04	Depositor
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.201 , 0.216	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2502	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: STU, SEP, TPO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	$\mathbf{lengths}$	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.41	0/2390	0.64	0/3210	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2352	0	2304	62	1
2	A	35	0	26	1	0
3	A	115	0	0	4	1
All	All	2502	0	2330	62	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1			$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:A:385:MET:CE	1:A:393:LYS:HD2	2.04	0.88



 $Continued\ from\ previous\ page...$

Continued from prev		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ (\mathring{\rm A})$	overlap (Å)
1:A:417:LEU:HD21	1:A:690:MET:HG3	1.59	0.84
1:A:385:MET:HE3	1:A:393:LYS:HD2	1.59	0.84
1:A:603:GLU:HG3	3:A:103:HOH:O	1.82	0.78
1:A:447:THR:HG22	1:A:696:PHE:N	1.98	0.77
1:A:392:GLY:HA3	1:A:411:LEU:HD23	1.67	0.75
1:A:450:THR:CG2	1:A:451:LYS:N	2.52	0.72
1:A:447:THR:N	1:A:696:PHE:O	2.22	0.72
1:A:392:GLY:CA	1:A:411:LEU:HD23	2.19	0.72
1:A:450:THR:HG22	1:A:451:LYS:N	2.07	0.70
1:A:696:PHE:C	1:A:696:PHE:CD1	2.69	0.65
1:A:450:THR:CG2	1:A:451:LYS:H	2.08	0.65
1:A:417:LEU:CD2	1:A:690:MET:HG3	2.25	0.65
1:A:391:PHE:HZ	1:A:425:THR:HA	1.62	0.64
1:A:391:PHE:CZ	1:A:425:THR:HA	2.33	0.64
1:A:450:THR:HG22	1:A:452:GLU:H	1.65	0.61
1:A:391:PHE:C	1:A:411:LEU:HD21	2.24	0.58
1:A:438:HIS:O	1:A:443:HIS:HE1	1.88	0.57
1:A:450:THR:HG23	1:A:451:LYS:H	1.70	0.56
1:A:391:PHE:O	1:A:409:LYS:HE3	2.06	0.55
1:A:499:GLY:HA3	1:A:531:LEU:HD21	1.89	0.54
1:A:454:LEU:HD22	1:A:694:PHE:CD2	2.43	0.54
1:A:447:THR:HG23	1:A:695:SEP:HA	1.89	0.54
1:A:411:LEU:HD12	1:A:456:PHE:HE1	1.72	0.53
1:A:423:GLU:O	1:A:427:VAL:HG23	2.08	0.53
1:A:389:GLY:O	1:A:390:SER:C	2.46	0.53
1:A:417:LEU:HG	1:A:691:PHE:HZ	1.74	0.53
1:A:513:ASP:HB3	1:A:515:ASP:OD1	2.09	0.52
1:A:402:THR:O	1:A:403:ASN:HB2	2.10	0.51
1:A:447:THR:HG22	1:A:696:PHE:HB3	1.92	0.51
1:A:605:GLU:HG2	3:A:103:HOH:O	2.09	0.51
1:A:391:PHE:O	1:A:411:LEU:HD21	2.11	0.51
1:A:392:GLY:HA3	1:A:411:LEU:CD2	2.40	0.50
1:A:391:PHE:O	1:A:409:LYS:CE	2.59	0.50
1:A:385:MET:HE3	1:A:393:LYS:CD	2.38	0.50
1:A:467:MET:O	1:A:471:GLN:HG3	2.12	0.49
1:A:696:PHE:O	1:A:696:PHE:HD1	1.96	0.48
1:A:605:GLU:HG3	1:A:635:ARG:NH2	2.29	0.48
1:A:447:THR:CG2	1:A:694:PHE:O	2.63	0.47
1:A:447:THR:HG21	1:A:694:PHE:O	2.14	0.47
1:A:438:HIS:CD2	1:A:440:PHE:H	2.32	0.47
1:A:496:HIS:CE1	1:A:558:HIS:HB2	2.50	0.46



Continued from previous page...

A + 1	A 4 0	Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \ (\mathring{\rm A})$	overlap (Å)
1:A:492:LEU:HD11	1:A:505:LEU:HD22	1.98	0.46
1:A:390:SER:HB2	1:A:522:ASP:OD1	2.16	0.45
1:A:391:PHE:C	1:A:411:LEU:CD2	2.84	0.45
1:A:392:GLY:N	1:A:411:LEU:HD23	2.32	0.45
1:A:617:GLU:OE1	1:A:619:GLU:HG3	2.17	0.45
1:A:607:LYS:O	1:A:611:VAL:HG23	2.17	0.44
1:A:450:THR:HG22	1:A:452:GLU:N	2.31	0.44
1:A:512:LEU:HD23	1:A:518:ILE:HG22	1.99	0.44
1:A:558:HIS:HD2	1:A:562:TRP:HE1	1.66	0.44
1:A:409:LYS:NZ	1:A:428:GLU:OE1	2.51	0.43
1:A:389:GLY:HA3	2:A:200:STU:H16	2.02	0.41
1:A:502:TYR:HE2	1:A:505:LEU:HA	1.85	0.41
1:A:499:GLY:CA	1:A:531:LEU:HD21	2.51	0.41
1:A:645:LYS:HA	3:A:34:HOH:O	2.20	0.41
1:A:454:LEU:HD22	1:A:694:PHE:CE2	2.55	0.41
1:A:600:ARG:HD3	3:A:42:HOH:O	2.20	0.41
1:A:638:ASN:CG	1:A:641:GLU:HG3	2.40	0.41
1:A:473:CYS:O	1:A:474:HIS:HB2	2.20	0.41
1:A:529:ASN:HD22	1:A:529:ASN:HA	1.57	0.41
1:A:692:ARG:O	1:A:693:ASN:HB2	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
3:A:114:HOH:O	3:A:114:HOH:O[2_656]	1.77	0.43
1:A:477:ASP:OD1	1:A:477:ASP:OD1[2_656]	2.18	0.02

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	276/345 (80%)	262 (95%)	12 (4%)	2 (1%)	22 16	

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	690	MET
1	A	390	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	252/312 (81%)	240 (95%)	12 (5%)	25 22	

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	390	SER
1	A	391	PHE
1	A	447	THR
1	A	450	THR
1	A	529	ASN
1	A	600	ARG
1	A	604	LYS
1	A	688	GLN
1	A	691	PHE
1	A	692	ARG
1	A	693	ASN
1	A	696	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	404	GLN
1	A	438	HIS



Continued from previous page...

Mol	Chain	Res	Type
1	A	443	HIS
1	A	529	ASN
1	A	554	GLN
1	A	558	HIS
1	A	582	GLN
1	A	595	ASN
1	A	630	GLN
1	A	688	GLN
1	A	693	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trme	Chain	Res	Link	В	ond leng	${ m gths}$	В	ond ang	les
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	SEP	A	695	1	8,9,10	2.27	1 (12%)	8,12,14	3.39	2 (25%)
1	TPO	A	538	1	8,10,11	3.06	1 (12%)	10,14,16	1.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	695	1	-	4/5/8/10	-
1	TPO	A	538	1	-	0/9/11/13	-

All (2) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
1	A	538	TPO	P-OG1	-8.42	1.43	1.59
1	A	695	SEP	P-OG	-6.16	1.40	1.60

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
1	A	695	SEP	O2P-P-OG	-6.49	89.47	106.73
1	A	695	SEP	O3P-P-OG	6.17	123.14	106.73

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	695	SEP	N-CA-CB-OG
1	A	695	SEP	CB-OG-P-O1P
1	A	695	SEP	CB-OG-P-O2P
1	A	695	SEP	CB-OG-P-O3P

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	695	SEP	1	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
IVIOI	Туре	Chain	rtes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	STU	A	200	-	30,42,42	4.37	23 (76%)	31,68,68	1.78	8 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	STU	A	200	-	-	0/4/42/42	-

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	Ideal(Å)
2	A	200	STU	C7-C6	8.82	1.57	1.43
2	A	200	STU	C11-C18	7.39	1.52	1.42
2	A	200	STU	C6-C19	6.29	1.50	1.42
2	A	200	STU	C12-C17	6.07	1.51	1.41
2	A	200	STU	C24-C23	6.07	1.62	1.53
2	A	200	STU	C5-C20	6.02	1.51	1.41
2	A	200	STU	C24-C25	5.32	1.62	1.51
2	A	200	STU	O6-C22	5.11	1.51	1.42
2	A	200	STU	C14-C15	4.87	1.50	1.38
2	A	200	STU	C10-C11	4.64	1.50	1.42
2	A	200	STU	C1-C20	4.62	1.50	1.41
2	A	200	STU	C3-C2	4.40	1.49	1.38
2	A	200	STU	C2-C1	4.35	1.46	1.36
2	A	200	STU	C8-N1	-4.27	1.31	1.35
2	A	200	STU	C14-C13	4.03	1.45	1.36
2	A	200	STU	C22-C23	3.90	1.56	1.52
2	A	200	STU	C16-C17	3.88	1.48	1.41
2	A	200	STU	C3-C4	3.59	1.44	1.36
2	A	200	STU	C4-C5	3.35	1.47	1.41
2	A	200	STU	C13-C12	3.32	1.47	1.41
2	A	200	STU	C15-C16	3.00	1.43	1.36
2	A	200	STU	C26-C21	2.67	1.55	1.51
2	A	200	STU	C28-N4	2.46	1.53	1.46

All (8) bond angle outliers are listed below:

\mathbf{Mol}	Chain	Res	Type	${f Atoms}$	\mathbf{Z}	$\operatorname{Observed}(^{o})$	$\mathbf{Ideal}(^o)$
2	A	200	STU	C9-N1-C8	4.55	118.23	113.85
2	A	200	STU	C26-C21-C22	-3.38	106.06	112.64



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
2	A	200	STU	C7-C8-N1	-2.99	103.35	106.37
2	A	200	STU	C11-C12-C17	2.95	109.60	106.37
2	A	200	STU	C24-C23-N4	-2.56	106.48	112.17
2	A	200	STU	C10-C9-N1	-2.56	99.16	101.76
2	A	200	STU	O4-C25-C24	-2.47	108.75	112.31
2	A	200	STU	C1-C20-N3	-2.34	129.45	132.25

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	200	STU	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

